



УДК / UDC 630  
УДК / UDC 635.9  
УДК / UDC 674

Online ISSN 1857-9507  
[www.sf.ukim.edu.mk/sumarski\\_pregled.htm](http://www.sf.ukim.edu.mk/sumarski_pregled.htm)

# ШУМАРСКИ ПРЕГЛЕД FOREST REVIEW

МЕЃУНАРОДНО НАУЧНО СПИСАНИЕ  
INTERNATIONAL SCIENTIFIC JOURNAL

80 ГОДИНИ ОД РАЃАЊЕТО НА РАДОСЛАВ РИЗОВСКИ  
YEARS OF RADOŠLAV RIZOVSKI'S BIRTH



Шум. преглед (Šum. pregled)  
For. review

Год. 46  
Vol. 46

Стр. 1-95  
Pag. 1-95

Скопје, 2015  
Skopje, 2015







УНИВЕРЗИТЕТ „СВ. КИРИЛ И МЕТОДИЈ“ ВО СКОПЈЕ  
Ss. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE

ШУМАРСКИ ФАКУЛТЕТ ВО СКОПЈЕ  
FACULTY OF FORESTRY IN SKOPJE



УДК / UDC 630  
УДК / UDC 635.9  
УДК / UDC 674

Online ISSN 1857-9507  
[www.sf.ukim.edu.mk/sumarski\\_pregled.htm](http://www.sf.ukim.edu.mk/sumarski_pregled.htm)

## ШУМАРСКИ ПРЕГЛЕД FOREST REVIEW

МЕЃУНАРОДНО НАУЧНО СПИСАНИЕ  
INTERNATIONAL SCIENTIFIC JOURNAL

Шум. преглед (Šum. pregled)  
For. review

Год. 46  
Vol. 46

Стр. 1-95  
Pag. 1-95

Скопје, 2015  
Skopje, 2015

<b>ШУМАРСКИ ПРЕГЛЕД</b>	<b>FOREST REVIEW</b>
Меѓународно научно списание	International Scientific Journal
Год. 46 / Стр. 1-95	Vol. 46 / Pag. 1-95
Скопје, 2015	Skopje, 2015
Online ISSN 1857-9507	Online ISSN 1857-9507
УДК 630	UDC 630
УДК 635.9	UDC 635.9
УДК 674	UDC 674
<b>Издавач</b>	<b>Publisher</b>
Универзитет „Св. Кирил и Методиј“ во Скопје	Ss. Cyril and Methodius University in Skopje
Шумарски факултет во Скопје	Faculty of Forestry in Skopje
Декан	Dean
Д-р Јане Ацевски	Jane Acevski PhD
<b>Главен и одговорен уредник</b>	<b>Editor-in-chief</b>
Д-р Бојан Симовски	Bojan Simovski PhD
<b>Уредувачки одбор</b>	<b>Editorial board</b>
Д-р Љупчо Несторовски (Скопје, Македонија)	Ljupčo Nestorovski PhD (Skopje, Macedonia)
Д-р Марилена Идојтиќ (Загреб, Хрватска)	Marilena Idžojić PhD (Zagreb, Croatia)
Д-р Милосав Анђелиќ (Подгорица, Црна Гора)	Milosav Anđelić PhD (Podgorica, Montenegro)
Д-р Милорад Даниловиќ (Белград, Србија)	Milorad Danilović PhD (Belgrade, Serbia)
Д-р Ирена Папазова-Анакиева (Скопје, Македонија)	Irena Papazova-Anakieva PhD (Skopje, Macedonia)
Д-р Роберт Брус (Љубљана, Словенија)	Robert Brus PhD (Ljubljana, Slovenia)
Д-р Чиприан Палагиану (Сучава, Романија)	Ciprian Palaghianu PhD (Suceava, Romania)
Д-р Влатко Андоновски (Скопје, Македонија)	Vlatko Andonovski PhD (Skopje, Macedonia)
Д-р Саша Орловиќ (Нови Сад, Србија)	Saša Orlović (Novi Sad, Serbia)
Д-р Маргарита Георгиева (Софија, Бугарија)	Margarita Georgieva PhD (Sofia, Bulgaria)
Д-р Зоран Говедар (Бања Лука, Р. Српска, БИХ)	Zoran Govedar PhD (Banja Luka, R. Srpska, BiH)
Д-р Јасминка Ризовска Атанасовска (Скопје, Македонија)	Jasminka Rizovska Atanasovska PhD (Skopje, Macedonia)
Д-р Мариус Димитров (Софија, Бугарија)	Marius Dimitrov PhD (Sofia, Bulgaria)
М-р Дејан Манџуковски (Скопје, Македонија)	Dejan Mandžukovski MSc (Skopje, Macedonia)
<b>Технички уредник</b>	<b>Technical editor</b>
М-р Дејан Манџуковски	Dejan Mandžukovski MSc
<b>Корица и насловна фотографија</b>	<b>Cover page and photography</b>
Д-р Бојан Симовски, <i>Ramonda nathaliae</i>	Bojan Simovski PhD, <i>Ramonda nathaliae</i>
Излегува еднаш годишно	Published once a year
<b>Интернет-страница</b>	<b>Web page (on-line)</b>
<a href="http://www.sf.ukim.edu.mk/sumarski_pregled.htm">www.sf.ukim.edu.mk/sumarski_pregled.htm</a>	<a href="http://www.sf.ukim.edu.mk/sumarski_pregled.htm">www.sf.ukim.edu.mk/sumarski_pregled.htm</a>
<b>Адреса на издавачот</b>	<b>Publisher's address</b>
УКИМ-Шумарски факултет во Скопје	UKiM Faculty of Forestry in Skopje
Редакција на Шумарски преглед	Editorial Board of the Forest Review
Ул. „16 Македонска бригада“ бр. 1	Ul. 16 Makedonska brigada br. 1
(П. факс 235)	(P.O. box 235)
1 000 Скопје	MK-1000 Skopje
Република Македонија	Republic of Macedonia
Е-пошта: <a href="mailto:sumpregled@sf.ukim.edu.mk">sumpregled@sf.ukim.edu.mk</a>	E-mail: <a href="mailto:sumpregled@sf.ukim.edu.mk">sumpregled@sf.ukim.edu.mk</a>
<a href="http://www.sf.ukim.edu.mk">www.sf.ukim.edu.mk</a>	<a href="http://www.sf.ukim.edu.mk">www.sf.ukim.edu.mk</a>

<b>ШУМАРСКИ ПРЕГЛЕД</b>	<b>FOREST REVIEW</b>
Меѓународно научно списание	International Scientific Journal
Год. 46 / Стр. 1-95	Vol. 46 / Pag. 1-95
Скопје, 2015	Skopje, 2015

Online ISSN 1857-9507	Online ISSN 1857-9507
УДК 630	UDC 630
УДК 635.9	UDC 635.9
УДК 674	UDC 674

<b>Научен и рецензентски одбор</b>	<b>Scientific and reviewers board</b>
Акад. Владо Матевски (Скопје, Македонија)	Vlado Matevski Acad. (Skopje, Macedonia)
Д-р Андраж Чарни (Љубљана, Словенија)	Andraž Čarni PhD (Ljubljana, Slovenia)
Д-р Дмитро Јакушенко (Зелена Гора, Полска)	Dmytro Iakushenko PhD (Zielona Góra, Poland)
Д-р Маријана Новаковиќ-Вуковиќ (Белград, Србија)	Marijana Novaković-Vuković PhD (Belgrade, Serbia)
Д-р Јане Ацевски (Скопје, Македонија)	Jane Acevski PhD (Skopje, Macedonia)
Д-р Југослав Брујиќ (Бања Лука, Р. Српска, БИХ)	Jugoslav Brujić PhD (Banja Luka, R. Srpska, BiH)
Д-р Марјан Никетиќ (Белград, Србија)	Marjan Niketić PhD (Belgrade, Serbia)
Д-р Рената Ќуштеревска (Скопје, Македонија)	Renata Ćušterevska PhD (Skopje, Macedonia)
Д-р Сеад Војниковиќ (Сараево, БИХ)	Sead Vojniković PhD (Sarajevo, BiH)
Д-р Бојан Симовски (Скопје, Македонија)	Bojan Simovski PhD (Skopje, Macedonia)
Д-р Јасница Медак (Загреб, Хрватска)	Jasnica Medak PhD (Zagreb, Croatia)
Д-р Митко Костадиновски (Скопје, Македонија)	Mitko Kostadinovski PhD (Skopje, Macedonia)

<b>Интернет-страница</b>	<b>Web page (on-line)</b>
<a href="http://www.sf.ukim.edu.mk/sumarski_pregled.htm">www.sf.ukim.edu.mk/sumarski_pregled.htm</a>	<a href="http://www.sf.ukim.edu.mk/sumarski_pregled.htm">www.sf.ukim.edu.mk/sumarski_pregled.htm</a>

<b>Адреса на издавачот</b>	<b>Publisher's address</b>
УКИМ-Шумарски факултет во Скопје	UKiM Faculty of Forestry in Skopje
Редакција на Шумарски преглед	Editorial Board of the Forest Review
Ул. „16 Македонска бригада“ бр. 1	Ul. 16 Makedonska brigada br. 1
(П. фах 235)	(P.O. box 235)
1 000 Скопје	MK-1000 Skopje
Република Македонија	Republic of Macedonia
Е-пошта: <a href="mailto:sumpregled@sf.ukim.edu.mk">sumpregled@sf.ukim.edu.mk</a>	E-mail: <a href="mailto:sumpregled@sf.ukim.edu.mk">sumpregled@sf.ukim.edu.mk</a>
<a href="http://www.sf.ukim.edu.mk">www.sf.ukim.edu.mk</a>	<a href="http://www.sf.ukim.edu.mk">www.sf.ukim.edu.mk</a>



## CONTENTS

### *Preface*

---

### ***Bibliography (monograph articles):***

---

Mandžukovski D. <b>THE LIFE AND WORK OF PROF. DR. RADOSLAV RIZOVSKI (16.IX 1935 – 20.VII 2008) 80 YEARS SINCE THE BIRTH OF AND OVER 40 YEARS OF EDUCATIONAL AND SCIENTIFIC ACTIVITY</b>	1
--	---

---

### ***Original Scientific Articles:***

---

Baričević D., Vukelić J., Puača M., Šapić I. <b>A PHYTOCOENOLOGICAL STUDY OF FORESTS OF HUNGARIAN OAK AND TURKEY OAK (QUERCETUM FRINETTO-CERRIDIS /RUDSKI 1949/ TRINAJSTIĆ ET AL. 1996) ON THE NORTHWEST BORDER OF ARRIVAL (NATURE PARK PAPUK, CROATIA)</b>	7
Dimitrov M., Natcheva R., Ganeva A., Gyurova D. <b>PLANT BIODIVERSITY OF SPHAGNUM-DOMINATED MIRES IN VITOSHA NATURE PARK</b>	15
Kutnar L., Eler K. <b>PLANT SPECIES DIVERSITY AND INVASIBILITY OF (PERI-)URBAN FORESTS OF LJUBLJANA, SLOVENIA</b>	30
Melovski Lj., Hristovski S. <b>FIRST RECORDS FOR SEVEN SPECIES AND ONE HYBRID FOR THE FLORA OF THE REPUBLIC OF MACEDONIA</b>	36
Nikolov I., Dimitrov M. <b>BOREO-MONTANE FOREST PHYTOCOENOSES IN CENTRAL STARA PLANINA MTS.</b>	43
Novaković-Vuković M., Milošević R. <b>ANALYSIS OF FLORISTIC COMPOSITION OF MOUNTAIN BEECH FOREST ON LIMESTONE AND SERPENTINE IN SERBIA</b>	55
Pedashenko H., Vassilev K., Bancheva S., Delcheva M., Vladimirov V. <b>FLORISTIC AND VEGETATION DIVERSITY IN KONGURA RESERVE (SOUTH-WEST BULGARIA)</b>	59
Teofilovski A., Nikolov Z., Mandžukovski D. <b>JUNCUS FILIFORMIS L. (JUNCACEAE), A NEW SPECIES IN THE FLORA OF MACEDONIA</b>	71
Vassilev K., Gavrilova A. <b>FLORA, HABITATS AND VEGETATION OF CHAMDZHA MANAGED RESERVE, CENTRAL BALKAN RANGE</b>	74
Vukelić J., Šapić I., Baričević D. <b>FLORISTIC-VEGETATIONAL VARIABILITY OF THE ASSOCIATION EPIMEDIO- CARPINETUM BETULI (HORVAT 1938) BORHIDI 1963 IN THE NORTH OF CROATIA</b>	88

---

### *Instructions to Authors*

---

## PREFACE

*Dear Colleagues and Readers,*

*It is a great pleasure to announce the online publication of the 46<sup>th</sup> volume of Forest Review.*

*As we mentioned in our past 46<sup>th</sup> volume, this 2015 is a special year because we celebrate one great jubilee: 80 years of Prof. Dr. Radoslav Rizovski's birth (1935-2015).*

*We are especially interested in your submitted articles concerning vegetation, flora and dendrology, particularly because these scientific disciplines were one of the favourable and most investigated fields of Prof. Dr. Rizovski's professional life (1935-2008). You make us proud for your interest to participate, and we are sincerely grateful for the respect you have shown. Thank you!*

*So far, we had an excellent collaboration, with the faculties of forestry from the Balkan and South-Eastern European countries. Next year we plan to double the efforts and start to publish both online and in hardcopy two issues per year. We really hope that you will continue with your hard work and our Review will find again a place for your research articles.*

*Special thanks to all authors and members of the Forest Review, as well as to all peer – reviewers for the participation in this dedicated volume.*

*On behalf of the Editorial Board,*



*Asst. Prof. Bojan Simovski PhD, Editor-in-Chief*



*Eighty years of Prof. Dr. Radoslav Rizovski's birth (1935-2015)...*



*... Many years memories of his scientific research and academic teaching*

**THE LIFE AND WORK OF PROF. DR. RADOŠLAV RIZOVSKI (16.IX 1935 – 20.VII 2008)  
80 YEARS SINCE THE BIRTH OF AND OVER 40 YEARS OF EDUCATIONAL AND SCIENTIFIC  
ACTIVITY**

MANDŽUKOVSKI D.

*Public enterprise Makedonski šumi, Skopje, Macedonia*

*Corresponding author e-mail address: d\_mandzukovski@yahoo.com*

**ABSTRACT:** Professor Radoslav Rizovski is one of the remarkable figures who made a mark in vegetation science in Macedonia. As a tireless researcher, he created invaluable tuition that will represent basis for future generations to learn from it but also to upgrade it. He has been active in nature conservation on the ground of the former Yugoslavia, especially in Macedonia. He has published 50 scientific papers, several working papers relating to vegetation mapping of part of Macedonia, and several monographs. This year marks 80 years since his birth and 40 years of teaching and research, and for that purpose this paper is dedicated to.

**Keywords:** Radoslav Rizovski, vegetation science, nature protection.

# 1 PROFESSIONAL PORTRAIT OF PROF. DR. RADOŠLAV RIZOVSKI

Professor Radoslav Rizovski was born on 16.IX 1935, in the village Rožden in Mariovo, part of a broader area of Municipality of Kavadarci. His father's roots are from Kavadarci, and from the mother's side, from Veles. His father finished studies in theology in Bitola and as highly valued man; he was a priest in three villages.

On his mother's side there were also scholars, among them teachers and clergymen.

At a young age (seven years old), he is left without a parent. After the death of his father, Rizovski with his closest family moves to Skopje, where he forms his primary and secondary education.

Persuaded by the forestry engineer Strashimir Lazarov (his first cousin), he enrolled at the Agricultural - Forestry Faculty in Skopje, on the Department of Forestry, where he undergraduated in 1958, and graduated in 1959.



**Figure 1:** Radoslav Rizovski (Rožden, 16.IX 1935 - 20.VII 2008, Bitola)

During this period and after graduation, he worked at the Institute for Forest Management - Skopje, where participates in the execution of the field works for the current year at Tuinska forest - Kicevo and field work at the Shar mountain until late autumn, where he was witnessing a difficult situation of major snow drifts at the end of field work, and complex withdrawal from the mountain.

For this short stay, the young, industrious engineer Rade, as the older colleagues from the Institute of Forest Management, proudly and respectfully called him. A short time period, Prof. Rizovski works at Forest management officec Karadzica in Skopje.

His postgraduate studies continued at the Faculty of Forestry in Zagreb, Republic of Croatia, which were realized under inter-Republic cooperation in the former Yugoslavia, for additional education of the young staff of the newly established Faculties of Forestry.

For this occasion, he stays in Zagreb for about 4 years, and under the leadership of Professor Milan Anic, prepares his master's thesis. His Master thesis was in 1969, on the topic titled: *Cenoze hrasta kitnjaka s običnim grabom i hrasta kitnjaka u Centralnoj Makedoniji i Centralnoj Hrvatskoj*- in English "Communities of sessile oak with hornbeam and sessile oak in Central Macedonia and Central Croatia".

The Doctoral dissertation entitled: *Шумите на дабот плоскач (Quercus farnetto Ten) во јужните краишта на СР Македонија*- in English "The forests of Hungarian oak (Quercus farnetto Ten) in the southern parts of the SR of Macedonia", he successfully defended 1973 in Skopje before the commission consisting of: Em, H., Fukarek, P. and Dzekov, S.

Professor Rizovski was a recognized expert in the field of floristry and phytocoenology in our state and abroad, in the broader area of the Balkan Peninsula. Published about fifty scientific and professional papers in the field of which he worked, from which several are international.

He participated in scientific committees of international congresses, editorials of scientific - professional journals and reviews of high school and university textbooks. He was also a member of Committees for several doctoral and master thesis papers of candidates, who are now leading experts in the field of phytocoenology on the territory of the former Yugoslavia.





**Figure 2:** Prof. Dr. Radoslav Rizovski with part of his family

## 2 EDUCATIONAL ACTIVITIES

During his studies, because of his interest in common names of plants and general plant affinity he was spotted by Professor Hans Em.

After the work at the Institute for Forest Management and forest economy office Karadzica, he is drafted for military service.

During his military service period, the Forestry department of then Agricultural-Forestry Faculty in Skopje opens admission for an assistant in the Department of Sylviculture on the subject of Phytocoenology. In his absence, the request for admission and other documentation, according to Professor Rizovski, was filled out and personally applied by Professor Em.

From 13.X 1961, becomes assistant of the course Phytocoenology. After his doctorate in 1973, becomes an assistant professor. From 1978, he is associate professor of the course Ecology and typology of the forests and pastures, pretty later from Phytocoenology, while for full professor was elected in 1983.

For the purposes of education in the subjects which he held on the Faculty of Forestry in Skopje, regularly visited with his students, several mountain ranges, among them focus was given to the western part of Macedonia, with emphasis on the Shar and Bistra mountains, because of the rich diversity that these mountains have.

Part of the field activities were undertaken in Katlanovo near Skopje, due to an introduction of the sub Mediterranean vegetation. These were his basic field - teaching facilities, and occasionally field trips were made to Golak - Delcevo, Jakupica and other parts of Macedonia.

Besides the courses of Phytocoenology, Professor Rizovski short period lectured Petrographics, and from the early nineties of the last century to retirement, held lectures on the course Parks with protection of the environment.

His approach towards students was genuine and uncommon. His lectures were good representation of his background knowledge from multiple fields of natural sciences, filled with evoking memories of former field works, of course, associated with the current theme that he lectured. As such, the lectures remain in the memories of a large number of students.

He was always accessible for consultation or for a general chat.

## 3 SCIENTIFIC RESEARCH ACTIVITY

The main area of science research for Professor Rizovski was the forest vegetation of Republic of Macedonia; however, he gives his contribution in research of the pastures, as well as the typology of the forests.

With the research made in the doctoral dissertation, Professor Rizovski determined that in the southern edges of Macedonia, on the move of from Demir Kapija canyon to the south, which is known as Dolno Povardarie, are developing two forest communities of Hungarian oak (*Q. farneto*), but without the presence of Turkey oak (*Q. cerris*): *Carici cuspidatae-Quercetum farneto* Rizovski and *Carpino orientalis - Quercetum farneto* Rizovski.

An important moment of his career is the research of the wild fruit dendroflora, where in cooperation with several authors worked on the genera such as: *Rubus*, *Cornus*, *Vitis*, *Prunus*, *Corylus* etc.

Important is the contribution of Professor Rizovski in the typology of the forests. From the example of the typological schools from the former Yugoslavia and Austria in 1978, built a concept that is most suited to the natural conditions in our country and the level of exploration of the vegetation. For this purpose, he made several researches and mapping of the parts of the mountains Platchkovitza and Kozuf at the site called Duditza, gathering satisfactory results.

Professor Rizovski in middle of the seventies of the last century to the beginnings of the eighties (1981) with Lj. Micevski, works on the phytosociological and ecological regionalization and typization of bare lands in order to define and classify the sites within the limits of previously established areas for afforestation. Parts of these results are published at a later date (1990).

During this period, he publishes two new communities for science. The first, on the beech (*F. moesiaca*) and small-leaved lime (*T. cordata*) and the second is community of Hop Hornbeam (*O. carpinifolia*) that develops on silicate substrate at the gorges in Macedonia.

How big was the love of the professor Rizovski towards floristic and vegetation conveys the information that he with his own personal funding purchased professional literature, herbarium materials and equipment for the research of vegetation.

An important period from the scientific work of Professor Rizovski is his work in the project called "Vegetation map of Yugoslavia, in the region of the Republic of Macedonia". This project was undertaken on the initiative of Forestry Institute from Zagreb - the Republic of Croatia, upon the before held international symposium in Stolzenau 1959, for the preparation of vegetation map of Europe. The same project on the territory of the former Yugoslavia started to be realized in 1963.

In the period of 1963 - 1987, Professor Rizovski was included as associate in the teams for mapping the vegetation of Macedonia in favor of preparing the vegetation map of Yugoslavia in scale 1: 200 000 while in the period from 1989 to 1991, he was leader of the project for the Republic of Macedonia.



**Figure 3:** Field trip with students in second year of studies at the Faculty of Forestry in the early 90's, site Kozarica - Shar Mountain (Prof. Dr. Radoslav Rizovski first from left)

The team was made up of well-known phytocoenologists in Macedonia on the front with Em and his associates: Dzekov, Nikolovski, Rizovski, Andonoski, later they are joined by Matveyeva, Lj. Micevski, Mirčevski, Batkoski, Gudeski, Dimovski, Manevski and others. From this study, Prof. Rizovski in cooperation with Professor Em in 1974, draft a *Prodromus of plant communities of SR Macedonia* (manuscript).

In 1985, Professor Rizovski participates at the colloquium held in Bribir-Ilok, which actually was a gathering of eminent phytocoenologists from the former Yugoslavia organized as Scientific Council of the Vegetation map of Yugoslavia, in order to settle the some problems in Phytocoenolgy, and progress the making of the vegetation map of Yugoslavia.

From this gathering comes out *Prodromus phytocoenosum Jugoslaviae ad mappam vegetationis m 1: 200 000*, which with certain adjustments for the newly described communities is still used today.

In accordance with the determined objective of the project "Vegetation map of Yugoslavia" undertaken were obligations to be developed 2 types of the map:

- Map of the natural potential climatic vegetation zones with scale of R 1: 500 000, which for the territory of the Republic of Macedonia was created by Professor Rizovski, and the same was been printed as appendix in the book "Climate vegetation soil zones (regions) in the Republic of Macedonia", published by the Macedonian Academy of Sciences and Arts, 1996

- Map of recent (real) vegetation in the measurement scale of 1: 200 000

The Project "Vegetation Map of Macedonia" continues to early nineties of the last century but it is still not finished.

Although this project remains unfinished, under the guidelines of Professor Em, Professor Rizovski and Professor Dzekov have been working on the refugium vegetation of Macedonia. From this project arised several described communities, as that of the Macedonian oak (*Q.macedonica*), the narrow-leaved ash (*F. angustifolia*) and several communities from the gorges of the major rivers.

Professor Rizovski due to prolonged sickness passed away on July 20<sup>th</sup> 2008.



**Figure 4:** Prof. Dr. Radoslav Rizovski on a field research



## 6 BIBLIOGRAPHY OF PROF. DR. RADOSLAV RIZOVSKI

1968

- [1] Гудески, А., Ризовски, Р.: Најниско наоѓалиште на елата во СР Македонија, ШП. Бр 3-4, стр. 44-47, 1968, Скопје 1970
- [2] Ризовски, Р.: Споредување на горун-габерови шуми од Северна Хрватска и СР Македонија. ГЗШФ, (шумарство), кн. 23, 1970

1970

- [3] Јоветиќ, Р., Трпков, Б., Ризовски, Р.: Еколошки карактеристики на еребицата камењарка на подрачјето на планината Жеден, ГЗШФ, кн.23, 1970 Скопје

1972

- [4] Спиоровски, Ј., Ризовски, Р.: За особеностите на почвите врз дијабаз и габро под некои типови горувни шуми во потесната област на Демир Капија. ГЗШФ, кн. 23, 1972, Скопје

1973

- [5] Цветковиќ, Д., Ризовски, Р.: Помолошке карактеристике неких дивљих врста купине (*Rubus ssp.*) у СР Македонији. Југословенско воћарство бр.25-26, 1973 Чачак.
- [6] Миновски, Д., Ризовски, Р.: Прилог кон проучувањето на автохтони форми дренки – *Cornus mas* L. во некои локалитети во СР Македонија. Социјалистичко земјоделство бр.7-9, 1973, Скопје
- [7] Миновски, Д., Ризовски, Р.: Прилог кон проучувањето на сливи џанки (*Prunus cerasifera* Ehrh) во некои локалитети во СР Македонија. Социјалистичко земјоделство бр.4-6, 1973, Скопје

1974

- [8] Ризовски, Р., Цветковиќ, Д.: Родот *Rubus* L. – капини. Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје
- [9] Ризовски, Р., Миновски, Д.: *Prunus spinosa* L. – трнка, тринка. Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје
- [10] Ризовски, Р., Миновски, Д.: *Corylus avellana* L. – леска. Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје
- [11] Ризовски, Р., Миновски, Д.: *Corylus colurna* L. – дива леска, мечја лека Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје
- [12] Миновски, Д., Ризовски, Р.: *Cornus mas* L. – леска, Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје
- [13] Миновски, Д., Ризовски, Р.: *Prunus cerasifera* Ehrh. во СР Македонија, Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје

- [14] Миновски, Д., Ризовски, Р.: *Vitis silvestris* L. дива лоза, Зборник на трудови – (посебно издание) на Зем. Шум. 1974 Скопје.
- [15] Ризовски, Р.: Ценози на дабот плоскач (*Quercus farnetto* Ten.) како засебен вегетациски појас во Долното Повардарје ГЗШФ (шумарство), кн. 26, 1974 Скопје.
- [16] Rizovski, R. Cenoze hrasta sladuna (*Quercus frainetto* Ten.) u submediteranskoj zoni donjeg Povardarja. Izvadak iz doktorske disertacije. Poroč. Vzhodno alp.-dinar. Preuč.veget. Tom 14, SAZU, Ljubljana.
- [17] Јоветиќ, Р., Трпков, Б., Ризовски, Р., Биолошки карактеристики на еребицата камењарка (*Alectoris graeca* M.), Ловец бр. 4-5-6, 1974 Скопје

1975

- [18] Спиоровски, Ј., Ризовски, Р.: Почвите под некои плоскачеви шуми во Долното Повардарје. Год. Збор. на Зем. Шум. Фак.1975 Скопје
- [19] Ризовски, Р.: Simpozij za organizaciju mreže trajno zaštićenih površina u Jugoslaviji i njihovo istraživanje, Š.L. 7-10, s.363

1977

- [20] Rizovski, R., Beitrag zur Kenntnis der Verbreitung von *Dioscorea balcanica* Koš. Glasn. Republic. Zav. Zašt. Prirode – Prirodnj. Muzeja, 10: Titograd

1978

- [21] Цеков, С., Ризовски, Р.: Шумската растителност во Малеш и Пијанец. МАНУ (посебно издание) Малеш и Пијанец I – Вегетација, Скопје
- [22] Ризовски, Р.: Фитоценологија со основи на типологија на шумите и пасиштата стр. 1-307 (скрипта), Шумарски факултет – Скопје

1982

- [23] Ризовски, Р.: Вегетацијата на планината Црна Гора во сливовите на Липковска и Летничка река. Шумарски преглед 1-2, Скопје.

1983

- [24] Rizovski, R.: Fitocenoza crnog graba (*Ostrya carpinifolia* Scop.) na silikatnoj podlozi u Makedoniji. Radovi, knj. 122, odeljenje prirodnih i matematičkih nauka, kn. 21 kn.72 ANUBiH, Sarajevo.
- [25] Rizovski, R.: Glavni tipova šuma SR Makedonije. Šum. Encikl., knj.2, Leksikogr. Zavod, Zagreb
- [26] Ризовски, Р.: Фитоценозата на бука и ситнолисна липа (*Tilio cordatae* – *Fagetum* Ass. nov.) во Македонија, МАНУ, Прилози IV 1-2 Оделение за биолошки и медицински науки, Скопје

- 1984
- [27] Ризовски, Р.: Нови наоѓалишта на ретки растенија, Културно наследство IX, Скопје, 160
- 1985
- [28] Ем, Х., Џеков, С., Ризовски, С.: За рефугијалната шумска вегетација во СР Македонија МАНУ, Прилози VI 1-2 Одделение за биолошки и медицински науки, Скопје
- 1986
- [29] Rizovski, R.: Šume i prerada drveta Jugoslavije – Endemska dendroflora Jugoslavije – Beograd  
[30] Jovanović, B., Lakušić, R., Rizovski, R., Trinajstić, I., Zupančić, M.: Prodrum phytocenosum Jugoslaviae ad mappam vegetationis M 1 : 2 000 000 Naučno veće vegetaciske karte Jugoslavije, Bribir-Ilok
- 1987
- [31] Rizovski, R., Dzekov, S.: Šumske zajednice Jugoslavije - Makedonija. Šum. Encikl., knj.III, Leksikogr. Zavod, Zagreb  
[32] Љ. Мицевски, Ризовски, Р.: Шумско-вегетациска карактеристика на Мариово. Зборник на трудови Мариово природни и социо-економски обележја и можности за развој-Друштво за наука и уметност –Прилеп-Битола.
- 1988
- [33] Em, H., Rizovski, R., Dzekov, S.: Ein Beitrag zur Kenntnis refugialer Vegetation Mazedoniens. Bushen – Hopfenbushenmischwald im Treskakanon, Atti del simposio della societa estalpino – dinarica di fitosociologia: Feltre 29 giugno – 3 luglio 1988
- 1989
- [34] Ризовски, Р.: Фитоценологија – (поглавје во средношколски учебник- Дендрологија), Просветно дело, Скопје  
[35] Ризовски, Р.: Еколошки и биолошки карактеристики на вегетациските подрачја во Македонија. Хортус, год 5, бр 9-10, Скопје
- 1990
- [36] Ризовски, Р., Џеков, С.: Шумска вегетација на Бистра. Едиција Бистра, кн. II, МАНУ, Скопје.  
[37] Ризовски, Р.: Пасиштата како одделна пејсажна структура во поврзувањето на земјоделските и шумските комплекси. Хортус, год 6, бр 11-12, Скопје
- 1996
- [38] Филиповски, Ѓ., Ризовски, Р., Ристевски, П.: Карактеристики на климатско-вегетациските зони (региони) во Република Македонија, МАНУ  
[39] Ризовски, Р., Групче, Љ., Ризовска - Атанасовска, Ј.: Рефугиуми во клисурите на р.
- Треска како простори за заштита на биолошката разновидност во Р. Македонија. Симпозиум: Национални паркови и нивната улога во заштитата на биодиверзитетот на Балканскиот полуостров, Охрид.
- 1997
- [40] Rizovski, R.: Vegetation and its importance for the protection of Prespa region. Internac. Symp. Towards Integrated Conservation and Sustainable Development of Transboundary Macro and Micro Prespa Lakes . Korcha, Albania. PPNEA, Tirana  
[41] Трпков, Б., Ризовски, Р., Здравковски, Т., Шумски резерват ЈАСЕН- (монографија)
- 1998
- [42] Групче, Љ., Лозановски, Р., Ризовски, Р.: Развој на екологијата како наука во Република Македонија. I Конгрес на еколозите на Македонија со меѓународно учество - Книга на пленарни реферати, Охрид  
[43] Ризовски, Р., Ризовска Атанасовска, Ј.: Заедницата на македонскиот даб (*Quercus trojana* Webb.) во долината на реката Раец. I Конгрес на еколозите на Македонија со меѓународно учество-Зборник на трудови Том 1  
[44] Меловски, Љ., Ризовски, Р., Караделев, М., Христовски С.: Медитерански карактеристики на флората, вегетацијата и габите на Островот Голем Град на Преспанското езеро. Публикација на Новинарски Еколошки центар (НЕЦ), Скопје.
- 2000
- [45] Ризовски, Р., Ризовска Атанасовска, Ј.: Прилог кон познавање на биодиверзитетот во рефугијалната област Нице-Кожуф. Јубилеен годишен зборник на Шумарски факултет-Скопје  
[46] Ризовски, Р., Групче, Љ., Ризовска Атанасовска, Ј.: Биолошка разновидност на охридско – преспанското подрачје како основа за прогласување на заштитен трансграничен регион. Меѓународен симпозиум Одржлив развој на преспанскиот регион.  
[47] Ризовски, Р., Ризовска Атанасовска, Ј.: Улогата на варовничко – доломитните црници за секуларната и рецентна сукцесија на вегетацијата во Република Македонија, МАНУ-Скопје
- 2004
- [48] Ризовски, Р., Ризовска Атанасовска, Ј.: Особениости на флората и вегетацијата на Церипашина Планина во сливот на Тетовска Река II Конгрес на еколозите на Македонија со меѓународно учество-Зборник на трудови

2009

- [49] Ризовски, Р.: Вегетациски подрачја на Република Македонија. Македонска енциклопедија кн. 1 МАНУ
- [50] Ризовски, Р.: Шумски заедници во Република Македонија. Македонска енциклопедија кн. 2 МАНУ

## 9 ACKNOWLEDGEMENTS

The Author is grateful to Blagoj Nikolov for his helpful cooperation about the English translation of the text, and to Joso Vukelić for the photo used in this article.



**Figure 5:** Prof. Dr. Radoslav Rizovski as a member of PhD thesis defence commission (Photo: J. Vukelić)



# A PHYTOCOENOLOGICAL STUDY OF FORESTS OF HUNGARIAN OAK AND TURKEY OAK (*QUERCETUM FRAINETTO-CERRIDIS* /RUDSKI 1949/ TRINAJSTIĆ ET AL. 1996) ON THE NORTHWEST BORDER OF ARRIVAL (NATURE PARK PAPUK, CROATIA)

<sup>1</sup>BARIČEVIĆ D., <sup>1</sup>VUKELIĆ J., <sup>2</sup>PUAČA M., <sup>1</sup>ŠAPIĆ I.

<sup>1</sup>University of Zagreb, Faculty of Forestry, Zagreb, Croatia

<sup>2</sup>Hrvatske šume, limited liability company, Zagreb, Croatia

Corresponding author e-mail address: dario.baricevic@zg.htnet.hr

**ABSTRACT:** The paper presents phytocoenological research of forests of Hungarian and Turkey oak in the locality of Otmanov Vis within Papuk Nature Park. These specific and very interesting areas has not so far been included in research of a broader distribution range of the community of Hungarian and Turkey oak and represents a unique feature because of very rare soil type (eutric ranker on trachyandesite). Phytocoenological sampling based on the principles of the Zürich-Montpellier School was performed on five plots. Our own relevés were compared with those from a broader Kutjevo area, Bosnia and Herzegovina and northwestern Serbia. Statistical analysis was accomplished using Syn-tax 2000 software. According to the results, the studied stands belong to the association *Quercetum frainetto-cerridis* (Rudski 1949) Trinajstić et al. 1996, or more precisely to its subassociation *ruscetosum* Jovanović et Dunjić 1951. Syntaxonically, they belong to the alliance *Quercion frainetto*, order *Quercetalia pubescentis*, and class *Quercio-Fagetea*. In addition to the characteristic species of the association, the subassociation in the study area is well defined by the following differential species of the subassociation: *Cornus mas*, *Helleborus odorus*, *Melica uniflora*, *Teucrium chamaedrys*, *Ruscus aculeatus*, *Physospermum cornubiense* and *Campanula persicifolia*. In terms of synecology, it is characterized by its occurrence in the colline belt, at elevations from 260 to 400 m and inclinations from 5 - 40°. As Otmanov Vis is the only area in Croatia where this subassociation is occurs this work also presents the first description of this subassociation in Croatia. The site should be placed under special protection due to exceptional biological diversity and beauty of the landscape.

**Keywords:** phytocoenological study, *Quercetum frainetto-cerridis*, Croatia, *ruscetosum*, first description.

## 1 INTRODUCTION

The forest of Hungarian oak and Turkey oak (*Quercetum frainetto-cerridis* Rudski /1949/ Trinajstić et al. 1996) is a climatogenic phytocoenosis occurring over a large part of the Balkan Peninsula. In Croatia, it has been studied in more detail by Trinajstić et al. [23] in the Slavonian uplands, which also form the north-western boundary of its occurrence. Otherwise, the Hungarian oak is a rare forest species in Croatia. It is distributed in Slavonia and in several places in Dalmatia, but it covers more extensive areas only on the southern slopes of Mt. Krndija. It appears in smaller enclaves surrounded by stands of sessile oak and common hornbeam, where it forms stable stands of high commercial and scientific value. Trinajstić et al. [23] define it as a mesophilic subassociation *carpinetosum betuli*, whereas the conclusion of the latest research [19] urges for a clear definition of plant community in the locality of Otmanov Vis, since it was observed that floristically, it differs fundamentally from the neighbouring, mesophilic subassociation studied by Trinajstić et al. [23]. This was corroborated by Vukelić [24] in his general description of the forest of Hungarian and Turkey oaks in Croatia.

The forest stands in the locality of Otmanov Vis, covering an area of 24.76 ha, occur under specific synecological conditions and differ significantly from the neighbouring stands in terms of flora and management. The specific synecological conditions have also enabled the growth of a large number of rare and protected species, making them a valuable object of research. In general, stands of Hungarian and Turkey oaks in the area of the Slavonian uplands cover 894 ha. It is interesting that Hungarian oak in this area was first recorded by the botanist P. Kitaibel as far back as 1808, which was also the first record of this species on the Balkan Peninsula [4]. Hungarian oak stands in the Slavonian uplands were also discussed by Fukarek [4, 5], Fukarek et al. [7],

Cestar et al. [3], and Zelić [26, 27].

All the above prompted us to carry out a phytocoenological study of these stands in the subject area, make a detailed analysis of the floristic composition, compare the new relevés with those from a few previously published studies, and thus provide their clear definition, nomination and description. This paper aims to present the undertaken studies.

## 2 RESEARCH AREA

The research area encompasses the locality of Otmanov Vis (45.43° N, 17.95° E) on the southern slopes of Mt. Krndija within Papuk Nature Park (Fig. 1). Mt. Krndija, together with Psunj and Papuk in the west, and Požeška Gora and Dilj in the south, builds a mountain chain that surrounds the Požega Valley and forms the mountains of Slavonia. The mentioned mountains are isle mountains that stretch between the Podravina and Posavina lowlands in the south-western boundary part of the Pannonian Plain.

The research area is characterized by hilly-colline features. Diverse microrelief, microclimatic, geological and edaphic conditions alternate over a relatively small area. Elevations range between 260 and 400 m a.s.l.

The basic hills of Krndija are made up of diverse metamorphic rocks from the Paleozoic era, which are transgressively followed by the lower Miocene marine sediments with the effusion of trachyandesite, while Quaternary sediments are developed in the lower parts of the terrain. Igneous rocks are covered with Quaternary layers of loess that erodes gradually [22].

According to research Puača et al [19], eutric ranker on trachyandesite is developed on the complex and multilayered lithological bedrock as the dominant soil type. The multi-component composition of pedocartographic units includes eutric ranker, regolithic and browned, eutric brown soil and eutric colluvium

(70:15:15). Pedological data analysis shows that this soil has very high humus content (according to Gračanin) of weak acidic reaction. The soil is very rich in nitrogen and potassium and rich in physiologically active phosphorus. In terms of mechanical composition, the analyzed soil is texturally marked as clayey loam. The limiting factors of soil productivity are its small depth and skeletal nature, which are conditioned by the properties of lithological bedrock, terrain inclination and constant erosion.



**Figure 1:** View at the site Otmanov Vis - spring aspect

According to the data for Požega meteorological station, the mean annual air temperature is about 11 °C, and the mean annual precipitation is between 750 and 800 mm. In the Köppen classification, the climate type is Cfbw"x". According to the Thornthwaite's classification system, the study area extends in the area of humid climate.

### 3 MATERIAL AND RESEARCH METHODS

Phytocoenological research into the community of Hungarian oak in the study area was performed in two stages. The first stage consisted of collecting phytocoenological relevés. Vegetation in the plots was sampled according to the principles of the Zürich-Montpellier Phytocoenological School [2]. Five phytocoenological plots were established in the locality of Otmanov Vis. Phytocoenological relevés were combined with the collection of individual synecological data in the field, such as the elevation, exposition, inclination and coordinate position of each relevé using a GPS device (Garmin Montana 650). The size of the relevés (plots) was 20m x 20m.


The second stage consisted of processing the data acquired in the field. The relevés were entered into the Turboveg database [10] together with 9 relevés from a wider surroundings of Kutjevo [23], 10 relevés from Bosnia and Herzegovina - including 3 relevés from the area of Croatia [7], and 10 relevés from north-western Serbia [13]. Statistical processing was performed using Syn-tax 2000 software [18]. Two methods of numerical analysis were applied: cluster analysis (Single linkage, Complete linkage, Average linkage method and Increment sum of squares method) and multidimensional scaling (Principal coordinates analysis). The Similarity ratio was used for both methods.

The species in the phytocoenological table were classified according to the social affiliation of the species


[24], while plant nomenclature was adjusted according to the Flora Croatica database [17].

### 4 RESEARCH RESULTS AND DISCUSSION

The results of classical analysis of 34 phytocoenological relevés and the applied cluster methods (Fig. 2) and the analysis obtained by multidimensional scaling (Fig. 3) showed that the relevés were placed in four groups. Our phytocoenological relevés were grouped into a separate set. In order to clearly define the differences and similarities between individual sets of relevés, the floral composition was analyzed and compared in detail. The results of these analyses clearly show that the relevés in the study area of Otmanov Vis systematically belong to the association *Quercetum frainetto-cerridis* (Rudski 1949) Trinajstić et al. 1996 (Table 1).



**Figure 2:** Dendrogram produced by cluster analysis – Incremental sum of squares



**Figure 3:** Multidimensional scaling - Principal coordinates analysis

Syntaxonomic affiliation to the association *Quercetum frainetto-cerridis* was confirmed by the species recorded there, including *Quercus frainetto*, *Quercus cerris*, *Pyrus pyraeaster*, *Carpinus betulus*, *Acer campestre*, *Fraxinus ornus*, *Crataegus monogyna*, *Ligustrum vulgare*, *Potentilla micrantha*, *Glechoma hirsuta*, *Brachypodium sylvaticum*, *Hieracium sabaudum*, *Carex flacca*, *Moehringia trinervia* and others, which are characteristic for this association (Table I). Furthermore,

differences in the floristic composition suggest that the association should be affiliated to a lower syntaxonomic unit. To define lower systematic units, we should refer to the already defined subassociations which would possibly correspond to the studied stands. Thus, the association *Quercetum frainetto-cerridis* is divided into several different types that are described as different subassociations. This is the result of diverse variations in the synecological conditions of this association, as already discussed by Glišić [8]. The basic type is the typical forest of Hungarian oak and Turkey oak (*Quercetum frainetto-cerridis* subas. *typicum*). In terms of soil humidity and acidity it is almost identical to the forest of sessile oak and common hornbeam. It is characterized by the relatively rich, luscious and diverse herb layer. The second type is the forest of Hungarian oak and Turkey oak with common hornbeam (*Quercetum frainetto-cerridis* subas. *carpinetosum betuli*). The first was described by Rudski [20], and it encompasses areas that border with the forest of sessile oak and common hornbeam. Their principal differential species are *Carpinus betulus* and some other species of the alliance *Fagion*, then *Acer campestre*, *Prunus avium*, *Moehringia trinervia* and others. The third type is the forest of Hungarian oak and Turkey oak with butcher's broom (*Quercetum frainetto-cerridis* subas. *ruscetosum*) which was described by Jovanović et Dunjić [11]. Ecologically and geographically, it is a separate phytocoenosis that occurs in the southern part of the Pannonian Plane and grows in thermophilic habitats [25]. Its principal differential species is butcher's broom (*Ruscus aculeatus*). The fourth type is the forest of Hungarian oak and Turkey oak with oriental hornbeam (*Quercetum frainetto-cerridis* subas. *carpinetosum orientale*). This is a phytocoenosis with xerophilic species with oriental hornbeam (*Carpinus orientalis*) featuring as the main discriminating species. It should be pointed out that owing to frequent degradation of these forests, their floral composition undergoes changes. The degraded stages include the subassociations *Quercetum frainetto-cerridis* subas. *nudum* and subas. *hieracietosum* [13]. Apart from these subassociations, the Habitat Handbook of Serbia [1] also lists the subassociations *comandretosum* B. Jovanović 1968, *fagetosum* E. Vukičević 1959, *juglandetosum* E. Vukičević 1974, *paeonietosum* M. Janković et Nikolić 1967, *physospermetosum* Rudski 1940, *pubescentosum* Jovanović, *petraeae* Z. Tomić 1989, *roboris* B. Jovanović et Z. Tomić 1978, *virgilianae* B. Jovanović et Vukičević 1977, and *scardicum* Krasniqi (1968) 1972 in Kosovo. In Macedonia, such forests are among the other described in Galičica National Park under the name of *Quercetum frainetto-cerris* Horvat 1954 [14] or *Quercetum frainetto-cerris macedonicum* Oberdorfer 1948 emend Horvat 1959 [15, 16]. Examples of geographic adjectives in the names of the subassociations are found in some other authors, e.g. *thracicum* = *bulgaricum* [9], *moesiicum* = *serbicum* [12]. In the latest analysis of the forest of Hungarian oak and Turkey oak in Bosnia and Herzegovina are described as *Quercetum frainetto-cerridis* (Rudski 1949) Trinajstić et al. 1996 because Trinajstić et al. [23] corrected its original illegitimate name "*Quercetum confertae-cerris serbicum*" [21]. They also conclude that in the southern B&H described association *Quercetum frainetto hercegovinum* [6] could not be separate from other *Quercetum frainetto-cerridis* forests.

The floristic composition of Hungarian oak-Turkey

oak forests in the proximity of the study area (the surroundings of Kutjevo) was described and investigated by Trinajstić et al. [23]. These stands are discriminated from the typically composed stands of the association from the centre of its range in Serbia by the absence of the species *Tilia tomentosa*, *Sorbus domestica*, *Rosa gallica*, *Danna cornubiensis*, *Lychnis coronaria*, *Silene viridiflora*, *Tanacetum corymbosum*, *Trifolium alpestre* and many others. Based on the analysis, they affiliate these stands to the subassociation *carpinetosum betuli*.

However, the association described by Trinajstić et al. [23] also differs fundamentally from the studied stands in the locality of Otmanov Vis in terms of its floral composition and habitat conditions. The analysis of the floristic composition in the study area clearly shows independence at the level of subassociation (Table I). Because of the specific habitat conditions in the study area, there are species that are absent from the previously described subassociation, such as *Tilia tomentosa*, *Danna cornubiensis*, *Tanacetum corymbosum*, *Euphorbia cyparissias*, *Melica uniflora*, *Helleborus odoratus*, *Dryopteris filix-mas*, *Physospermum cornubiense*, *Teucrium chamaedrys*, *Asplenium adiantum-nigrum*, *Calamintha nepeta*, *Poa nemoralis*, *Dictamnus albus*, *Spiraea chamaedryfolia*, *Ruscus aculeatus* and others. In terms of floral composition and the presence of characteristic and differential species, the community in the study area best corresponds to the forest of Hungarian oak and Turkey oak with butcher's broom *Quercetum frainetto-cerridis* subassociation *ruscetosum*, which was described in the work by Jovanović et Dunjić [11]. According to this description, the subassociation is developed over flat and mildly sloping terrains, of weak acid soil reaction. There is occurrence of thermophilic, Mediterranean floral elements, which were preserved during the ice age owing to the warm climate on the shores of the former Pannonian Sea. The described community contains species which were found in the study area, thus determining its affiliation to the subassociation of the forest of Hungarian oak with Turkey oak and butcher's broom. Differential species of this subassociation include *Cornus mas*, *Helleborus odoratus*, *Melica uniflora*, *Teucrium chamaedrys*, *Ruscus aculeatus*, *Physospermum cornubiense* and *Campanula persicifolia*.

The study locality is specifically characterized by a very high abundance of the species *Digitalis grandiflora*, *Carex divulsa*, *Alliaria petiolata*, *Fallopia convolvulus*, *Dictamnus albus* and *Spiraea chamaedryfolia*, which will be the subject of further vegetation and floristic research. Of additional interest are the beautiful seasonal aspects that begin with the awakening of the vegetation in early March, as well as the dominance of hellebore (*Helleborus odoratus*), dogwood (*Cornus mas*) and spring flowers. April is characterized by the dominance of the facies *Alliaria officinalis*. The end of April is marked by the abundant appearance of leopard's bane (*Doronicum columnae*). The beginning of May is characterized by luscious ground vegetation, when the main ridge is covered with nationally protected species (*Dictamnus albus*), early purple orchid (*Orchis mascula*), martagon lily (*Lilium martagon*) and others. Manna ash (*Fraxinus ornus*), germander meadowsweet (*Spiraea chamaedryfolia*) and dog rose (*Rosa canina*) are in full bloom. Large yellow foxglove (*Digitalis grandiflora*) appears in abundance at the beginning of June.



**Table I:** Comparison of the relevés from the study area with the relevés of other types of forests of Hungarian oak and Turkey oak

Nr. of releve		30	31	32	33	34	Presence degree	Presence degree	Presence degree	Presence degree
Date (year/month)		2014/6	2014/6	2014/6	2014/6	2014/6				
Coordinate x		6496766	6496759	6496487	6496407	6496234				
Coordinate y		5031094	5031061	5031006	5031022	5030898				
Releve area (m <sup>2</sup> )		400	400	400	400	400				
Elevation (m)		375	395	380	375	340				
Exposition (degrees)		0	0	300	0	180				
Inclination (degrees)		35	0	25	40	10				
Cover layer of trees (%)		70	50	90	60	70				
Cover layer of shrubs (%)		40	20	5	15	10				
Cover layer of ground vegetation (%)		60	80	30	60	70				
Source							New relevés Otmanov Vis	Trinajstić et al. 1996	Fukarek et al. 1974	Jovanović 1967
Floral composition:										
Characteristic species of association:										
<i>Quercus frainetto</i>	A	4	1	3	1	3	V	V	V	V
<i>Fraxinus ornus</i>		2	2	2	4	3	V	I		III
<i>Quercus cerris</i>		+	3	.	.	1	III	III	V	V
<i>Tilia tomentosa</i>		.	.	3	.	.	I		III	
<i>Ligustrum vulgare</i>	B	.	+	+	.	+	III	V	V	II
<i>Fraxinus ornus</i>		.	.	.	1	1	III	IV	V	III
<i>Acer campestre</i>		+	+	.	.	.	II	V	V	II
<i>Crataegus monogyna</i>		.	+	.	+	.	II	V	V	III
<i>Rosa arvensis</i>		.	.	.	+	+	II	III	IV	
<i>Quercus frainetto</i>		.	.	.	.	+	I	IV		V
<i>Chamaecytisus supinus</i>		.	.	.	.	.		IV		
<i>Quercus cerris</i>		.	.	.	.	.		I		V
<i>Potentilla micrantha</i>	C	1	1	1	1	+	V	V	V	IV
<i>Hypericum perforatum</i>		+	+	+	+	+	V	II		IV
<i>Brachypodium sylvaticum</i>		1	1	1	.	+	IV	III		II
<i>Calamintha nepeta</i> agg.		1	2	1	.	1	IV			II
<i>Fraxinus ornus</i>		+	.	.	.	.	I	II		IV
<i>Quercus frainetto</i>		1	.	.	.	.	I	III	IV	V
<i>Veronica chamaedrys</i>		+	.	.	.	.	I	V		IV
<i>Clinopodium vulgare</i>		.	+	.	.	.	I		V	II
<i>Rosa arvensis</i>		.	+	.	.	.	I			
<i>Trifolium medium</i>		.	.	.	.	+	I	II		
<i>Tilia tomentosa</i>		.	.	.	.	.		I		II
<i>Quercus cerris</i>		.	.	.	.	.		I	III	IV
Differential species of subassociation <i>carpinetosum betuli</i> :										
<i>Prunus avium</i>	A	+	.	.	.	.	I	I	III	
<i>Carpinus betulus</i>		.	.	.	.	.		III		
<i>Acer campestre</i>		.	.	.	.	.		III		
<i>Carpinus betulus</i>	B	.	+	.	.	.	I	V	V	V
<i>Prunus avium</i>		.	.	.	.	.		III		
<i>Moehringia trinervia</i>	C	+	.	+	.	.	II	III	I	
<i>Helleborus croaticus</i>		.	.	.	.	.		V		
<i>Prunus avium</i>		.	.	.	.	.		III		II
<i>Carpinus betulus</i>		.	.	.	.	.		II	III	II
<i>Acer campestre</i>		.	.	.	.	.				II
Differential species of subassociation <i>ruscetosum</i> :										
<i>Cornus mas</i>	B	3	2	.	1	+	IV	II	I	
<i>Ruscus aculeatus</i>		+	.	.	.	1	II			
<i>Helleborus odoratus</i>	C	2	1	2	+	1	V		IV	
<i>Melica uniflora</i>		1	3	1	3	1	V		I	I
<i>Teucrium chamaedrys</i>		.	+	+	+	+	IV		II	
<i>Physospermum cornubiense</i>		+	.	+	.	.	II		III	IV
<i>Campanula persicifolia</i>		+	+	.	.	.	II	I	I	III
<i>Quercetalia pubescentis</i> :										
<i>Sorbus torminalis</i>	B	+	+	.	.	.	II	II	III	I
<i>Acer tataricum</i>		.	.	.	.	.		III	V	II
<i>Sorbus domestica</i>		.	.	.	.	.		II		I
<i>Viola alba</i>	C	+	+	+	.	.	III	III		
<i>Tanacetum corymbosum</i>		+	.	+	+	.	III	I	I	
<i>Lathyrus venetus</i>		+	.	.	.	.	I	II	III	
<i>Tamus communis</i>		.	+	.	.	.	I	I	II	III

BARIČEVIĆ D., VUKELIĆ J., PUAČA M., ŠAPIĆ I.  
A PHYTOCOENOLOGICAL STUDY OF FORESTS OF HUNGARIAN OAK AND TURKEY OAK (*QUERCETUM FRAINETTO-CERRIDIS* /RUDSKI 1949/ TRINAJSTIĆ ET AL. 1996) ON THE NORTHWEST BORDER OF ARRIVAL (NATURE PARK PAPUK, CROATIA)

continuation of Table I

<i>Lathyrus niger</i>		.	.	.	.	.	.	I	V	IV
<i>Acer tataricum</i>		.	.	.	.	.	.	I		III
<i>Convallaria majalis</i>		.	.	.	.	.	.	I		
<i>Polygonatum odoratum</i>		.	.	.	.	.	.	I		
<i>Sorbus torminalis</i>		.	.	.	.	.	.			III
<i>Oenanthe pimpinelloides</i>		.	.	.	.	.	.	I		
<i>Fagetalia</i> :										
<i>Fagus sylvatica</i>	A	.	.	.	.	.	.	I		
<i>Fagus sylvatica</i>	B	.	.	.	.	.	.	III		
<i>Mycelis muralis</i>	C	.	+	+	2	+	IV	III		I
<i>Stellaria holostea</i>		+	+	+	1	.	IV	I	III	
<i>Glechoma hirsuta</i>		+	1	.	.	.	II	V	II	
<i>Lilium martagon</i>		+	.	.	.	.	I	I		
<i>Symphytum tuberosum</i> agg.		.	.	.	.	.		III	I	
<i>Viola reichenbachiana</i>		.	.	.	.	.		III		I
<i>Primula vulgaris</i>		.	.	.	.	.		III	IV	
<i>Carex sylvatica</i>		.	.	.	.	.		III	I	
<i>Pulmonaria officinalis</i>		.	.	.	.	.		III	II	
<i>Galium odoratum</i>		.	.	.	.	.		II		
<i>Polygonatum multiflorum</i>		.	.	.	.	.		I		
<i>Sanicula europaea</i>		.	.	.	.	.		I	I	
<i>Neottia nidus-avis</i>		.	.	.	.	.		I		
<i>Scrophularia nodosa</i>		.	.	.	.	.		I		
<i>Geranium robertianum</i>		.	.	.	.	.		II	I	
<i>Dentaria bulbifera</i>		.	.	.	.	.		I		
<i>Melica nutans</i>		.	.	.	.	.		I		
<i>Calamintha grandiflora</i>		.	.	.	.	.		I		
<i>Epimedium alpinum</i>		.	.	.	.	.		I	II	II
<i>Stachys sylvatica</i>		.	.	.	.	.		I		
<i>Circaea lutetiana</i>		.	.	.	.	.		I		
<i>Asarum europaeum</i>		.	.	.	.	.		I	I	
<i>Salvia glutinosa</i>		.	.	.	.	.		I		
<i>Quercetalia robori-petraeae</i> :										
<i>Castanea sativa</i>	A	.	.	.	.	.				I
<i>Castanea sativa</i>	B	.	.	.	.	.		I		I
<i>Festuca heterophylla</i>	C	2	.	1	1	1	IV	I	III	III
<i>Hieracium sabaudum</i>		+	.	.	.	.	I	III		IV
<i>Veronica officinalis</i>		+	.	.	.	.	I	III	II	V
<i>Luzula forsteri</i>		+	.	.	.	.	I	III	II	III
<i>Melampyrum sylvaticum</i>		.	.	.	.	.		III	III	
<i>Serratula tinctoria</i>		.	.	.	.	.		I	I	
<i>Castanea sativa</i>		.	.	.	.	.				I
<i>Quercu-Fagetea</i> :										
<i>Quercus petraea</i>	A	+	.	.	.	.	I	IV	IV	IV
<i>Quercus robur</i>		.	.	.	.	.		I	I	
<i>Pyrus pyraister</i>	B	.	+	.	.	+	II	IV	V	
<i>Quercus petraea</i>		.	.	.	.	.		I		III
<i>Rubus hirtus</i> s.lat.		.	.	.	.	.		I		
<i>Corylus avellana</i>		.	.	.	.	.		I	I	I
<i>Euonymus europaeus</i>		.	.	.	.	.		I	II	
<i>Cruciata glabra</i>	C	.	+	.	.	.	I	III		III
<i>Quercus petraea</i>		.	.	.	.	.		III	III	III
<i>Pyrus pyraister</i>		.	.	.	.	.				III
<i>Corylus avellana</i>		.	.	.	.	.				I
<i>Galium sylvaticum</i>		.	.	.	.	.		III		
<i>Fragaria vesca</i>		.	.	.	.	.		I		II
<i>Cephalanthera longifolia</i>		.	.	.	.	.		I		
<i>Galium schultesii</i>		.	.	.	.	.		I		
<i>Hedera helix</i>		.	.	.	.	.		I		I
<i>Rhamno-Prunetea</i> :										
<i>Cornus sanguinea</i> ssp. <i>hungarica</i>	B	.	.	.	.	.		IV		
<i>Clematis vitalba</i>		.	.	.	.	.		III	III	
<i>Cornus sanguinea</i>		.	.	.	.	.		II	III	
<i>Crataegus nigra</i>		.	.	.	.	.		II		
<i>Prunus spinosa</i>		.	.	.	.	.		II	II	
<i>Euonymus europaeus</i>	C	.	+	.	.	.	I	I		
<i>Prunus spinosa</i>		.	.	.	.	.		I		
<i>Crataegus monogyna</i>		.	.	.	.	.				II

continuation of Table I

<i>Ligustrum vulgare</i>		-	-	-	-	-	-	-	-	II
Other species:										
<i>Genista tinctoria</i>	B	1	-	-	-	-	I	III	IV	
<i>Rubus ulmifolius</i>		-	-	-	-	-	-	IV		
<i>Juniperus communis</i>		-	-	-	-	-	-	II		III
<i>Dactylis glomerata</i>	C	2	2	1	-	+	IV	V	V	
<i>Geum urbanum</i>		-	+	-	-	-	I	III		
<i>Carex flacca</i>		-	-	-	-	+	I	III		
<i>Galium lucidum</i>		+	-	-	-	-	I	II		
<i>Vicia species</i>		+	-	-	-	-	I	II		
<i>Silene nutans s.lat.</i>		-	-	-	+	-	I	I		
<i>Vincetoxicum hirsutidaria</i>		-	+	-	-	-	I	I		
<i>Ajuga reptans</i>		-	-	-	-	-	-	V	IV	I
<i>Ranunculus ficaria</i>		-	-	-	-	-	-	I	I	
<i>Prunella grandiflora</i>		-	-	-	-	-	-	IV		
<i>Galeopsis pubescens</i>		-	-	-	-	-	-	III		
<i>Astragalus glycyphyllos</i>		-	-	-	-	-	-	III	I	
<i>Lysimachia nummularia</i>		-	-	-	-	-	-	III		
<i>Urtica dioica</i>		-	-	-	-	-	-	I		
<i>Galium species</i>		-	-	-	-	-	-	I		
<i>Galium mollugo</i>		-	-	-	-	-	-	I	IV	I
<i>Symphytum officinale</i>		-	-	-	-	-	-	I		
<i>Myosotis ramosissima</i>		-	-	-	-	-	-	I		
<i>Erigeron annuus</i>		-	-	-	-	-	-	I		
<i>Limodorum abortivum</i>		-	-	-	-	-	-	I		
<i>Lysimachia nemorum</i>		-	-	-	-	-	-	I		
<i>Agrimonia agrimonoides</i>		-	-	-	-	-	-	I		
<i>Viola mirabilis</i>		-	-	-	-	-	-	I		
<i>Lysimachia vulgaris</i>		-	-	-	-	-	-	I		
<i>Galium aparine</i>		-	-	-	-	-	-	I		
<i>Hieracium murorum</i>		-	-	-	-	-	-	I		II
<i>Hieracium species</i>		-	-	-	-	-	-	I	V	
<i>Origanum vulgare</i>		-	-	-	-	-	-	I	I	
<i>Juniperus communis</i>		-	-	-	-	-	-	-	II	III
<i>Genista tinctoria</i>		-	-	-	-	-	-	-		II
Other specific species the study area:										
<i>Spiraea chamaedryfolia</i>	B	-	-	+	1	-	-	II		
<i>Rubus canescens</i>		-	+	-	-	-	-	I	III	
<i>Malus sylvestris</i>		+	-	-	-	-	-	I	I	
<i>Alliaria petiolata</i>	C	3	2	2	3	3	V	I		
<i>Digitalis grandiflora</i>		3	1	+	-	+	IV	I		
<i>Fallopia convolvulus</i>		+	1	+	1	+	V	I		
<i>Dictamnus albus</i>		-	2	1	-	2	III			
<i>Carex divulsa</i>		+	-	-	+	+	III	I		
<i>Euphorbia cyparissias</i>		-	-	+	-	+	II		II	
<i>Dryopteris filix-mas</i>		+	-	-	+	-	II		I	
<i>Asplenium adiantum-nigrum</i>		+	-	-	+	-	II			
<i>Poa nemoralis</i>		1	-	+	-	-	II			
<i>Achillea millefolium</i>		-	+	-	-	-	I		I	
<i>Thymus pulegioides ssp. montanus</i>		-	-	-	+	-	I			II
<i>Rubus canescens</i>		+	-	-	-	-	I			I
<i>Prenanthes purpurea</i>		-	+	-	-	-	I			
<i>Trifolium repens</i>		-	+	-	-	-	I			
<i>Allium species</i>		-	-	-	+	-	I			
<i>Asplenium trichomanes</i>		-	-	-	I	-	I			
<i>Ranunculus bulbosus</i>		-	-	-	+	-	I			
<i>Sedum maximum</i>		-	-	-	+	-	I			
<i>Doronicum columnae</i>		-	-	-	-	+	I			
<i>Veronica montana</i>		-	-	+	-	-	I		II	

Apart from these species are accompanied the following species, with presence degree: Fukarek (*Galium aristatum* V, *Brachypodium pinnatum* V, *Carex pilosa* V, *Prunella vulgaris* IV, *Cytisus nigricans* IV, *Fragaria moschata* III, *Viola hirta* III, *Dorycnium germanicum* II, *Silene italica* II, *Agrimonia eupatoria* II, *Melittis melissophyllum* II, *Peucedanum cervaria* II, *Potentilla erecta* I, *Thymus serpyllum* I, *Aristolochia clematitis* I, *Buglossoides purpureoaeerulea* I, *Anemone nemorosa* I, *Euphorbia amygdaloides* I, *Populus tremula* I, *Tilia cordata* I, *Digitalis species* I, *Solidago virgaurea* I, *Stachys species* I, *Ulmus minor* I, *Aposotis foetida* I, *Centaurea species* I, *Chamaecytisus hirsutus* I, *Orlaya species* I, *Galeobdolon luteum* I, *Quercus polycarpa* I, *Conyza canadensis* I, *Erigeron candidus* I, *Myosotis sylvatica* I, *Persicaria maculosa* I, *Plantago media* I, *Solanum dulcamara* I, and Jovanović (*Campanula patula* V, *Hieracium bauhini* V, *Genista ovata* IV, *Chamaecytisus hirsutus* IV, *Rumex acetosella* IV, *Chamaecytisus austriacus* III, *Pteridium aquilinum* III, *Rubus hirtus* III, *Sedum cepaea* III, *Fagus moesiaca* II, *Luzula campestris* II, *Silene viridiflora* II, *Melampyrum pratense* II, *Polygala vulgaris* I, *Centaurium erythraea* I, *Epilobium lanceolatum* I, *Luzula pilosa* I, *Anthoxanthum odoratum* I, *Festuca drymeia* I, *Hypericum montanum* I, *Carex caryophyllaea* I, *Lathyrus vernus* I, *Poa angustifolia* I).



#### 4 CONCLUSIONS

The conducted phytocoenological analysis of the forests of Hungarian oak and Turkey oak in the study area show that the studied stands belong to the association *Quercetum frainetto-cerridis* Rudski (1949) Trinajstić et al. 1996, or more precisely, to its subassociation with butcher's broom *ruscetum* Jovanović et Dunjić 1951.

Syntaxonomically, this community belongs to the class *Querceto-Fagetea* Braun-Blanquet et Vlieger 1937, order *Quercetalia pubescentis* Klika 1933 and alliance *Quercion frainetto* Horvat 1954.

Along with the characteristic species of the association, the subassociation in the study area is well defined by the differential species of the subassociation *ruscetum*: *Cornus mas*, *Helleborus odoratus*, *Melica uniflora*, *Teucrium chamaedrys*, *Ruscus aculeatus*, *Physospermum cornubiense* and *Campanula persicifolia*. Synecologically, it characteristically occurs in the colline belt, in the Central European vegetation zone of thermophilic forests. It grows at elevations between 260 and 400 m and inclinations of 5-40° and on the dominantly eutric ranker on a trachyandesite base. The limiting factors of soil productivity are its depth and skeletal nature, which are conditioned by the properties of the bedrock, terrain slope and constant erosion. As a consequence, the stands are of poorer quality and lower commercial value.

As Otmanov Vis is the only area in Croatia in which this subassociation was detected, we provide the first description of this subassociation in Croatia. It should be placed under special protection owing to exceptional biological diversity and beauty of the landscape.

#### 6 REFERENCES

- [1] J. Blaženčić et al., Staništa Srbije - Priručnik sa opisima i osnovnim podacima, Institut za botaniku i botanička bašta "Jevremovac", Biološki fakultet Univerziteta u Beogradu (2005), pag. 684, Beograd.
- [2] J. Braun-Blanquet, Pflanzensoziologie, Grundzüge der Vegetationskunde (1964), pag. 865, Springer, Wien – New York.
- [3] D. Cestar, V. Hren, Z. Kovačević, J. Martinović, Z. Pelcer, Tipološke značajke šuma slavonskog gorja, Radovi Šum. instituta 39 (1979), pag. 78, Zagreb.
- [4] P. Fukarek, Prilog poznavanju nomenklature i rasprostranjenosti hrasta sladuna (*Quercus conferta* Kit. in Schult. = *Q. farnetto* Ten.), Radovi XXII. N. D. BiH. knj. 6 (1963), pag. 113-166, Sarajevo.
- [5] P. Fukarek, Sjeverozapadna granica današnje rasprostranjenosti hrasta sladuna (*Quercus conferta* Kit.). Šumarski list 88/3-4 (1964), pag. 109-123, Zagreb.
- [6] P. Fukarek, Das *Quercetum confertae hercegovinicum* im Narenta-Tal. - Angew. Pflanzensoziol. 18 (1966), pag. 37-45.
- [7] P. Fukarek, B. Fabijanić, N. Janjić, Nova nalazišta sladuna (*Quercus conferta* Kit.) u Bosni i Hercegovini, Naučni skupovi Srpske akademije nauka i umjetnosti 1 (1974), pag. 69-83, Beograd.
- [8] M. Glišić, Prilog poznavanju areala šuma hrastova cera i sladuna (*Quercetum confertae-cerris* Rudski) u severo-istočnoj Bosni. Nar. šumar 10 (1956), pag. 21-26, Sarajevo.
- [9] S. Gančev, La végétation de la zone orophyte du Rila nord-ouest. Izv. Bot. Inst. Ban, Sofia (1963), pag. 15-99.
- [10] S. M. Hennekens, J. H. J. Schaminée, TURBOVEG, a comprehensive data base management system for vegetation data, J. Veg. Sci. 12 (2001), pag. 589-591.
- [11] B. Jovanović, R. Dunjić, Prilog poznavanju fitocenoza hrastovih šuma Jasenice i okoline Beograda. Zbornik radova Instituta za ekologiju i biogeografiju Srpske akademije nauka 2 (1951), pag. 203-230, Beograd.
- [12] B. Jovanović, O klimatogenoj šumi jugoistočne Srbije, Institut za ekologiju i biogeografiju, Zbornik radova 7/6 (1956), pag. 3-35, Beograd.
- [13] B. Jovanović, Neke šumske fitocenoze severozapadne Srbije. Institut za šumarstvo i drvenu industriju, Zbornik, knjiga VI (1967), pag. 19-72, Beograd.
- [14] Matevski et al., Forest vegetation of the Galičica mountain range in Macedonia, Biološki institut Jovana Hadžija ZRC SAZU, Makedonska akademija na naukite i umetnostite (2011), pag. 200, Ljubljana.
- [15] A. Medwecka-Kornaš, R. Drenovski, Lj. Grupce, M. Mulev, The oak forest *Quercetum frainetto-cerris* on the foothills of the mountain range Galičica (Macedonia), Acta Societatis Botanicorum Poloniae 55/3 (1986), pag. 325-342.
- [16] Lj. Melovski, Lj. Grupce, M. Mulev, Nutrient Cycling *Quercetum Frainetto-Cerris Macedonicum* Ecosystem In Galičica National Park, Book, Responses of Forest Ecosystems to Environmental Changes (1992), pag. 405-416.
- [17] T. Nikolić, (ur.), Flora Croatica, baza podataka. On-line (<http://hirc.botanic.hr/fcd>). Botanički zavod, Prirodoslovno-matematički fakultet, Sveučilište u Zagrebu, Accessed december, 2015.
- [18] J. Podani, SYN-TAX 2000. Computer Programs for Data Analysis in Ecology and Systematics, User s Manual (2001), pag. 53, Budapest.
- [19] B. Puača, Ž. Najvirt, A. Miličević, Neke pedološko-florističke i gospodarske značajke šumskih sastojina na lokalitetu Otmanov Vis. Šumarski list 135/3-4 (2011), pag. 161-167, Zagreb.
- [20] I. Rudski, Tipovi lišćarskih šuma jugoistočnog dela Šumadije. Prir. Muz. Srps. Zem. 25 (1949), pag. 3-67.
- [21] V. Stupar, Đ. Milanović, J. Brujić, A. Čarni, Formalized classification and nomenclatural revision of thermophilous deciduous forests (*Quercetalia pubescentis*) of Bosnia and Herzegovina. Tuexenia 35 (2015), pag. 85-130, Göttingen, available online at [www.tuexenia.de](http://www.tuexenia.de)
- [22] A. Takšić, Geologija slavonskog gorja i Požeške kotline. U: Strbašić, M. (ur.) Požega 1277-1977 (1977), pag. 16-28, Požega.
- [23] I. Trinajstić, J. Franjić, J. Samardžić, I. Samardžić, Fitocenološke značajke šuma sladuna i cera (As. *Quercetum frainetto-cerris* Rudski 1949) u Slavoniji (Hrvatska). Šumarski list 120/7-8 (1996), pag. 299-306, Zagreb.
- [24] J. Vukelić, Šumska vegetacija Hrvatske, Sveučilište u Zagrebu, Šumarski fakultet, Državni zavod za zaštitu prirode (2012), pag. 403, Zagreb.

- [25] E. Vukićević, Šumske fitocenoze Cera. Glas. Muz. Šum. Lova 6 (1966), pag. 95–124.
- [26] J. Zelić, Preliminarne volumne tablice za hrast sladun (*Quercus frainetto* Ten), Radovi Šumarskog instituta 38/1 (2003), pag. 97–117, Jastrebarsko.
- [27] J. Zelić, Hrast sladun na Krndiji, Hrvatske šume d.o.o. - Zagreb, Uprava šuma-Podružnica Požega (2011), Požega.

PLANT BIODIVERSITY OF *SPHAGNUM*-DOMINATED MIRES IN VITOSHA NATURE PARK<sup>1</sup>DIMITROV M., <sup>2</sup>NATCHEVA R., <sup>2</sup>GANEVA A., <sup>3</sup>GYUROVA D.<sup>1</sup>Department of Dendrology, University of Forestry, Sofia, Bulgaria,<sup>2</sup>Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Sofia, Bulgaria,<sup>3</sup>Vitosha Nature Park Directorate, Sofia, Bulgaria

Corresponding author e-mail address: mariusdimitrov@abv.bg

**ABSTRACT:** Eighty-four *Sphagnum*-dominated mire complexes on the territory of Vitosha Nature Park were studied. It included spatial inventory as well as full account of plant species and plant community diversity in each mire. The distribution of mires with respect to elevation and exposure was analyzed. We found 121 vascular plant species and 89 bryophytes. This is a substantial biodiversity considering the small total area of this type of habitats in the Park (0.4% of its territory) and their patchiness. We recorded 24 species of conservation importance. The registered phytocoenoses belong to three classes, three orders, five alliances, six plant associations, three subassociations and five plant communities. The major identified threats for *Sphagnum*-dominated mires were changes in the hydrological regime due to natural drainage and climate change, as well as anthropogenic drainage and water catchment. Other negative impacts are deposition of airborne pollutants, trampling by tourists, wastes, grazing and burning. Our work highlights the role of Vitosha Nature Park for the conservation of *Sphagnum*-dominated mires in Bulgaria and the importance of proper management for their protection.

**Keywords:** plant biodiversity, phytocoenoses, diagnostic species, bryophytes, conservation, Bulgaria.

## 1 INTRODUCTION

*Sphagnum*-dominated mires on Mt Vitosha are among the most representative and in the most intact state in Bulgaria [8]. The characteristic climatic conditions and historical development of the mountain determine the formation of various plant communities that are dominated or participated by *Sphagnum* and/or other peat-forming mosses. These communities harbor a large number of species, many of which are of conservation importance. The specific hydrological regime and their vulnerability to various biotic and abiotic factors highlight a number of problems related to adequate management and conservation of *Sphagnum*-dominated mires on the territory of Vitosha Nature Park.

The first mention of *Sphagnum*-dominated mires in Bulgaria, and on Mt Vitosha in particular, was made by Adamovič [1]. He gave a brief description of vegetation and a short list of typical plant species for this type of habitat. Adamovič divided high-mountain mires into two groups – subalpine and alpine. However, he did not treat the mires on Mt Vitosha in more details.

The first extensive treatment of the bryophyte flora of the mires on Mt Vitosha was made by Podpéra [25]. He listed 62 species. Plant communities were briefly mentioned in relation to their bryophyte flora.

Brief information about the flora and vegetation of some mire complexes was provided by [2, 7, 19, 22, 30, 32, 33, 34, 37]. Ruskov [28] studied the development of spruce forests in the *Sphagnum*-mires with respect to their forestry value. Petrov [23] treated in detail four mire complexes in the spruce forests of the mountain. He presented the first comprehensive study of floristic diversity (including bryophytes), hydrology, origin and development of these mires. Hájek et al. [39] made the recent treatment of the vegetation of *Sphagnum*-dominated mires in Mt Vitosha. The authors studied the syntaxonomic diversity of 53 mires and reported 45 bryophyte and 102 vascular plant species. Information about the floristic diversity of Vitosha mires was provided also by Hájková et al. [12], Hájková & Hájek [11], and Hájek et al. [9, 10].

Vitosha mires are young formations [5, 6, 19, 23, 35]. They started to form in late Subboreal and developed intensively during the Subatlantic.

Despite the relatively large list of publications mentioning in various respects *Sphagnum*-dominated mires on Mt Vitosha, there is still missing a complete inventory of this habitat type. The aim of this study was to make a comprehensive inventory of the distribution, vegetation and flora of *Sphagnum*-dominated mires in Vitosha Nature Park, to evaluate their nature conservation state and to draw attention to the most important threats for this extremely vulnerable and important habitat type.

## 2 MATERIAL AND METHODS

## 2.1 Study site

Vitosha Nature Park occupies the larger part of Mt Vitosha (Fig. 1). This is the oldest designated national Park in Bulgaria. Mt Vitosha is situated in the south-western part of Bulgaria. The average elevation is 1317 m a.s.l., the highest peak is Cherni vrah (2290 m a.s.l.). The average precipitation is ca 1115 mm (841 mm at 1300 m and 1175 mm at 2286 m a.s.l.), the average temperature is ca. 3°C (7°C at 1300 m and 0,3°C at 2286 m a.s.l.). Mt Vitosha is a relatively young dome-shaped mountain containing an older upper Cretaceous crystalline plutonic core, surrounded by volcanic and volcano-sedimentary rocks [36]. Characteristic for the mountain is the presence of large plateau morphostructures that were further modelled by denudation and cryogenic processes. These structures harbor the largest mire complexes in Mt Vitosha and are the major water source of the mountain.

## 2.2 Spatial inventory

For the initial spatial inventory we used published data as well as the information for the distribution of habitat type 7140 and 91D0 for Mt Vitosha from the project on the Natura 2000 network in Bulgaria “Mapping and assessment of the conservation status of habitats and species. Phase I”. A preliminary map was drawn which was used for the field trips. At each site the geographical coordinates and elevation of a central point were measured via GPS. This data coupled with current


orthophoto images was used to draw polygons of the studied sites.

### 2.3 Inventory of plant diversity

At each visited polygon in area with typical mire vegetation was selected a relevé of 25 m<sup>2</sup> for description of plant communities. For communities along springs and brooks the size of relevés was 16 m<sup>2</sup>. Additionally, floristic diversity was studied in transects along the entire site and the recorded additional species were assigned an abundance score of “+”.

The nomenclature of vascular plants follows [4], of mosses [13], and of liverworts [27].

The classification of plant communities was done using the Braun-Blanquet's approach [3, 38]. The abundance/ dominance of taxa was evaluated using an extended nine-grade Braun-Blanquet scale [14]. Cluster analysis was made using the software Syn-Tax [25]. We applied UPGMA, floristic similarity was calculated using the Penrose index. Some relevé were moved manually considering common diagnostic species. The identified clusters were compared with vegetation descriptions provide by Hájek et al. [9, 39]. If general floristic composition, diagnostic, constant and dominating species coincided, the vegetation clusters were assigned to the respective syntaxa and syntaxonomic categories.



**Figure 1:** Map of the study area

## 3 RESULTS


### 3.1 Distribution of Sphagnum-dominated mires

We studied a total 84 mires on Mt Vitosha (Fig. 1, Annex I). All of them were situated within Vitosha Nature Park. They are distributed in the altitudinal range of 1230-2240 m. The largest number of mires is at 1700-1850 m and above 1900 m a.s.l. with a northern component of exposure (Fig. 2 and Fig. 3). Fewer peatlands are located on the southern slopes due to the prevalence of calcareous bedrock, higher insulation, and lower moisture. These conditions reflect on the vegetation composition and diversity. The total area of all studied mires is 73,3 ha. The average size is 0,9 ha (minimum 0,7 ha, maximum 17,8 ha).


### 3.2 Plant communities

Plant communities were described at 65 mire polygons. The syntaxonomic analysis revealed that the studied communities belong to three classes, three orders, five alliances, six plant associations, three subassociations and five plant communities (Annex I). Additionally were found plant groups belonging to the rangless category "community". Two of these groups

included five and three relevés, the remaining three were recorded at a single relevé.



**Figure 2:** Distribution of Sphagnum-dominated mires in Vitosha Nature Park according to elevation



**Figure 3:** Distribution of Sphagnum-dominated mires in Vitosha Nature park according to exposure

Of the syntaxa described by Hajek at al. [39] only the *Drepanocladetum exannulati* Krajina 1933 was not recorded. The authors point that this association, documented from the Vitosha Mt only by one small relevé, represents the oligotrophic mires dominated by sedges and brown mosses indicating initial, albeit stable, and strongly waterlogged stages of mire formation.

Other communities represented by one locality in Vitosha Mt and distributed mainly in other mountains in Bulgaria [9], were also not identified: ass. *Carici echinatae-Sphagnetum* Soó 1944, ass. *Caricetum nigrae* Braun 1915 subass. typicum, var. *Sphagnum teres*, ass. *Dactylorhiza cordigerae-Eriophoretum latifolii* Hájek et al. 2008.

The association *Primula exiguae-Caricetum echinatae* Roussakova 2000 was found at two relevés. Here are found the diagnostic for this association *Primula \*exigua*, *Sphagnum platyphyllum*, *Pinguicula balcanica*, and *Pseudorchis frivaldii*. Bryophytes were represented mostly by the genera *Sphagnum*, *Warnstorfia*, and *Polytrichum*.

A large part of mire complexes in Mt Vitosha in the altitudinal range of 1750–1900 m a.s.l. belong to ass. *Cirsio heterotrichi-Caricetum nigrae* (Soo 1957) Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005 [39]. During the present study, this association was found at 18 localities. All of the diagnostic species of the association were registered. Among mosses the largest participation had *Sphagnum warnstorffii*, *Sphagnum capillifolium*, *Sphagnum flexuosum*, *Sphagnum contortum*, *Sphagnum palustre*, *Sphagnum teres*, *Aulacomnium palustre*,



*Polytrichum commune*, *Warnstorfia exannulata*, *Warnstorfia sarmentosa*, etc. All known successional stages, described as subassociations were present. Subass. *sphagnetosum subsecundi* makes successional and syntaxonomic transition with ass. *Primulo exiguae-Caricetum echinatae*. It was found at 11 localities. The major diagnostic species *Sphagnum subsecundum* was well represented while the remaining diagnostic species *Carex echinata*, *Eriophorum angustifolium*, and *Gentianella bulgarica* had lower constancy and abundance as compared to the original diagnosis of the association. The typical subassociation was present at four relevés. Subass. *eriphoretosum vaginatae* was reported at four relevés and represents the most advanced successional stages within the association.

The largest number of relevés in the present study belonged to ass. *Bruckenthalio-Sphagnetum capillifolii* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005. It represents a more advanced successional stage characterized by shrub and semi-shrub species such as *Juniperus sibirica*, *Vaccinium vitis-idaea*, *Vaccinium myrtillus*, *Vaccinium uliginosum*, *Bruckenthalia spiculifolia*, as well as a number of species typical for wet meadows – *Deschampsia caespitosa*, *Agrostis capillaris*, *Alchemilla vulgaris* agg., *Geum coccineum*, *Allium sibiricum*, *Nardus stricta*, etc. This association was the optimum in the distribution of *Sphagnum capillifolium*. High constancy and abundance had also *Sphagnum platyphyllum*, *Sphagnum subsecundum*, *Warnstorfia exannulata*, *Warnstorfia sarmentosa*, etc. This association was distributed mostly at 1900–2200 m a.s.l.

Association *Angelico pancicii-Calthetum laetae* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005 develops around springs and small water courses. Two relevés belonged to this association. Since the relevés were chosen in order to reveal bryophyte diversity, the abundance of *Caltha palustris* and *Angelica pancicii* was relatively low. Besides the diagnostic *Brachythecium rivulare* and *Rhizomnium punctatum* were detected 10 bryophyte species albeit with lower abundance.

To the spring vegetation belongs the association *Saxifragetum stellaris* Deyl 1940. It was represented at four relevés at 1775–1860 m a.s.l. Of the diagnostic species were found *Saxifraga alpigena*, *Philonotis seriata*, and *Soldanella montana*. Other differential species were *Scapania undulata* and *Oncophorus virens*.

Besides the typical mire and spring vegetation types, *Sphagnum* mosses develop in hygrophilic communities of order *Molinietalia*. Such were the phytocoenoses of ass. *Scirpetum sylvatici* Ralski 1931, which was found by Hájek et al. [39] as well as in the current study. Differential for this association are *Scirpus sylvaticus*, *Lysimachia vulgaris*, *Scutellaria alpina*, and *Sphagnum fimbriatum*. Other bryophytes are *Dicranella palustris*, *Sphagnum girgensohnii*, *Sphagnum squarrosum*, *Polytrichum commune*, and *Philonotis fontana*.

At some sites were observed phytocoenoses characteristic for advanced successional stages and indicating more mesic conditions – lesser representation of *Sphagnum* mosses, higher proportion of mesophilic grasses, shrubs and trees. The coenoses of comm. with *Holcus mollis* had relatively high abundance and constancy of species from *Caricetalia fuscae* (*Carex nigra*, *Carex echinata*, *Eriophorum vaginatum*, *Eriophorum latifolium*), as well as species from *Molinietalia* (*Deschampsia caespitosa*, *Angelica pancicii*, *Juncus effusus*, *Scirpus sylvaticus*, *Juncus conglomeratus*,

*Potentilla erecta*, *Filipendula ulmaria*, etc.). At better-drained places were developed facies dominated by or with higher abundance of *Holcus mollis*, *Calamagrostis arundinacea*, *Rubus idaeus*, *Pinus sylvestris*, and *Alnus glutinosa*. These species indicate a succession towards formation of shrub and forest vegetation types. Differential species were *Holcus mollis* and *Calliergonella cuspidata*. Of bryophytes often occur *Sphagnum warnstorffii*, *Sphagnum subsecundum*, *Scapania undulata*, *Warnstorfia exannulata*, *Sphagnum flexuosum*, *Sphagnum contortum*, etc. Similar species composition had the coenoses of comm. of *Festuca rubra*, where high constancy and relatively high abundance had *Festuca rubra* and *Agrostis capillaris*.

Community of *Carex rostrata* was found at one locality. It developed at a water saturated site at the edge of permanent water pool. Differential and dominant species were *Carex rostrata* and *Carex curta*. Bryophytes with highest abundance were *Sphagnum teres*, *Sphagnum russowii*, *Sphagnum inundatum*, *Polytrichum commune*, *Warnstorfia exannulata*, *Aulacomnium palustre*, and *Philonotis fontana*.

Community of *Carex nigra* and *Luzula sudetica* is transitional between ass. *Primulo exiguae-Caricetum echinatae* and ass. *Saxifragetum stellaris*. Major dominants were *Carex nigra*, *Sphagnum capillifolium*, *Luzula sudetica*, *Carex echinata*, and *Sphagnum flexuosum*. Higher participation had also *Sphagnum contortum*, *Sphagnum inundatum*, *Scorpidium cossonii*, *Philonotis seriata*, *Rhizomnium punctatum*, etc.

Community of *Fissidens adianthoides* and *Hamatocaulis vernicosus* was described at one site with lower cover of vascular plants. A large number of bryophytes without a clear dominant also participated in this community. Differential species were *Fissidens adianthoides* and *Hamatocaulis vernicosus*. The optimum development of the latter species is where herbaceous vascular plants are absent or their abundance is low [31].

### 3.3 Plant diversity and conservation species

In *Sphagnum*-dominated mires on the territory of Vitosha Natural Park were found a total of 210 plant species (Table I). Of them 121 were vascular plants and 89 were bryophytes. Nine vascular plants and 15 bryophytes are of conservation importance (Table I). Two species are Critically Endangered, three Endangered, 17 Vulnerable, one Near Threatened, and one Data Deficient [18, 24, 21]. Further three species are included in Annex II and 22 species are included in Annex 2a of Bulgarian Biodiversity Act. One species is included in Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora.

Further five species are Balkan endemics or sub-endemics (*Angelica pancicii*, *Senecio pancicii*, *Gentianella bulgarica*, *Crocus veluchensis*, and *Cirsium heterotrichum*). Ten bryophyte species (*Sphagnum auriculatum*, *Sphagnum compactum*, *Sphagnum subnitens*, *Calypogeia neesiana*, *Campylium polygamum*, *Plagiomnium ellipticum*, *Scorpidium cossonii*, *Scorpidium revolvens*, *Drepanocladus polygamum*, and *Dichelyma falcatum*) are new to the flora of Mt. Vitosha.

## 4 DISCUSSION

### 4.1 Importance of Vitosha Nature Park for the conservation of *Sphagnum*-dominated mires

**Table I:** Vascular plant and bryophyte species of conservation importance in *Sphagnum*-dominated mires on the territory of Vitosha Nature Park. Abbreviations: CR – Critically Endangered, EN – Endangered, VU – Vulnerable, NT – Near Threatened, DD – Data Deficient, RL – Red list, RDB – Red Data Book of Republic of Bulgaria, BA – Biodiversity Act of Bulgaria.

№	Species	Category	National conservation status	Other
<b>Vascular plants</b>				
1.	<i>Angelica pancici</i>	VU	RL	Balkan endemic
2.	<i>Drosera rotundifolia</i>	VU	RL	
3.	<i>Potentilla palustris</i>	VU	RL	
4.	<i>Pseudorchis albida</i>	VU	RL, CITES	
5.	<i>Pseudorchis frivaldii</i>	VU	RL, CITES	
6.	<i>Salix pentandra</i>	CR	RL, RDB, BA	Glacial relict
7.	<i>Senecio pancici</i>	NT	RL	Balkan endemic
8.	<i>Swertia perennis</i>	EN	RL, RDB	Glacial relict
9.	<i>Utricularia minor</i>	EN	RL, RDB, BA	
<b>Bryophytes</b>				
1.	<i>Bryum weigelii</i>	VU	RL	
2.	<i>Calypogeia sphagnicola</i>	VU	RL	
3.	<i>Drepanocladus polygamum</i>	VU	RL	
4.	<i>Cephalozia connivens</i>	VU	RL	
5.	<i>Cephalozia pleniceps</i>	DD	RL	
6.	<i>Cephaloziella hampeana</i>	CR	RL, RDB	
7.	<i>Hamatocaulis vernicosus</i>	VU	RL, RDB, BA, Directive 92/43	
8.	<i>Meesia uliginosa</i>	VU	RL	
9.	<i>Plagiothecium denticulatum</i> var. <i>undulatum</i>	VU	RL	
10.	<i>Riccardia incurvata</i>	VU	RL	
11.	<i>Riccardia multifida</i>	VU	RL	
12.	<i>Scorpidium revolvens</i>	VU	RL, RDB	
13.	<i>Sphagnum fallax</i>	VU	RL	
14.	<i>Sphagnum subnitens</i>	VU	RL, RDB	
15.	<i>Tomentypnum nitens</i>	EN	RL, RDB	

Vitosha Nature Park is an important area for the conservation of *Sphagnum*-dominated mires and their plant species and community diversity in Bulgaria. Mt Vitosha is the richest of *Sphagnum*-mosses phytogeographic region in Bulgaria with 23 species. Despite the restricted and scattered distribution and small size, this type of habitat is characterized by a considerable plant species and plant community diversity. For the first time Stefanoff [32] paid attention to the process of drying out of Vitosha mires. He suggested that measures must be undertaken for their restoration in the light of their importance for the water regime of the mountain. In addition to their role for biodiversity conservation, Vitosha mires provide a number of valuable ecosystem services. The most important of them are related to the water holding and water regulation capacity of *Sphagnum* cover and the peat layer. The rain and snowmelt water penetrate slowly in the peat and thus it supports the even river flow. Most rivers of Mt Vitosha have their sources in *Sphagnum*-dominated mires.

The most important threat to the mires in Mt Vitosha is drying. This is mostly a natural process due to two factors: climate change and lowering of water table due to water erosion of stream beds. The natural processes were enhanced by building of a number of water catchments for drinking water in the spring areas of many mires in the 1950-s and 1960-s. Drying has a twofold effect. On the one hand, it suppresses the development of the major habitat component – *Sphagnum* mosses. On the other hand drying stimulates the development of more competitive vascular plants. The overall effect is gradual displacement of the typical mire species with species of mesophilic meadows. The process of displacement is

enhanced by increased nutrient input from air pollution with nitrogen and sulphur from the nearby urban areas (the city of Sofia and the industrial area of town Pernik) which was considerable during the middle of XIX by the end of XX century.

Proper management of *Sphagnum*-dominated mires on the territory of the Park is needed in order to protect this unique and rich type of habitat. This include mostly preventive measures: careful consideration when new water catchments or infrastructures are built in order to prevent disturbance of water regime, taking measures to avoid passing of tourists through mires, and last but not least raising awareness for the importance and vulnerability of *Sphagnum*-mires. Active restoration of some sites with disturbed water regime may also be undertaken. Such recommendation could be found also in the report for habitat 7140 Transition mires and quaking bogs (distribution and ecological status) as a result of above mentioned project on mapping habitats and species in Natura 2000 sites in Bulgaria (<http://natura2000.moew.government.bg>).

#### 4.2 Comments on some species of conservation importance

*Drosera rotundifolia*. The species was reported for Mt. Vitosha in 1883 [20]. Since then there are occasional reports [e.g. 23] but the general view was that it is rare. Our study showed that *D. rotundifolia* is a relatively regular member of *Sphagnum*-dominated mires in Mt Vitosha having being found at 18 sites. All of them are situated in the altitudinal range of 1300-1886 m a.s.l. At most sites the populations are numerous and in very good state.

*Potentilla palustris*. In Mt. Vitosha it has been found for the first time in 1930 [37], and has not been reported since then for the flora of the mountain. We found it at three sites. The populations were represented by a small number of individuals with scattered shoots often barely to be seen above the *Sphagnum* cover. Only at one site the plants were observed with flowers.

*Sphagnum subnitens*. This is the first report of the species for Mt. Vitosha. The species is known only from Pirin [29] and Rila Mts [17]. On Mt. Vitosha *S. subnitens* was found at one site, represented by few small patches.

*Drepanocladus polygamum*. This is the first report of the species for Mt. Vitosha. The species is reported only from Pirin Mt [15] but it was not supported by herbarium specimen. The species has not been observed in Bulgaria since 1966.

*Hamatocaulis vernicosus*. The occurrence of *H. vernicosus* on Mt. Vitosha has been doubtful. It was reported by Podpéra [26] for loc. Dragalevsko blato, which is located around the springs of Dragalevska river at ca. 1800 m a.s.l. and is today called loc. Kapaklivets. The attempts of the authors to verify the occurrence of the species at this locality proved unsuccessful. Since the conditions at the site differ from the conditions, where *H. vernicosus* is found in Bulgaria at present and is not supported by herbarium specimen, this report was treated as probable misidentification. The location of *H. vernicosus* on the south-western slopes of Mt. Vitosha in the current study is the first confirmed occurrence of the species in Vitosha Nature Park.

*Dichelyma falcatum*. This species was recently added to the list of bryophytes in Bulgaria. It was found for the first time in Rila Mts at 2300 m a.s.l. [16]. This is the second location of *D. falcatum* in Bulgaria and the first report for Mt. Vitosha. The species had similar ecological requirements on Mt. Vitosha – high elevation (2230 m a.s.l.) growing on acidic rocks in streams that dry out in late summer flowing through *Sphagnum*-mires. The species is not Red-listed because it was found in Bulgaria after the publication of the most recent Red list [18]. However, since its discovery it proved to be rare and grows at extremely vulnerable sites. It will therefore be considered in the next evaluation of Bulgarian bryophytes and will be assigned to a threat category.

## 5 ACKNOWLEDGMENT

This study was done within project DIR-5113326-4-98 “Actions for sustainable management of Nature Park Vitosha” financed by European Regional Development Fund and the national Operational Programme “Environment” 2007 – 2013.

## 6 REFERENCES

- [1] L. Adamovič, 1909. Vegetationsverhältnisse der Balkanländer (Mösische Länder) umfassend Serbien, Altserbien, Bulgarien, Ostrumelien, Nordthrakien und Nordmacedonien. Vegetation der Erde, XI. Leipzig: Engelmann. (1909), pag. 567.
- [2] I. Bondev, T. Meshinev, N. Andreev, and M. Lyubenova, The vegetation in “Torfeno branishte” reserve, Vitosha Mt. In: V. Velchev (ed.), National Theoretic Conference for Environmental Conservation and Restoration Vol. 1 (1982), pag. 80-84. Sofia (in Bulgarian).
- [3] J. Braun-Blanquet, 1964. Pflanzensoziologie. Grundzüge der Vegetationskunde. 3rd ed. Springer-Verlag, Wien, AT. (1964), pag. 865.
- [4] D. Delipavlov and I. Cheshmedzhiev (Eds.), Key to the Plants in Bulgaria. Acad. Press Agrarian Univ., Plovdiv. (2003), pag. 591. (in Bulgarian)
- [5] L. Filipovich, Palynological investigation of a peat bog above “Kumata” chalet, Vitosha Mountain. Fitologiya 21. (1982), pag. 27-33. (in Bulgarian)
- [6] L. Filipovitch, Palynological investigation of the peat bog under the Reznovete peaks in Vitosha Mountain. Nauka za Gorata 18(3). (1981), pag. 9-15. (in Bulgarian)
- [7] T. Georgieff, Phytogeographische Skizze des Vitošageirges. Jahrbuch der Universität Sofia, Landwirtschaftliche Fakultät, B, 6. (1928), pag. 179-209.
- [8] C. Gushev, D. Dimitrova, and S. Tzoneva, NATURA 2000 – European Ecological Network in Vitosha Mountain. EcoArc, Sofia. (2005).
- [9] M. Hájek, P. Hájková, and I. Apostolova, New plant associations from Bulgarian mires. Phytologia Balcanica 14. (2008), pag. 377-399.
- [10] M. Hájek, P. Hájková, D. Sopotlieva, I. Apostolova, and N. Velez, The Balkan wet grassland vegetation: a prerequisite to better understanding of European habitat diversity. Plant Ecology 195. (2008), pag. 197-213.
- [11] P. Hájková and M. Hájek, Sphagnum distribution patterns along environmental gradients in Bulgaria. Journal of Bryology 29. (2007), pag. 18-26.
- [12] P. Hájková, M. Hájek, and I. Apostolova, Diversity of wetland vegetation in the Bulgarian high mountains, main gradients and context-dependence of the pH role. Plant Ecology 184. (2006), pag. 111-130.
- [13] M. O. Hill, N. Bell, M. A. Bruggeman-Nannenga, M. Brugués, M. J. Cano, J. Enroth, K. I. Flatberg, J.-P. Frahm, M. T. Gallego, R. Garilleti, J. Guerra, L. Hedenäs, D. T. Holyoak, J. Hyvönen, M. S. Ignatov, F. Lara, V. Mazimpaka, J. Munõz, and L. Söderström, An annotated checklist of the mosses of Europe and Macaronesia. Journal of Bryology 28. (2006), pag. 198-267.
- [14] E. van der Maarel, Transformation of cover-abundance in phytosociology and its effects on community similarity. Vegetatio 39(2). (1979), pag. 97-114.
- [15] J. Mickiewicz, I. Rejment-Grochowska, and D. Sobotka, Résultats des recherches bryologiques en Bulgarie. Acta Societatis Botanicorum Poloniae 35(1). (1966), pag. 111-127.
- [16] R. Natcheva, Dichelyma falcatum: a new aquatic moss to the bryophyte flora of Bulgaria. Phytologia Balcanica 13(3). (2007), pag. 311-312.
- [17] R. Natcheva, R. Records 9-20. In: R. Natcheva (compiler), New bryophyte records in the Balkans: 5. Phytologia Balcanica 14(3). (2008), pag. 426-428.
- [18] R. Natcheva, A. Ganeva, and G. Spiridonov, Red list of the bryophytes in Bulgaria. Phytologia Balcanica 12(1). (2006), pag. 55-62.
- [19] N. Nedeva, Contemporary development of the peat bogs in the Vitosha Mountain. Fitologiya 1. (1975), pag. 34-46. (in Bulgarian)

- [20] J. Pančić, *Elementa ad floram principatus Bulgariae*. Beograd. (1883).
- [21] D. Peev, and al. (eds), *Red Data Book of Republic of Bulgaria*. Vol. I. Plants and fungi. BAS & MOEW, Sofia. (2015).
- [22] I. Penev, *Grass Associationen über dem Waldgürtel der Witoscha*. *Godishnik na Sofiiskiia Universitet, Biologicheski Fakultet*. Kniga 2, 58. (1964), pag. 59-94. (in Bulgarian)
- [23] S. Petrov, *Quellmoore in den Fichtenwäldern des Vitosha-Gebirges*. *Izvestiya na Botanicheskaya Institut*. Sofia 5. (1956), pag. 296-346. (in Bulgarian)
- [24] A. Petrova and V. Vladimirov, *Red List of Bulgarian vascular plants*. *Phytologia Balkanica* 15(1). (2009), pag. 63-94.
- [25] J. Podani, *SYN-TAX 2000*. Computer Program for Data Analysis in Ecology and Systematics. User`s Manual. Scientia Publ., Budapest. (2001), pag. 53.
- [26] J. Podpéra, *Ein Beitrag zu der Kryptogamenflora der bulgarischen Hochgebirge*. Beihefte zum Botanischen Centralblatt, Abt. 2, 28. (1911), pag.173-224.
- [27] R. M. Ros, V. Mazimpaka, U. Abou-Salama, M. Aleffi, T. L. Blockeel, M. Brugués, M. J. Cano, R. M. Cros, M. G. Dia, G. M. Dirkse, W. El Saadawi, A. Erdağ, A. Ganeva, J. M. González-Mancebo, I. Herrnstadt, K. Khalil, H. Kürschner, E. Lanfranco, A. Losada-Lima, M. S. Refai, S. Rodríguez-Nuñez, M. Sabovljević, C. Sérgio, H. Shabbara, M. Sim-Sim, and L. Söderström, *Hepatics and Anthocerotae of the Mediterranean, an annotated checklist*. *Cryptogamie, Bryologie* 28(4). (2007), pag. 351-43.
- [28] M. Ruskov, *Sphagnummoore in den Fichtenwälder des Witosha-Gebirges*. *Forstwirtschaft Gedanke* 3-4. (1934), pag. 147-156. (in Bulgarian)
- [29] J. Šmarda, *Complements à la flore muscinale de la Bulgarie*. *Revue bryologique et lichénologique* 37(1). (1970), pag.33-46.
- [30] R. Soó, *Pflanzengesellschaften aus Bulgarien*. I. – *Annales Universitatis Scientiarum Budapestinensis de Rolando Eötvös Nominatae, Sectio biologica* 1. (1957), pag. 231-239.
- [31] T. Štechová, M. Štech, and J. Kučera, *The distribution of *Hamatocaulis vernicosus* (Mitt.) Hedenäs (Calliergonaceae, Bryophyta) in the Czech Republic*. *Bryonora* 49. (2012): pag. 5-16.
- [32] B. Stefanoff, *La couverture végétale de Vitocha comme objet de l'élevage, de la defense et de l'utilisation*. *Annuaire de L'Université de Sofia, Livre 2*, 17. (1939), pag. 225-254.
- [33] N. Stojanoff, *Die vegetaiton des Bistrizaer Naturschutzgebietes*. *Sbornik na sjusa sa saschtschita na prirodata* 2. (1939) pag. 13-46. (in Bulgarian)
- [34] B. Iwanow, Al. K. Drenowski. *Über die Pflanzenformationen der alpinen Region des Witoschaberges in Bulgarien*. *Allgemeine Botanische Zeitschrift* 11-12. (1910), pag. 1-4.
- [35] N. Stojanoff and T. Georgieff, *Pollenanalytische Untersuchungen auf dem Vitoša-Gebirge*. *Spisanie BAN* 47. (1934), pag. 73-104.
- [36] T. Tsankov, *Rock structure*. In: A. Popov (ed.), *The Natural Bounty of Vitosha Nature Park*. Vitosha Nature Park Directorate, Sofia. (2014), pag. 17-24. (in Bulgarian)
- [37] I. Urumov, *Flora von Vitoša Planina*. *Sbornik na Bălgarskata Akademiya na Naukite* 26. (1930), pag.1-143. (in Bulgarian)
- [38] V. Westhoff and E. van der Maarel, *The Braun-Blanquet approach*. 2nd. Ed. In: R. Whittaker (ed.), *Classification of plant communities*. Junk, The Hague. (1978), pag. 287-399.
- [39] M. Hájek, R. Tzonev, P. Hájková, A. Ganeva, and I. Apostolova, *Plant communities of the subalpine mires and springs in the Vitosha Mt*. *Phytologia Balcanica* 11(2). (2005), pag. 193-20.



**Annex I:** Area, elevation and geographic coordinates of the investigated mire in Vitosha Nature Park. Mire numbers correspond to Annex II.

Nr.	Area (ha)	Elevation (m a.s.l.)	Longitude	Latitude
1	0.05	1800	23.291063	42.58905
2	0.01	2240	23.282532	42.562478
3	0.06	2235	23.281997	42.562313
4	0.23	2225	23.28138	42.561382
5	0.16	2225	23.28331	42.559605
6	1.73	2150	23.289183	42.551828
7	0.56	2150	23.2935	42.546433
8	0.51	1957	23.279205	42.58437
9	0.28	2130	23.293127	42.544017
10	3.65	2130	23.293842	42.542563
11	0.02	2060	23.29362	42.55549
12	0.31	2060	23.294455	42.555632
13	6.1	1790	23.256981	42.59076
14	0.1	1824	23.232778	42.581372
15	0.01	1952	23.277223	42.5847
16	0.28	1822	23.234676	42.58076
17	1.59	1814	23.230481	42.581557
18	0.04	1727	23.233761	42.587235
19	0.99	1510	23.241319	42.602244
20	0.03	1833	23.246824	42.58457
21	0.25	1887	23.246224	42.58239
22	11.07	1926	23.268523	42.58455
23	0.04	1885	23.243607	42.58194
24	0.11	1787	23.249498	42.58665
25	0.03	1775	23.249945	42.58826
26	0.27	1774	23.249263	42.588952
27	0.02	1755	23.240201	42.58725
28	0.15	1725	23.238814	42.58735
29	0.14	1526	23.239467	42.60144
30	0.4	1545	23.237193	42.59924
31	0.11	1656	23.223465	42.58983
32	0.16	1700	23.224869	42.58817
33	0.33	1776	23.24276	42.58618
34	0.02	1787	23.24408	42.5863
35	0.57	1780	23.244945	42.58667
36	2.73	1793	23.246718	42.58666
37	0.01	1787	23.266442	42.60308
38	0.01	1786	23.266137	42.60312
39	0.01	1790	23.265989	42.60296
40	0.07	1794	23.265603	42.60301
41	0.01	1789	23.265068	42.60321
42	0.18	1785	23.263199	42.6045
43	3.86	1805	23.263971	42.59937
44	5.69	1830	23.284095	42.58901
45	0.52	1840	23.285689	42.59171
46	0.04	1818	23.283812	42.59409
47	0.08	1806	23.282432	42.59687

Nr.	Area (ha)	Elevation (m a.s.l.)	Longitude	Latitude
48	0.11	1815	23.281437	42.59702
49	17.08	1840	23.278615	42.59497
50	0.06	1815	23.296888	42.55645
51	0.04	2010	23.295881	42.55655
52	0.02	2030	23.295526	42.55583
53	0.02	1940	23.307565	42.55028
54	0.01	1950	23.307448	42.54984
55	0.09	1920	23.309799	42.54926
56	0.12	1920	23.310269	42.54851
57	0.22	2140	23.29859	42.549707
58	4.29	1970	23.265713	42.55256
59	0.96	1990	23.261705	42.55737
60	0.04	1573	23.224503	42.56128
61	0.03	1701	23.24201	42.5579
62	0.02	1705	23.2438	42.55762
63	0.03	1706	23.244265	42.55748
64	0.11	1711	23.247182	42.55536
65	0.08	1718	23.247104	42.55622
66	0.01	1230	23.245056	42.50663
67	0.06	1318	23.252863	42.49297
68	0.06	1348	23.262691	42.50016
69	0.04	1879	23.283993	42.58691
70	0.17	2117	23.273196	42.56725
71	0.1	2046	23.270814	42.57024
72	1.62	2040	23.268327	42.56941
73	0.02	2030	23.266009	42.56885
74	0.48	2052	23.259509	42.56709
75	0.02	2051	23.257008	42.56617
76	0.06	2038	23.256381	42.5665
77	0.22	1994	23.248799	42.56785
78	0.81	1996	23.253211	42.56966
79	0.26	1994	23.251895	42.57214
80	0.49	1976	23.25419	42.57277
81	1.62	1954	23.259068	42.57493
82	0.33	1910	23.26425	42.57935
83	0.04	1912	23.264295	42.58301
84	11.07	1926	23.268523	42.58455
85	0.7	1830	23.303	42.571

**Annex II:** Synoptic table of the phytocoenoses from Vitosha Mts. Mire numbers correspond to Annex I. 1. Syntaxa: 1. ass. *Primula exiguae*-*Caricetum echinatae* Roussakova 2000; 2. ass. *Cirsio heterotrichi*-*Caricetum nigrae* (Soo 1957) Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005 subass. *sphagnetosum subsecundi* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005; 3. ass. *Cirsio heterotrichi*-*Caricetum nigrae* (Soo 1957) Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005 subass. *typicum* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005; 4. ass. *Cirsio heterotrichi*-*Caricetum nigrae* (Soo 1957) Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005 subass. *eriphoretosum vaginatae* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005; 5. ass. *Bruckenthalio-Sphagnetum capillifolii* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005; 6. ass. *Angelico pancicii*-*Calthetum laetae* Hájek, Tzonev, Hájková, Ganeva and Apostolova 2005; 7. comm. with *Holcus mollis*; 8. ass. *Saxifragetum stellaris* Deyl 1940; 9. comm. of *Festuca rubra*; 10. comm. of *Carex rostrata*; 11. ass. *Scirpetum sylvatici* Ralski 1931; 12. comm. of *Carex nigra* and *Luzula sudetica*; 13. comm. of *Fissidens adianthoides* and *Hamatocaulis vernicosus*.

Syntaxa		1	2	3	4	5	6	7	8	9	10	11	12	13
Number of relevés		2	10	4	4	26	2	5	5	3	1	1	1	1
Number of species		75	110	103	64	147	42	78	75	45	24	12	26	26
Taxa	Mire number	CONSTANCY/ABUNDANCE												
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>Primula *exigua</i>	1, 3-6, 10, 12, 16, 25, 27, 38, 39, 49, 57, 64, 65, 75, 82	V <sup>+-1</sup>	II <sup>+</sup>			III <sup>+</sup>								
<i>Sphagnum platyphyllum</i>	2, 3, 9-11, 16, 25, 49, 50, 69-71, 74, 76, 78, 79	V <sup>2m-2b</sup>	I <sup>2a</sup>			III <sup>2b-3</sup>								
<i>Carex nigra</i>	1, 2, 4, 6-8, 10-13, 15-17, 20, 21, 27-29, 32-35, 37, 43, 57, 61, 63-65, 69-71, 73-75, 78, 82	V <sup>+-1</sup>	V <sup>+-5</sup>	V <sup>+-3</sup>	V <sup>+-3</sup>	V <sup>+-4</sup>	V <sup>2b-3</sup>	III <sup>2a-2b</sup>			+		4	
<i>Molinia caerulea</i>	1, 13, 14, 16, 17, 19-23, 25, 27, 28, 32, 36, 38, 39, 42-44, 49, 50, 60, 69, 71	V <sup>+</sup>	IV <sup>+-4</sup>	V <sup>2m-3</sup>	IV <sup>+-4</sup>	I <sup>+-3</sup>								+
<i>Succisa pratensis</i>	1, 13, 14, 16-23, 25, 27, 29, 32, 33, 36, 38, 39, 42-44, 60, 63, 65, 69, 77, 81	III <sup>+</sup>	V <sup>+-2b</sup>	V <sup>+-2b</sup>	V <sup>+-3</sup>	II <sup>+-2b</sup>		I <sup>2a</sup>			+			+
<i>Cirsium heterotrichum</i>	1, 3, 4, 6-8, 10, 13, 14-17, 20-22, 25, 27, 29, 32, 34, 35, 37, 42, 43, 49, 50, 57, 61, 64, 65, 69-71, 75, 77-79, 81, 82, 85	V <sup>+</sup>	IV <sup>+</sup>	IV <sup>+-2a</sup>	II <sup>+</sup>	IV <sup>+-3</sup>	V <sup>+</sup>	I <sup>2a</sup>	I <sup>+</sup>	IV <sup>+</sup>	+			
<i>Potentilla erecta</i>	4, 6-10, 12-21, 23, 25, 27-29, 32-39, 42-44, 49, 50, 57, 60, 61, 64, 65, 67-71, 74-79, 81, 82, 85	V <sup>+</sup>	V <sup>+-2b</sup>	V <sup>+-2b</sup>	III <sup>2b-3</sup>	V <sup>+-3</sup>	V <sup>+</sup>	IV <sup>+-2a</sup>	IV <sup>+</sup>	V <sup>2a-3</sup>	+			+
<i>Dactylorhiza cordigera</i> s.l.	1, 5, 10, 11, 13, 16, 17, 21, 23, 25, 27, 29, 32, 33, 37-39, 42, 43, 49, 60, 61, 62, 63, 65, 69, 75, 77, 82	V <sup>+</sup>	V <sup>+-1</sup>	II <sup>1</sup>	III <sup>+</sup>	II <sup>+-2a</sup>		III <sup>+-1</sup>			+			+
<i>Sanguisorba officinalis</i>	1, 7, 13, 14-18, 20, 23, 25, 27, 29, 62, 63, 69, 71, 74, 76, 81, 82, 78	III <sup>+</sup>	II <sup>+</sup>	IV <sup>+-2b</sup>	III <sup>+</sup>	III <sup>+-2a</sup>		I <sup>+</sup>						
<i>Sphagnum warnstorffii</i>	1, 2, 4, 9, 13, 14, 17, 19-21, 23, 27, 28, 33, 42, 43, 49, 50, 60, 61, 62, 70, 78, 81	III <sup>+</sup>	II <sup>2m-2a</sup>	IV <sup>2m-2a</sup>	III <sup>2-m</sup>	II <sup>2m-2b</sup>		III <sup>2m-2a</sup>	I <sup>2b</sup>					1
<i>Sphagnum subsecundum</i>	2, 4-6, 10, 16-18, 25, 27-29, 34, 35, 38, 42, 50, 57, 57, 62, 64, 67-71, 73-75, 78, 79, 82, 85	III <sup>2m</sup>	V <sup>+-3</sup>		II <sup>2a</sup>	III <sup>2a-2b</sup>	V <sup>2b-3</sup>	III <sup>2a-2b</sup>	I <sup>2b</sup>					
<i>Carex echinata</i>	1, 4, 6, 8-11, 13, 16-18, 20, 21, 25, 28, 34-36, 38, 39, 42, 50, 57, 61, 62, 66, 68, 71, 74, 76, 77, 81, 82, 85	III <sup>+</sup>	II <sup>+-1</sup>	III <sup>+</sup>	V <sup>+-2a</sup>	IV <sup>+-3</sup>	V <sup>+-2b</sup>	II <sup>+-2a</sup>	I <sup>+</sup>		+		2a	
<i>Eriophorum angustifolium</i>	11, 18, 28, 35, 64, 71, 74	III <sup>+</sup>	II <sup>+-2a</sup>	III <sup>+-2m</sup>	II <sup>+</sup>	I <sup>3</sup>								
<i>Gentianella bulgarica</i>	1, 4, 8, 10, 13, 14, 17, 18, 21, 23, 25, 44, 49, 50, 69-71,	III <sup>+</sup>	II <sup>+</sup>	IV <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>	III <sup>+</sup>				+			

continued

DIMITROV M., NATCHEVA R., GANEVA A., GYUROVA D.  
PLANT BIODIVERSITY OF *SPHAGNUM*-DOMINATED MIRES IN VITOSHA NATURE PARK

continuation of Annex II

	74-79, 81, 82													
<i>Eriophorum vaginatum</i>	2, 3, 5, 6, 8, 10, 12, 15, 19, 22, 23, 25, 35, 37-39, 43, 44, 49, 50, 57, 61, 63, 70, 71, 74, 78, 79, 81, 82	III <sup>+</sup>	III <sup>+-2a</sup>	II <sup>+</sup>	V <sup>2b-4</sup>	III <sup>+-4</sup>	III <sup>+</sup>	I <sup>1</sup>	II <sup>+-1</sup>	II <sup>+</sup>				
<i>Juniperus sibirica</i>	1, 2, 4, 5, 7-10, 13, 14-16, 18-21, 23, 25, 27, 29, 37, 42, 44, 49, 57, 62-65, 69-71, 73-75, 79	V <sup>+</sup>	III <sup>+-2a</sup>	V <sup>+</sup>	IV <sup>+</sup>	IV <sup>+-3</sup>	III <sup>3</sup>	I <sup>2a</sup>	II <sup>+</sup>				+	
<i>Bruckenthalia spiculifolia</i>	4, 7, 9, 10, 12, 13, 16, 20, 21, 23, 25, 28, 29, 32, 37-39, 43, 44, 61, 65, 69, 70, 71, 74	III <sup>2m</sup>	IV <sup>+-2a</sup>	IV <sup>+-1</sup>	II <sup>+</sup>	III <sup>+-2a</sup>		I <sup>+</sup>	II <sup>+</sup>					
<i>Vaccinium myrtillus</i>	4, 7-10, 13, 16, 17, 21, 25, 27, 29, 36, 42, 44, 50, 57, 65, 69-71, 74, 79, 82	III <sup>+</sup>	II <sup>+</sup>	III <sup>+-2m</sup>		III <sup>+-2b</sup>	III <sup>+</sup>		I <sup>2a</sup>					
<i>Vaccinium vitis-idaea</i>	2, 7, 10, 13, 14-17, 20, 23, 25, 29, 32, 38, 39, 43, 44, 50, 57, 65, 69, 71, 74	III <sup>+</sup>	III <sup>+</sup>	IV <sup>+-2m</sup>	II <sup>+</sup>	II <sup>+</sup>			I <sup>2a</sup>					
<i>Sphagnum capillifolium</i>	1-3, 6, 7, 10-13, 15-17, 19-23, 25, 29, 32, 33, 43, 44, 49, 50, 57, 62-65, 70, 71, 73-76, 78, 79, 81, 82	V <sup>2m-2b</sup>	III <sup>2a-3</sup>	IV <sup>2b-3</sup>	V <sup>2b</sup>	V <sup>2m-3</sup>		II <sup>2b</sup>	II <sup>2a-2b</sup>				3	
<i>Homogyne alpina</i>	2-6, 9, 13, 16, 21, 27, 29, 44, 57, 70, 71, 73-76	III <sup>+</sup>		II <sup>+</sup>		III <sup>+</sup>			III <sup>+</sup>				+	
<i>Caltha palustris</i> agg.	4, 6, 13, 18, 21, 33, 37, 57, 60, 69, 77		I <sup>+</sup>	II <sup>+</sup>		I <sup>+</sup>	III <sup>2b</sup>	I <sup>+</sup>	II <sup>+-2a</sup>		+			+
<i>Rhizomnium punctatum</i>	12, 36, 73, 75, 85					I <sup>2m</sup>	III <sup>2a</sup>		II <sup>2m</sup>	II <sup>2a</sup>			2m	
<i>Brachythecium rivulare</i>	8, 11, 17, 37-39, 65, 85		II <sup>+-1</sup>			I <sup>2m</sup>	V <sup>2a</sup>							
<i>Angelica pancicii</i>	1, 10-13, 16-18, 20, 32, 35, 44, 61-63, 65, 69, 75, 85	III <sup>+</sup>	II <sup>+</sup>	IV <sup>+-1</sup>	II <sup>+</sup>	II <sup>+</sup>	III <sup>+</sup>	II <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>				
<i>Holcus molis</i>	13, 33, 60, 61, 62, 67, 68			II <sup>1</sup>				V <sup>+-2b</sup>						+
<i>Calliergonella cuspidata</i>	14, 16-18, 23, 25, 28, 33, 60-62, 64, 68, 69	III <sup>2a</sup>	II <sup>2m-2a</sup>	II <sup>2a</sup>	II <sup>2a</sup>	I <sup>2m-2a</sup>		IV <sup>2m-2b</sup>	I <sup>2a</sup>					2m
<i>Calamagrostis arundinacea</i>	14, 62			II <sup>+</sup>				I <sup>3</sup>						
<i>Rubus idaeus</i>	17, 18, 44, 62		I <sup>+</sup>	II <sup>+</sup>				I <sup>2a</sup>						
<i>Pinus sylvestris</i>	1, 13, 14, 16-18, 68	III <sup>+</sup>	I <sup>+</sup>	III <sup>+</sup>		I <sup>+</sup>		I <sup>2b</sup>						
<i>Alnus glutinosa</i>	68							I <sup>2a</sup>						
<i>Saxifraga *alpigena</i>	11, 16, 21, 28, 29, 57, 70, 73	III <sup>+</sup>				I <sup>+</sup>			IV <sup>+-2b</sup>				+	
<i>Philonotis seriata</i>	5-7, 12, 16, 17, 21, 28, 29, 42, 44, 49, 57, 67, 73, 75, 85	V <sup>+-2m</sup>	II <sup>2m-1</sup>	II <sup>2m</sup>		I <sup>2m-2b</sup>	III <sup>2m</sup>	I <sup>2a</sup>	IV <sup>2a</sup>				2a	
<i>Soldanella montana</i>	4, 7, 11, 13, 16, 28, 57, 64, 70, 75, 76	III <sup>+</sup>		II <sup>+</sup>		II <sup>+</sup>			III <sup>+-3</sup>					
<i>Scapania undulata</i>	1, 2, 11-13, 16, 17, 19-21, 25, 28, 44, 49, 57, 61, 62, 64, 69, 70, 75, 79	V <sup>2m</sup>	II <sup>1-2m</sup>	IV <sup>1-2m</sup>	II <sup>1</sup>	II <sup>2m-2a</sup>		II <sup>1</sup>	V <sup>2m-2a</sup>					
<i>Oncophorus virens</i>	6, 12, 57					I <sup>2m</sup>			III <sup>2m-2a</sup>					
<i>Festuca rubra</i> agg.	2-4, 6, 13, 18-21, 27-29, 32, 34-37, 44, 62, 69, 70, 73, 81		II <sup>+-2a</sup>	IV <sup>+-2m</sup>	II <sup>+</sup>	II <sup>+-1</sup>		I <sup>+</sup>	II <sup>1</sup>	V <sup>+-3</sup>			+	
<i>Juncus effusus</i>	13, 17, 29, 33-35, 66, 68		I <sup>+</sup>	II <sup>1</sup>		I <sup>+</sup>		II <sup>+-1</sup>		IV <sup>+-2a</sup>		1		
<i>Carex rostrata</i>	4										4			
<i>Carex curta</i>	29, 77					I <sup>+</sup>					2b			
<i>Scirpus sylvaticus</i>	33-35, 66							I <sup>2b</sup>		IV <sup>+-2a</sup>		4		
<i>Lysimachia vulgaris</i>	14, 17, 66		I <sup>+</sup>	II <sup>+</sup>								2a		

continued

continuation of Annex II

<i>Scutellaria alpina</i>	66											+		
<i>Sphagnum fimbriatum</i>	66											2b		
<i>Luzula sudetica</i>	2, 3, 6, 7, 11, 21, 32, 39, 42, 50, 57, 64, 65, 69-71, 73-78		II <sup>+</sup>			III <sup>+-2a</sup>			IV <sup>+-2a</sup>		+		2a	
<i>Fissidens adianthoides</i>	60, 74					I <sup>r</sup>								1
<i>Hamatocaulis vernicosus</i>	60													1
<b>Other vascular plants</b>														
<i>Achillea millefolium</i>	17, 18, 62		I <sup>+</sup>					I <sup>+</sup>						
<i>Agrostis capillaris</i>	1, 2, 4-9, 13, 14, 17-21, 23, 25, 27, 32-36, 44, 50, 57, 62, 64, 69, 70, 74, 75, 78, 79, 85		III <sup>+-3</sup>	V <sup>+</sup>	III <sup>+</sup>	IV <sup>+-1</sup>	V <sup>+</sup>	II <sup>+-2b</sup>	I <sup>+</sup>	V <sup>+-3</sup>				
<i>Alchemilla vulgaris</i> agg.	1, 5, 7, 11-13, 25, 27, 28, 33, 57, 65, 69-71, 82, 85		I <sup>+</sup>	II <sup>+</sup>		II <sup>+</sup>	III <sup>+</sup>	I <sup>+</sup>	III <sup>+</sup>					
<i>Allium sibiricum</i>	2, 3, 5-7, 10-14-17, 20, 22, 27, 28, 32, 37-39, 43, 44, 50, 57, 69, 70, 71, 74, 75, 78, 79, 82	III <sup>+</sup>	III <sup>+-2b</sup>	V <sup>+-2a</sup>	II <sup>+</sup>	IV <sup>+-3</sup>			IV <sup>+-2b</sup>					
<i>Athyrium filix-femina</i>	13, 16, 17, 66	III <sup>+</sup>	I <sup>+</sup>	II <sup>2m</sup>								+		
<i>Betula pendula</i>	17, 32		I <sup>+</sup>											
<i>Bistorta major</i>	2, 7, 13, 15-17, 19-21, 29, 43, 44, 49, 82	V <sup>+-2a</sup>	II <sup>+</sup>	IV <sup>+-2m</sup>	II <sup>+</sup>	I <sup>+-2a</sup>								
<i>Briza media</i>	67							I <sup>+</sup>						
<i>Campanula abietina</i>	32, 44		I <sup>+</sup>	II <sup>+</sup>										
<i>Cardamine rivularis</i>	1, 4, 11, 13, 17, 37, 57, 70		I <sup>+</sup>	II <sup>2m</sup>		I <sup>+</sup>			I <sup>1</sup>					
<i>Carex ovalis</i>	32		I <sup>+</sup>											
<i>Carex pallescens</i>	2, 5, 13, 50, 70			II <sup>1</sup>		I <sup>+</sup>								
<i>Cirsium appendiculatum</i>	1, 4, 21, 28		I <sup>+</sup>			I <sup>+</sup>			II <sup>+</sup>					
<i>Crepis paludosa</i>	1, 2, 8, 29					I <sup>+-2a</sup>	III <sup>+</sup>							
<i>Crocus veluchensis</i>	3-6, 18, 44, 81		I <sup>+</sup>	II <sup>+</sup>		I <sup>+</sup>								
<i>Cruciata glabra</i>	1, 13, 18		I <sup>+</sup>	II <sup>+</sup>										
<i>Deschampsia caespitosa</i>	1-15, 17, 18, 20, 21, 23, 25, 28, 32, 34-37, 42-44, 49, 50, 57, 60, 62, 64, 68, 70, 71, 73-79, 81, 82	III <sup>+</sup>	IV <sup>+-1</sup>	V <sup>+-2a</sup>	II <sup>+</sup>	V <sup>+-2b</sup>	III <sup>4</sup>	II <sup>+-3</sup>	IV <sup>+-1</sup>	V <sup>1-3</sup>	2b		+	+
<i>Deschampsia flexuosa</i>	8, 13, 15, 17, 21, 35, 37, 42, 44, 57, 69-71, 74		II <sup>+</sup>	III <sup>+-2m</sup>		I <sup>+-1</sup>	III <sup>+</sup>		I <sup>1</sup>	II <sup>1</sup>				
<i>Dianthus superbus</i>	1, 15, 60-64, 69, 79				II <sup>+</sup>	I <sup>+-1</sup>		II <sup>1-2a</sup>						+
<i>Doronicum austriacum</i>	4					I <sup>+</sup>								
<i>Drosera rotundifolia</i>	13, 16, 17, 23-26, 28, 32, 43, 49, 61-65, 67, 68	V <sup>+</sup>	II <sup>+</sup>	II <sup>2m</sup>	III <sup>+</sup>	I <sup>+</sup>		II <sup>+</sup>						
<i>Epilobium angustifolium</i>	14, 17, 44, 49, 71	III <sup>+</sup>	I <sup>+</sup>	III <sup>+</sup>										
<i>Epilobium palustre</i>	4, 11, 13, 18, 20, 22, 34, 36, 69, 74-76, 79		I <sup>+</sup>	III <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>				IV <sup>+</sup>				
<i>Equisetum hiemale</i>	61, 63, 66				II <sup>+</sup>			I <sup>+</sup>				+		
<i>Equisetum palustre</i>	13, 17, 23, 33-35, 44, 50, 68, 77		I <sup>+</sup>	III <sup>+-1</sup>	II <sup>+</sup>	I <sup>+</sup>		II <sup>+</sup>		IV <sup>+</sup>	+			
<i>Eriophorum latifolium</i>	1, 2, 4, 7, 10-14, 16, 17, 27, 29, 37-39, 44, 49, 57, 61,	V <sup>+-2a</sup>	II <sup>+-1</sup>	IV <sup>+-2a</sup>		IV <sup>+-3</sup>	III <sup>4</sup>	I <sup>3</sup>	IV <sup>+-2b</sup>				+	

continued



DIMITROV M., NATCHEVA R., GANEVA A., GYUROVA D.  
PLANT BIODIVERSITY OF *SPHAGNUM*-DOMINATED MIRES IN VITOSHA NATURE PARK

continuation of Annex II

	64, 65, 69-71, 73-76, 79, 82, 85													
<i>Euphrasia rostkoviana</i>	6, 11, 16, 32, 64, 69, 71, 82	III <sup>+</sup>	I <sup>+</sup>											
<i>Festuca valida</i>	8						III <sup>+</sup>							
<i>Filipendula ulmaria</i>	1, 17, 18, 33, 34		I <sup>+</sup>			I <sup>+</sup>		I <sup>2a</sup>		II <sup>+</sup>				
<i>Galium palustre</i>	1, 32-34		I <sup>+</sup>			I <sup>+</sup>		I <sup>1</sup>		II <sup>+</sup>				
<i>Gentiana pneumonanthe</i>	6, 10, 11, 16, 49, 57, 68, 75, 78, 82	V <sup>+</sup>				II <sup>+</sup>		I <sup>+</sup>	I <sup>+</sup>					
<i>Geum coccineum</i>	1-13, 16-18, 21, 23, 25, 28, 29, 32, 34-37, 42-44, 50, 57, 61, 64, 65, 69, 71, 73-77, 81, 82, 85	III <sup>+</sup>	IV <sup>+</sup>	III <sup>+1</sup>	II <sup>+</sup>	IV <sup>+2a</sup>	V <sup>+</sup>	I <sup>+</sup>	V <sup>+</sup>	V <sup>+</sup>	+		+	
<i>Geum montanum</i>	4, 7					I <sup>+</sup>								
<i>Hieracium cymosum</i>	21								I <sup>+</sup>					
<i>Hypericum maculatum</i>	6-8, 17, 18, 20, 21, 32, 62, 69, 73		I <sup>+</sup>	II <sup>+</sup>		I <sup>+</sup>	III <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>				+	
<i>Hypericum perforatum</i>	17, 18, 67		I <sup>+</sup>					I <sup>+</sup>						
<i>Juncus articulatus</i>	1, 7, 8, 11, 12, 14-16, 29, 33, 49, 50, 61, 63, 65	V <sup>+2m</sup>		II <sup>+</sup>	II <sup>+</sup>	II <sup>+1</sup>	III <sup>+</sup>	II <sup>+</sup>	I <sup>+</sup>					
<i>Juncus conglomeratus</i> agg.	1, 8, 11, 14, 16-18, 21, 25, 28, 35, 36, 61-64, 67, 69, 85	III <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>	I <sup>+</sup>	V <sup>+2b</sup>	III <sup>+</sup>	II <sup>2a</sup>	IV <sup>+</sup>				
<i>Juniperus communis</i>	68							I <sup>+</sup>						
<i>Lathyrus pratensis</i>	1, 8, 32-35, 61, 62		I <sup>+</sup>			I <sup>+</sup>	III <sup>+</sup>	III <sup>+</sup>		IV <sup>+</sup>				
<i>Leontodon autumnalis</i>	3-5, 10-12, 18, 25, 27, 32, 33, 61, 62, 64, 65, 67, 69, 71, 75, 76, 78, 82		I <sup>+</sup>			III <sup>+</sup>		IV <sup>+</sup>	I <sup>+</sup>					
<i>Luzula forsteri</i>	43		I <sup>+</sup>											
<i>Luzula luzuloides</i>	27					I <sup>+</sup>								
<i>Luzula sylvatica</i>	13, 18, 27, 29		I <sup>+</sup>	II <sup>1</sup>		I <sup>+</sup>								
<i>Lysimachia nummularia</i>	18, 35, 37		I <sup>+</sup>							II				
<i>Lythrum salicaria</i>	13, 17, 18, 60, 67		I <sup>+1</sup>	II <sup>2m</sup>				I <sup>+</sup>						+
<i>Matricaria caucasica</i>	2, 4, 5, 9, 67					I <sup>+</sup>		I <sup>+</sup>						
<i>Mentha arvensis</i>	67							I <sup>+</sup>						
<i>Meum mutellina</i>	2-7, 10-12, 17, 18, 33, 34, 62, 69, 74, 82		I <sup>+</sup>			III <sup>+2a</sup>		II <sup>+</sup>	I <sup>+</sup>	II <sup>1</sup>				
<i>Myosotis scorpioides</i> agg.	1, 11, 15, 21, 33, 34, 37, 57, 69		I <sup>+</sup>			I <sup>+</sup>		I <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>				
<i>Nardus stricta</i>	1-5, 6, 8, 13, 18, 21, 29, 44, 62, 64, 68, 69-71, 73-76, 78, 79, 85		I <sup>+1</sup>	III <sup>+1</sup>		IV <sup>+3</sup>	V <sup>+</sup>	II <sup>+</sup>					1	
<i>Parnassia palustris</i>	1, 7, 8, 10, 12, 13, 16, 17, 25, 32, 57, 60, 63, 64, 77-79, 81, 82, 85	III <sup>+</sup>	I <sup>+</sup>	II <sup>2m</sup>	II <sup>+</sup>	II <sup>+</sup>	V <sup>+</sup>		II <sup>+</sup>		+			+
<i>Persicaria maculata</i>	6, 10, 11					I <sup>+</sup>								
<i>Phleum alpinum</i>	2, 23, 25, 28, 44, 49	III <sup>+</sup>		II <sup>+</sup>	II <sup>+</sup>	I <sup>+</sup>			I <sup>+</sup>					
<i>Phragmites australis</i>	33							I <sup>+</sup>						
<i>Picea abies</i>	8, 13, 14, 16-19, 23, 32, 42, 44, 49	V <sup>+</sup>	II <sup>+</sup>	IV <sup>+</sup>	III <sup>+</sup>	I <sup>+</sup>	III <sup>+</sup>							

continued

continuation of Annex II

	65													
<i>Pinguicula balcanica</i>	3-5, 7-13, 16, 21, 29, 49, 57, 60, 62, 64, 69, 77, 78	V <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>		III <sup>+</sup>	III <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>		+			+
<i>Pinus peuce</i>	42		I <sup>+</sup>											
<i>Plantago gentianoides</i>	6, 35					I <sup>+</sup>				II <sup>+</sup>				
<i>Potentilla palustris</i>	15, 22, 43		I <sup>+</sup>		II <sup>2b</sup>	I <sup>+</sup>								
<i>Potentilla ternata</i>	3, 4					I <sup>+</sup>								
<i>Pseudorchis albida</i>	16, 39, 42, 44, 65	III <sup>+</sup>	I <sup>+</sup>	II <sup>+</sup>		I <sup>+</sup>								
<i>Pseudorchis frivaldii</i>	1, 4, 5, 8-11, 16, 23, 25, 29, 32, 37-39, 44, 49, 65, 70, 71, 79, 81	V <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>	III <sup>+</sup>	III <sup>+</sup>							
<i>Ranunculus sartorianus</i>	2, 4-6, 9, 32-34, 69		I <sup>+</sup>			II <sup>+</sup>		I <sup>+</sup>		II <sup>+</sup>				
<i>Rhynanthus angustifolius</i>	13, 17, 32, 49	III <sup>+</sup>	I <sup>+</sup>	I <sup>+</sup>										
<i>Rumex acetosa</i>	1, 4, 18, 73		I <sup>1</sup>			I <sup>+</sup>							+	
<i>Salix caprea</i>	17-19, 32, 36, 63, 65		I <sup>+</sup>		III <sup>+</sup>	I <sup>+</sup>				II <sup>+</sup>				
<i>Salix lapponum</i>	1, 6, 10, 12-17, 35-39, 42-44, 49, 57, 65, 67, 69, 71, 75, 76, 78, 81	V <sup>+-2m</sup>	II <sup>+-2a</sup>	IV <sup>+-1</sup>		III <sup>+</sup>		I <sup>2a</sup>	II <sup>+-2a</sup>	IV <sup>+</sup>				
<i>Salix waldsteiniana</i>	12-14, 16, 17, 32, 44, 49, 57, 71	V <sup>+-2m</sup>	I <sup>+-1</sup>	IV <sup>+</sup>		I <sup>+</sup>			II <sup>+</sup>					
<i>Senecio nemorensis</i>	3, 7, 10, 20, 23, 50, 70, 79			II <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>								
<i>Senecio pancicii</i>	3, 5, 10, 13, 16, 17, 25, 28, 44, 49, 69-71, 79	V <sup>+</sup>	I <sup>+</sup>	III <sup>+</sup>		II <sup>+</sup>			I <sup>+</sup>					
<i>Sesleria comosa</i>	2, 4, 6, 12, 22, 44, 50, 57, 70, 71, 75, 76, 78, 79, 81, 82			II <sup>+</sup>	II <sup>+</sup>	III <sup>+</sup>			II <sup>+</sup>					
<i>Stellaria graminea</i>	32		I <sup>+</sup>											
<i>Swertia perennis</i>	6, 10, 63, 82				II <sup>+</sup>	I <sup>+</sup>								
<i>Taraxacum appeninum</i>	3, 4, 70					I <sup>+</sup>								
<i>Thalictrum aquilegifolium</i>	8						III <sup>+</sup>							
<i>Thlaspi praecox</i>	6, 44			II <sup>+</sup>		I <sup>+</sup>								
<i>Trichophorum caespitosum</i>	4, 7-10, 12, 13, 16, 20, 21, 23, 25, 29, 38, 42, 44, 49, 50, 70, 71, 74-76, 78, 79, 81, 82	V <sup>2a-2b</sup>	II <sup>2a-4</sup>	IV <sup>2m-2b</sup>	II <sup>3</sup>	III <sup>+-4</sup>	III <sup>+</sup>		I <sup>+</sup>					
<i>Trifolium michelii</i>	10, 32, 69, 85		I <sup>+</sup>			I <sup>+</sup>	III <sup>+</sup>							
<i>Trifolium pratense</i>	12, 28, 32, 69, 82		I <sup>+</sup>			I <sup>+</sup>			II <sup>+</sup>					
<i>Trifolium spadiceum</i>	10, 12, 13, 18, 28, 32, 62-64, 77, 82		I <sup>+</sup>	II <sup>+</sup>	II <sup>+</sup>	I <sup>+-2a</sup>		I <sup>2a</sup>	II <sup>+</sup>		+			
<i>Trisetum flavescens</i>	35									II <sup>+</sup>				
<i>Trolius europaeus</i>	1					I <sup>2a</sup>								
<i>Utricularia minor</i>	13, 14, 25, 37			III <sup>+</sup>										
<i>Vaccinium uliginosum</i>	1, 3, 4, 6, 7, 8, 10-13, 15-17, 23, 28, 29, 32, 37, 39, 42, 44, 50, 57, 69-71, 74-76, 79, 82, 85	III <sup>+</sup>	II <sup>+-2a</sup>	III <sup>+-1</sup>	II <sup>+-2a</sup>	IV <sup>+-2b</sup>	V <sup>+</sup>		III <sup>+-2b</sup>					
<i>Veratrum lobelianum</i>	1, 8, 12, 13, 17, 19, 20, 21, 23, 25, 27, 32, 34, 35, 37,		II <sup>+</sup>	IV <sup>+</sup>	III <sup>r-+</sup>	II <sup>+</sup>	III <sup>+</sup>		II <sup>+</sup>	IV <sup>+</sup>			+	

continued

DIMITROV M., NATCHEVA R., GANEVA A., GYUROVA D.  
PLANT BIODIVERSITY OF *SPHAGNUM*-DOMINATED MIRES IN VITOSHA NATURE PARK

continuation of Annex II

	42-44, 50, 64, 69, 71, 73, 74, 79, 81, 82													
<i>Viola dacica</i>	4					I <sup>+</sup>								
<b>Other mosses</b>														
<i>Aneura pinguis</i>	13			II <sup>2m</sup>										
<i>Aulacomnium palustre</i>	2, 3, 5, 6, 9, 12, 13, 15-19, 22, 32, 36, 42, 44, 49, 60, 63, 67-69, 73, 77, 78, 85	V <sup>2a</sup>	II <sup>2m-2a</sup>	III <sup>2m-2a</sup>	III <sup>2m-2a</sup>	II <sup>2m-2a</sup>	III <sup>2m</sup>	II <sup>2m</sup>	II <sup>2m</sup>	II <sup>2a</sup>	2a		2m	1
<i>Barbilophozia atlantica</i>	12								I <sup>2m</sup>					
<i>Brachythecium mildeanum</i>	19					I <sup>2m</sup>								
<i>Brachythecium reflexum</i>	79					I <sup>1</sup>								
<i>Brachythecium</i> sp.	36									II <sup>2m</sup>				
<i>Bryum pallens</i>	3, 12, 61, 62					I <sup>1</sup>		II <sup>1</sup>	I <sup>1</sup>					
<i>Bryum pseudotriquetrum</i>	1, 6, 7, 11-13, 16, 17, 21, 25, 36, 49, 57, 60, 64, 65, 70, 79	V <sup>2m</sup>	II <sup>+-2m</sup>	II <sup>2m</sup>		II <sup>1-2m</sup>			IV <sup>2m</sup>	II <sup>1</sup>				1
<i>Bryum weigelii</i>	11					I <sup>+</sup>								
<i>Calliergon cordifolium</i>	33							I <sup>1</sup>						
<i>Calypogeia azurea</i>	28, 71, 77, 79					I <sup>2m</sup>			I <sup>2m</sup>		2m			
<i>Calypogeia muehleriana</i>	13, 71, 76			II <sup>2m</sup>		I <sup>2m</sup>								
<i>Calypogeia neesiana</i> *	36									II <sup>2m</sup>				
<i>Calypogeia sphagnicola</i>	28, 42, 71		I <sup>1</sup>			I <sup>1</sup>			I <sup>1</sup>					
<i>Campylium polygamum</i> *	73												2m	
<i>Cephalozia bicuspidata</i>	12, 13, 28, 71, 75, 79			II <sup>2m</sup>		I <sup>2m</sup>			II <sup>2m</sup>					
<i>Cephalozia connivens</i>	71					I <sup>2m</sup>								
<i>Cephalozia pleniceps</i>	6, 35, 70					I <sup>+-2m</sup>				II <sup>2m</sup>				
<i>Cephaloziella divaricata</i>	3					I <sup>r</sup>								
<i>Cephaloziella hampeana</i>	79					I <sup>2m</sup>								
<i>Chiloscyphus pallescens</i>	10, 11, 13, 16, 19, 20, 35, 49, 61, 62, 69	V <sup>2m</sup>		III <sup>1-2m</sup>	II <sup>1</sup>	I <sup>2m-2a</sup>		II <sup>2m</sup>		II <sup>2m</sup>				
<i>Chiloscyphus polyanthos</i>	65, 69, 73					I <sup>1-2m</sup>							2m	
<i>Climacium dendroides</i>	1, 6, 8, 15, 19, 20, 27, 33, 60, 64, 69, 73, 81, 82			II <sup>2m</sup>	II <sup>2m</sup>	II <sup>2m-2a</sup>	III <sup>2m</sup>	II <sup>2a</sup>					2m	2m
<i>Dichelyma falcatum</i>	3	I <sup>2m</sup>												
<i>Dicranella palustris</i>	13, 16, 25, 44, 64, 66, 70, 71, 75, 81	III <sup>2m</sup>	II <sup>2m</sup>	III <sup>2m-2a</sup>		I <sup>2m-2a</sup>						1		
<i>Dicranum bonjeanii</i>	2, 6, 7, 10, 21, 22, 43, 70, 75		I <sup>1</sup>		II <sup>2a</sup>	II <sup>1-2a</sup>								
<i>Dicranum scoparium</i>	13, 70, 71, 75			II <sup>2a</sup>		I <sup>2m-2a</sup>								
<i>Jungermannia leiantha</i>	75					I <sup>2m</sup>								
<i>Kiaeria starkei</i>	12, 25, 60, 70, 75					I <sup>2m</sup>			I <sup>1</sup>					+

continued

continuation of Annex II

<i>Lophocolea heterophylla</i>	36									II <sup>2m</sup>				
<i>Lophozia obtusa</i>	12								I <sup>2m</sup>					
<i>Lophozia ventricosa</i>	70, 71, 75, 79					I <sup>2m-2a</sup>								
<i>Marchantia polymorpha</i>	19					I <sup>2m</sup>								
<i>Meesia uliginosa</i>	12								I <sup>1</sup>					
<i>Palustriella decipiens</i>	6-8, 10, 11, 16, 17, 21, 57, 75	III <sup>2m</sup>	I <sup>2m</sup>			I <sup>2m-2a</sup>	III <sup>2a</sup>		II <sup>2m</sup>					
<i>Pellia epiphylla</i>	16, 17, 35, 44, 69	III <sup>2m</sup>	I <sup>2m</sup>	II <sup>2m</sup>		I <sup>2m</sup>				II <sup>2m</sup>				
<i>Pellia neesiana</i>	22, 25, 36		I <sup>2m</sup>		II <sup>2m</sup>					II <sup>2m</sup>				
<i>Philonotis fontana</i>	6, 8, 21, 60-62, 66		I <sup>1</sup>			I <sup>2m</sup>	III <sup>2m</sup>	II <sup>1</sup>				1		1
<i>Plagiochila porelloides</i>	36									II <sup>2m</sup>				
<i>Plagiomnium affine</i>	61, 62							II <sup>2m</sup>						
<i>Plagiomnium ellipticum</i>	33, 36-39, 60		I <sup>1</sup>					I <sup>2a</sup>		II <sup>2a</sup>				2m
<i>Plagiomnium undulatum</i>	36, 60									II <sup>2b</sup>				2m
<i>Plagiothecium denticulatum</i> var. <i>undulatum</i>	35									II <sup>r</sup>				
<i>Pohlia nutans</i>	61							I <sup>r</sup>						
<i>Polytrichum commune</i>	2, 5, 8, 9, 12-17, 19, 20, 22, 32, 35, 37-39, 43, 44, 49, 50, 60-62, 66, 68, 69, 71, 77, 79, 85	V <sup>2a</sup>	II <sup>2m-2a</sup>	V <sup>2m-2a</sup>	III <sup>2a</sup>	II <sup>2m-2b</sup>	V <sup>2a</sup>	III <sup>2m-2a</sup>	I <sup>2a</sup>	II	2a	2a		2m
<i>Polytrichum strictum</i>	6, 15, 16, 43, 44, 49, 62, 70, 74, 79, 81	V <sup>2m-2a</sup>	I <sup>2a</sup>	II <sup>2m</sup>		II <sup>2a-2b</sup>		I <sup>2a</sup>						
<i>Rhizomnium magnifolium</i>	8, 13			II <sup>2m</sup>			III <sup>2m</sup>							
<i>Rhizomnium pseudopunctatum</i>	19					I <sup>2m</sup>								
<i>Rhytidiadelphus squarrosus</i>	73												2m	
<i>Riccardia incurvata</i>	13, 57			II <sup>2m</sup>					II <sup>1</sup>					
<i>Riccardia multifida</i>	13, 60-62			II <sup>2m</sup>				II <sup>+</sup>						1
<i>Sanionia uncinata</i>	75					I <sup>2m</sup>								
<i>Schistidium rivulare</i>	21		I <sup>r</sup>											
<i>Scorpidium cossonii</i>	6, 73, 76, 82					I <sup>2a</sup>							2a	
<i>Scorpidium revolvens</i>	12, 13, 76			II <sup>2m</sup>		I <sup>2m</sup>			I <sup>2a</sup>					
<i>Sphagnum angustifolium</i>	13, 16, 19-21, 23, 42, 43	III <sup>2a</sup>	II <sup>1-2m</sup>	III <sup>+-2m</sup>	III <sup>+-2m</sup>									
<i>Sphagnum auriculatum</i>	3, 9, 10, 12-14, 16, 17, 22, 37-39, 49, 57, 71, 74, 76, 82	V <sup>2m-2a</sup>	II <sup>2m-2a</sup>	III <sup>2a-2b</sup>	II <sup>2a</sup>	II <sup>2a-2b</sup>			III <sup>2b</sup>					
<i>Sphagnum centrale</i>	17, 21, 25, 32, 36, 68, 71		I <sup>2m-2a</sup>			I <sup>2m</sup>		I <sup>2b</sup>		II <sup>2b</sup>				
<i>Sphagnum compactum</i>	50					I <sup>2m</sup>								
<i>Sphagnum contortum</i>	9, 10, 13, 14, 16, 17, 19, 20, 22, 23, 29, 33, 44, 49, 57, 61, 62, 73, 78, 81	V <sup>2a-2b</sup>	I <sup>2b</sup>	V <sup>2a-2b</sup>	IV <sup>2b</sup>	I <sup>2a-2b</sup>		III <sup>2a-3</sup>	III <sup>2a</sup>				2a	



DIMITROV M., NATCHEVA R., GANEVA A., GYUROVA D.  
PLANT BIODIVERSITY OF *SPHAGNUM*-DOMINATED MIRES IN VITOSHA NATURE PARK

continuation of Annex II

<i>Sphagnum fallax</i>	16, 21, 23, 32, 35, 42, 71	III <sup>2a</sup>	II <sup>2m-2a</sup>		II <sup>2a</sup>	I <sup>2a</sup>				II <sup>2m</sup>				
<i>Sphagnum flexuosum</i>	13, 14, 16, 17, 21, 23, 25, 27, 28, 32, 33, 35, 36-39, 42-44, 49, 60, 62-65, 68, 73, 81, 82	V <sup>2a-2b</sup>	III <sup>2a-3</sup>	IV <sup>2b-3</sup>	III <sup>2a-3</sup>	I <sup>2a-2b</sup>		III <sup>2b</sup>	II <sup>3</sup>	IV <sup>2b</sup>			2b	1
<i>Sphagnum fuscum</i>	22, 44			II <sup>2b</sup>	II <sup>2a</sup>									
<i>Sphagnum girgensohnii</i>	8, 11, 13, 16, 17, 32, 35-39, 42-44, 49, 62, 66, 68, 70, 71, 85	V <sup>2m-2a</sup>	III <sup>2m-2a</sup>	III <sup>2a-2b</sup>		I <sup>2a</sup>	V <sup>2a</sup>	II <sup>2m-3</sup>		IV <sup>2a</sup>		2a		
<i>Sphagnum inundatum</i>	1, 9, 60, 68, 69, 73, 77, 78					I <sup>2m-2a</sup>		I <sup>2m</sup>			2a		2a	2m
<i>Sphagnum magellanicum</i>	44, 49, 71	III <sup>3</sup>		II <sup>2b</sup>		I <sup>2a</sup>								
<i>Sphagnum palustre</i>	13, 19-21, 23, 25, 35, 36, 43, 44, 49, 50, 65, 71, 79	III <sup>2b</sup>	I <sup>2a-2b</sup>	IV <sup>2a-2b</sup>	III <sup>2a-2b</sup>	I <sup>2a-2b</sup>				IV <sup>2a</sup>				
<i>Sphagnum quinquefarium</i>	1	III <sup>1</sup>												
<i>Sphagnum russowii</i>	13-15, 17, 21, 25, 36, 43, 49, 71, 77, 85	III <sup>2a</sup>	I <sup>1-2a</sup>	III <sup>2m-2a</sup>		I <sup>2m-2a</sup>	III <sup>2a</sup>			II <sup>2a</sup>	2a			
<i>Sphagnum squarrosum</i>	8, 10, 19-21, 32, 34-39, 42, 49, 66, 82	III <sup>2m</sup>	III <sup>2m-2a</sup>	II <sup>2m</sup>	II <sup>2m</sup>	I <sup>2m-2a</sup>	III <sup>2m</sup>			V <sup>2a</sup>		2m		
<i>Sphagnum subnitens</i>	75					I <sup>2m</sup>								
<i>Sphagnum teres</i>	10, 13, 14, 16, 17, 19-22, 23, 25, 29, 32-39, 44, 62, 63, 70, 77, 82	III <sup>2a</sup>	II <sup>2a-2b</sup>	V <sup>2m-2b</sup>	V <sup>2a-2b</sup>	I <sup>2a-2b</sup>		II <sup>2a</sup>		V <sup>2b</sup>	2b			
<i>Straminergon stramineum</i>	1, 6, 13, 16, 17, 19, 20, 22, 28, 36-39, 49, 50, 69-71, 74, 82	V <sup>2m</sup>	II <sup>2m</sup>	III <sup>2m-2a</sup>	III <sup>2m-2a</sup>	II <sup>2m-2a</sup>			I <sup>2a</sup>	II <sup>2a</sup>				
<i>Tomentypnum nitens</i>	10, 44			II <sup>2m</sup>		I <sup>2a</sup>								
<i>Warnstorfia exannulata</i>	2, 4-6, 9, 10, 12-16, 21-23, 25, 28, 29, 32, 33, 37-39, 43, 44, 49, 50, 61-64, 67, 69-71, 74-78, 81, 82	V <sup>2m</sup>	III <sup>2m-2a</sup>	IV <sup>2m-2a</sup>	IV <sup>2m-2a</sup>	IV <sup>1-2a</sup>		IV <sup>2m-2b</sup>	II <sup>2m-2a</sup>		2a			
<i>Warnstorfia sarmentosa</i>	4, 6, 9, 10, 12-14, 16, 19, 20, 23, 25, 28, 49, 50, 57, 61, 62, 70, 71, 75, 78, 79, 81	V <sup>2m-2a</sup>	II <sup>2a</sup>	IV <sup>2m-2a</sup>	III <sup>2m-2a</sup>	III <sup>2a-2b</sup>		II <sup>1</sup>	IV <sup>2a</sup>					

## PLANT SPECIES DIVERSITY AND INVASIBILITY OF (PERI-)URBAN FORESTS OF LJUBLJANA, SLOVENIA

<sup>1</sup>KUTNAR L., <sup>2</sup>ELER K.

<sup>1</sup>Slovenian Forestry Institute, Department of Forest Ecology, Ljubljana, Slovenia

<sup>2</sup>Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia & Slovenian Forestry Institute, Ljubljana, Slovenia

Corresponding author e-mail address: lado.kutnar@gozdis.si

**ABSTRACT:** Vascular plant species were assessed in the (peri-)urban forests of the Ljubljana region. Among the forests investigated were alluvial and riparian forests with high levels of plant species diversity. However, due to the vicinity of urban areas, the human impact on these forests is intense, and they are also highly exposed to the invasion of alien species. Within three forest locations in urban and peri-urban forests in the Ljubljana region, the vascular plant species were recorded, and their cover and origin status were estimated. In total, 161 vascular plant species were recorded at three study sites. Among them, 13 alien species were recognized. The majority of them were declared to be invasive species in Slovenia, which adversely affect the forest habitats and outcompete native plant species, and consequently, decrease the provisioning of some important ecosystem services such as erosion protection, recreation usage, biodiversity hosting, etc. The most abundant invasive species in the studied sites were *Solidago gigantea* Ait. and *Fallopia japonica* (Houtt.) Ronse Decr. The (peri-)urban forests of Ljubljana are under the significant impact of invasive plant species. The dynamics of expansion of invasive species should be carefully monitored.

**Keywords:** plant species, diversity, invasive species, (peri-)urban forest, Ljubljana.

### 1 INTRODUCTION

In urban environments, the proportion of alien species is closely related to the disturbance level (Chytrý *et al.*, 2008; Kowarik, 2008). In many heavily disturbed habitats (brownfields, urban wastelands, railway corridors, and industrial sites), alien species may appear as dominant species. Late-successional habitats are expected to host smaller populations of aliens; however, woody vegetation on urban sites may also include a surprisingly high amount of aliens (Kowarik, 2008).

In Slovenia, there are roughly 50 invasive vascular plant species (Jogan *et al.*, 2012). Black locust (*Robinia pseudoacacia* L.) with 0.6% of the total forest growing stock is the most widespread alien (non-native or non-indigenous) tree species in Slovenian forests (Kutnar & Pisek, 2013). It is widely distributed in lowlands and hilly areas, and in some urban areas. Due to ongoing climate change, even higher share of black locust (Kutnar & Kobler, 2013) and some other invasive plant species can be expected in the future (Dakskobler *et al.*, 2016).

Habitats vary considerably in the level of invasion (number or proportion of alien plant species they contain), which depends on local habitat properties, propagule pressure, and climate (Chytrý *et al.*, 2008). Lowland forests in Slovenia, such as riverine and floodplain forests woods, are highly exposed to the invasion and sub-spontaneous spread of invasive alien species (Dakskobler *et al.*, 2013, 2016). Rivers may act as the dispersal corridors of invasive alien species (Säumel & Kowarik, 2010). These invasive species may adversely affect the forest habitats and outcompete native plant species. Many invasive taxa have transformed both the structures and functions of ecosystems (Rejmánek *et al.*, 2005).

Among the (peri-)urban forests of the Ljubljana region, there are also alluvial and riparian forest types with high level of plant species diversity. However, due to the vicinity of urban areas, the human impact on these forests is intense, and they are also exposed to the invasion of alien species. Therefore, the aim of this study is to address the plant species diversity and to determine the number of alien plant species in selected urban and peri-urban forests in the Ljubljana region. Within the

same EMoNFu Project as in the Ljubljana region, the vegetation and flora in urban forests of Lombardy, Italy were also studied (Digiovinazzo & Padoa-Schioppa, 2014).

### 2 STUDY AREA AND METHODS

#### 2.1 Study plots and vegetation assessment

In the Ljubljana region, plots for the plant species assessment were set in three different forest sites: i) Gameljne site-1 (G1), ii) Gameljne site-2 (G2), and iii) Rožnik site (R) (Fig. 1).



**Figure 1:** Position of studied locations in Ljubljana region (Google Earth); G1 - Gameljne site-1; G2 - Gameljne site-2; R - Rožnik site

The Gameljne site-1 was selected on the upper terrace of the Sava River. It is a relatively dry site (gravel bedrock) without direct river influences (Fig 2). The second location, the Gameljne site-2 was selected in the flood area on the lower terrace of the Sava River (Fig 3). Both Gameljne sites are in the peri-urban area. The third

location was the Rožnik site in an urban area, close to the city centre (Fig 4).



**Figure 2:** Forest stand on the Gameljne site-1 with *Pinus sylvestris* L. in the upper tree layer, and various broadleaves in the lower tree layer (Photo: L. Kutnar)




**Figure 3:** The Gameljne site-2 near Ljubljana in spring aspect; it is under direct impacts of the Sava River (Photo: L. Kutnar)



**Figure 4:** Forest stand at the Rožnik site is located in Ljubljana urban area (Photo: L. Kutnar)

The size of each study site (location) was 2,500 m<sup>2</sup> (50 meters × 50 meters). Within each site, two different plots were installed; the large plot (400 m<sup>2</sup>) was set in the centre; 10 small plots (4 m<sup>2</sup>) were randomly selected around the large plot (Fig. 5).

In April 2013, and between June and July 2013, the assessments of plants and vegetation were performed on large and small plots.



**Figure 5:** Sampling design; distribution of large plot (in the centre) and 10 randomly distributed small plots around the centre

The cover of all vertical vegetation-layers (moss, herb, shrub, and tree layer) was estimated. The vertical vegetation-layers were defined according to the ICP-Forests protocol (Canullo *et al.*, 2011). The percentage cover of each of the above-mentioned layers, as well as the cover of bare soil and of surface rocks, were visually estimated.

All vascular plant species were recorded separately in each vertical layer (herb, shrub, and tree layer) (Canullo *et al.*, 2011). On large plots, the plant species cover was estimated using the Barkman *et al.* (1964) method, and the modified Londo (1975) method for small plots. The sources of the plant species nomenclature were Flora Europaea (Tutin *et al.*, 1964-1980; 1993) and national flora (Martinčič *et al.*, 2007).

## 2.2. Data analysis

The field data (vegetation relevés) of two temporal repetitions (spring and summer 2013) were merged into one common dataset for each plot. For further vegetation analysis, the cover estimations, for example from 1 to 5 of the Barkman (1964) scale and from 0.1 to 10 of the Londo (1975) scale, have been transformed to cover estimations in percentage, for example from 0.01% to 87.5% of the Barkman scale, and from 0.5% to 97.5% of the Londo scale.

The following measures of diversity were investigated:

- 1) Species richness (N) refers to the number of all plant species within a given plot;
- 2) The Shannon diversity index (H') is a measure that describes the structural composition of communities:

$$H' = -\sum_{i=1}^R p_i \ln p_i$$

where  $p_i$  is a relative cover of species  $i$  in a record.

The plant species diversity, diversity index, and share of invasive alien species were analysed for all large and small plots. Based on the plant species composition in vertical vegetation layers, 30 small plots and main diversity gradients were plotted in non-metric multidimensional scaling (NMS ordination) (McCune & Mefford, 2011).



### 3 RESULTS AND DISCUSSION

#### 3.1 Plant species composition and diversity

In the secondary pine forest of the Gameljne site-1, *Pinus sylvestris* L. was the dominant species in the upper tree layer, and *Carpinus betulus* L., *Tilia cordata* Mill., and *Corylus avellana* L. were recorded in the lower tree layer. In the forest stand of the Gameljne site-2 near the Sava River, *Alnus glutinosa* (L.) Gaertn., *Salix eleagnos* Scop., *Acer pseudoplatanus* L., and *Fraxinus excelsior* L. were among the most abundant tree species. Plots of the Rožnik site were overgrown by *Quercus petraea* (Matt.) Liebl., *Picea abies* (L.) Karst., *Castanea sativa* Mill., and *Robinia pseudacacia* L.

The most abundant shrub species on the Gameljne site-1 were *Ligustrum vulgare* L., *Lonicera caprifolium* L., and *Viburnum lantana* L.; *Rubus caesius* L. and *Cornus sanguinea* L. on the Gameljne site-2; and *Rubus hirtus* W. & K. on the Rožnik site.

The most common herb-layer species on the Gameljne site-1 were *Carex alba* Scop., *Calamagrostis varia* (Schr.) Host, *Anemone trifolia* L., *Carex flacca* Schreb., *Galanthus nivalis* L., and *Melica nutans* L.. The dominant herb-layer species on the Gameljne site-2 were *Solidago gigantea* Ait., *Galanthus nivalis* L., *Allium ursinum* L., *Aegopodium podagraria* L., *Fallopia japonica* (Houtt.) Ronse Decr., *Lamium orvala* L., and *Ranunculus ficaria* L. The dominant herb-layer species on the Rožnik site were *Anemone nemorosa* L., *Erythronium dens-canis* L., *Pteridium aquilinum* (L.) Kuhn, *Crocus vernus* (L.) Hill subsp. *vernus*, and *Maianthemum bifolium* L.

On all large and small plots on three sites (Gameljne site-1 and Gameljne site-2, Rožnik site), 161 vascular plant species were recorded in total (Table I). The number of vascular species varied between 61 (Rožnik site) and 85 (Gameljne site-2) per site.

**Table I:** Species diversity at three sites in the (peri)-urban forest

	All sites	Gameljne site-1	Gameljne site-2	Rožnik site
All plots	n=33	n=11	n=11	n=11
N Vascular species	161	69	85	61
Tree species	36	21	14	22
Shrub species	21	14	15	7
Herb species	104	34	56	32
Invasive species	9	0	8	3
Large plot (400 m <sup>2</sup> )	n=3	n=1	n=1	n=1
N species per plot	63.3	66	68	56
Shannon H'	3.39	3.17	3.51	3.48
Small plot (4 m <sup>2</sup> )	N=30	N=10	N=10	N=10
N species per plot	19.3	17.7	22.1	18.1
Shannon H'	2.24	2.26	2.34	2.09

In total, in all vertical vegetation layers; herb, shrub, and tree layers, 36 tree species were identified. The number of all shrub species, including climbers, was 21. Altogether, 104 herb layer species (including only non-woody species) were documented.


On average, 63 plant species per large plot, and 19 plant species per small plot were documented.

#### 3.2 Alien species

On the Gameljne site-2 near the Sava River, and on the Rožnik urban site, several alien plant species were found (Table I, Fig. 6). The majority of them have been declared to be invasive species in Slovenia. These invasive plant species, which represent 6% of the flora recorded (Fig. 7), probably come from different urban areas, e.g. gardens and parks. The majority of invasive species identified on studied sites are of North American or Asian origin.



**Figure 6:** In summer time, the forest floor of the Gameljne site-2 was covered by a dense layer of invasive alien species, *Solidago gigantea*, *Fallopia japonica*, *Rudbeckia laciniata* L. and others (Photo: L. Kutnar)



**Figure 7:** Alien and invasive species in the floristic spectrum of three (peri)-urban forests in the Ljubljana region

Among three sites in (peri)-urban forests in the Ljubljana region in Slovenia, the level of plant invasion was the highest at the Gameljne site-2 (9% or 8 species) (Table I). The most abundant invasive species on this site were *Solidago gigantea* and *Fallopia japonica* (including hybrid *F. x bohemica*) (Fig. 8 and Fig. 9). Other invasive species were *Rudbeckia laciniata* (Fig. 10), *Erigeron annuus* (L.) Pers., *Helianthus tuberosus* L., *Impatiens glandulifera* Royle, *Impatiens parviflora* DC., and *Physocarpus opulifolius* (L.) Maxim.


Besides the invasive species *Robinia pseudacacia*, *Berberis thunbergii* DC., and *Impatiens parviflora*, three alien but non-invasive species were also recorded on the Rožnik site, namely *Aesculus hippocastanum* L., *Prunus laurocerasus* L., and *Juglans mandshurica* Maxim.



**Figure 8:** *Solidago gigantea*, and similar species *Solidago canadensis* L. are invasive alien species of riverine, floodplain and swamp forests in Slovenia (Photo: L. Kutnar)



**Figure 9:** *Fallopia japonica* and *F. x bohemica* are invasive species that colonize different riparian ecosystems, roadsides, and waste places. (Photo: L. Kutnar)



**Figure 10:** *Rudbeckia laciniata* is one of the most widespread species on the Gameljne site-2 and also in Slovenia (Photo: L. Kutnar)


### 3.2 Main ecological and diversity gradients

Distribution of plots in the NMS ordination space reflects the main ecological and diversity gradients (Fig. 11). Axis 2 correlates closely with the gradient of water availability. At the lower values of Axis 2, the Gameljne

site-2 plots are placed, which are under significant river impact, including occasional flooding. The forest vegetation of this site belongs to a *Salici-Populetum* s. lat. association.

At the upper part of the ordination space (Fig. 11), plots on the Gameljne site-1 and Rožnik site, where direct water impacts are not pronounced, may be found. The forest vegetation of these two sites may be considered a *Quercus-Carpinetum* s. lat. plant community. However, the forest community composition of both sites is rather different, mostly due to differences in soil and bedrock properties. Both sites were exposed to significant human impacts, such as the provision of timber and fuel wood, litter gathering for farm use, collecting of non-timber forest products, and urbanisation, industrialisation, recreation activities mostly in the last decades, and therefore these forest stands were modified or even converted to secondary forests.

Additionally, Figure 11 depicts the main diversity gradients. On average, plots at the bottom of the ordination space are characterized by higher plant species diversity (see also Table I). The abundance of invasive plant species was higher on the Gameljne site-2 than on the Rožnik site, and on the Gameljne site-1 where no invasive species were found. Values of the Shannon diversity index were slightly higher on the Gameljne site-2 (2.34) and the Gameljne site-1 (2.26) than on the Rožnik site (2.09) (Table I, Fig. 11).



**Figure 11:** NMS ordination of small plots. Three test sites (locations) are represented by symbols as follows: G1 – Gameljne site-1; G2 – Gameljne site-2; R – Rožnik site. Scaling of plots was based on the plant species composition and species abundance in vertical vegetation layers. The main diversity gradients (the number of all vascular species ( $N_{total}$ ), the number of invasive alien species ( $N_{invasive\ sp.}$ ), and Shannon diversity index ( $H'$ )) are plotted. The lengths of arrows depict the increase in species number or in the index value.

### 4 CONCLUSIONS

In studied sites of the Ljubljana (peri-)urban forest area, high levels of vascular plant species diversity were determined. If the diversity of these forests is compared with the other main forest types of Slovenia (e.g. ICP



Forest Level II plots; Kutnar, 2006) the species diversity of (peri-)urban forests is, on average, higher.

The total number of plant species (161 species) recorded on three sites in Ljubljana region is also significantly higher than in the sites within different lowland forests, urban plantations, and parks in the Lombardy region, Italy (120 species on 14 sites; Digiovinazzo & Padoa-Schioppa, 2014). In both regions, the same harmonised method for evaluation of plant species diversity was applied.

However, the significant impact of invasive alien plant species was determined in the (peri-)urban forests of Ljubljana. Among all the plant species identified on studied sites 8% were alien and 6% of all were invasive. In comparison to the studied Ljubljana forests, the portion of alien species in Lombardy urban forests and parks was higher (15%; Digiovinazzo & Padoa-Schioppa, 2014). Kowarik (2008) reported an even higher amount of alien species number (33%) in urban stands dominated by *Robinia pseudoacacia*, but it is known that disturbed human-made habitats with herbaceous vegetation are the most invaded by aliens (Chytrý *et al.*, 2008).

It was suggested that there is not necessarily a relationship between the invasibility of a plant community and the number of species present in that community (Crawley *et al.*, 1999; Davis *et al.*, 2000). Other studies show that such a relationship exists: positive at the landscape scale (Stohlgren *et al.*, 1999; Sax, 2002) and negative at scales of < 1 m<sup>2</sup> (Levine, 2000). Taking into account 30 small plots (4 m<sup>2</sup>) studied in (peri-)urban forests in the Ljubljana region, the weak positive correlation (Pearson  $r = 0.254$ ) between total species number and the number of invasive species was established.

Invasive species may outcompete native plant species, may change forest habitats, and may disturb natural forest regeneration, ecosystem services, and functions (Rejmánek *et al.*, 2005). Among forest services in urban areas, the recreation use is one of the most significant. Besides the considerable change of the plant species composition of forest vegetation in the Sava River, and partly also in the urban forest on the Rožnik site, we determined that the dense cover of invasive species may negatively affect the accessibility and use of paths along river and the forest alone, and invasive species may affect the whole physiognomy of the riparian forests.

Therefore, in Ljubljana (peri-)urban forests, the dynamics of expansion of invasive species should be carefully monitored, and appropriate measures for its limitation need to be established in the near future. More forest management and silvicultural activities should be engaged with regard to the invasive species that already disturb the forest development, structure, and functions of (peri-)urban forests.

## 5 ACKNOWLEDGEMENTS

This research was financially supported by the LIFE+ EMoNFUr Project (10/ENV/IT/000399, 'Establishing a monitoring network to assess lowland forest and urban plantation in Lombardy and urban forest in Slovenia'). The preparation of the paper was partly supported by the Research Programmes P4-0107 and P4-0085, financed by the Slovenian Research Agency.

We would like to thank Matej Rupel, Andrej Verlič, Urša Vilhar and other colleagues from the Slovenian

Forestry Institute for help in the field and other assistance with this study.

The manuscript has undergone revision for the English language by Terry T. Jackson.

## 6 REFERENCES

- [1] Barkman, J.J., Doing, H., Segal, S., Kritische Bemerkungen und Vorschläge zur quantitativen Vegetationsanalyse, *Acta bot. neerl.*, 13, (1964), 394–419.
- [2] Canullo, R., Starlinger, F., Granke, O., Fischer, R., Aamlid, D., Neville, P., Assessment of ground vegetation. Manual Part VII-SP1, In: ICP Forests. Manual on methods and criteria for harmonized sampling, assessment, monitoring and analysis of the effects of air pollution on forests, UNECE ICP Forests Programme Coordinating Centre, Hamburg, (2011), [http://www.icp-forests.org/pdf/FINAL\\_GV.pdf](http://www.icp-forests.org/pdf/FINAL_GV.pdf)
- [3] Chytrý, M., Jarošík, V., Pyšek, P., Hájek, O., Knollová, I., Tichý, L., Danihelka, J., Separating habitat invasibility by alien plants from the actual level of invasion. *Ecology*, 89, (2008): 1541–1553.
- [4] Crawley, M.J., Brown, S.L., Heard, M.S., Edwards, G.R., Invasion-resistance in experimental grassland communities: species richness or species identity? *Ecology Letters*, 2, 3, (1999), 140–148.
- [5] Dakskobler, I., Kutnar, L., Šilc, U., Poplavni, močvirni in obrežni gozdovi v Sloveniji = Floodplain woods, swamp woods and riverine forests in Slovenia, Ljubljana: Silva Slovenica, Gozdarski inštitut Slovenije: Zveza gozdarskih društev Slovenije - Gozdarska založba, (2013), pag. 127. (In Slovenian with English abstract).
- [6] Dakskobler, I., Kutnar, L., Šilc, U., Vreš, B., Prisotnost in pogostnost tujerodnih rastlinskih vrst v gozdnih rastiščnih tipih Slovenije = Presence and frequency of alien plant species in forest site types of Slovenia. In: Jurc, M. (Ed.). *Invazivne tujerodne vrste v gozdovih ter njihov vpliv na trajnostno rabo gozdnih virov: zbornik prispevkov posvetovanja z mednarodno udeležbo = Invasive alien species in forests and their impact on the sustainable use of forest resources: lectures presented at the conference with international participation*, Ljubljana: Biotehniška fakulteta, Oddelek za gozdarstvo in obnovljive gozdne vire (2016), 125–141.
- [7] Davis, M.A., Grime, J.P., Thompson, K., Fluctuating resources in plant communities: a general theory of invisibility, *Journal of Ecology*, 88, (2000), 528–534.
- [8] Digiovinazzo, P., Padoa-Schioppa, E., Vegetation and flora monitoring in urban forests of Lombardy, Presentation at LIFE+ EMoNFUr Final Conference, Milano, 17. 6. 2014. (2014).
- [9] ICP Forests, 2013. the International Co-operative Programme on Assessment and Monitoring of Air Pollution Effects on Forests operating under the UNECE Convention on Long-range Transboundary Air Pollution plots, <http://icp-forests.net/>
- [10] Jogan, N., Eler, K., Novak, Š., Priročnik za sistematično kartiranje invazivnih tujerodnih rastlinskih vrst, Zavod Symbiosis, Nova vas, (2012), pag. 53. (In Slovenian).

- [11] Kowarik, I., On the Role of Alien Species in Urban Flora and Vegetation, In: Marzluff, J.M., Shulenberg, E., Endlicher, W., Alberti, M., Bradley, G., Ryan, C., ZumBrunnen, C., Simon, U. (Eds.), Urban Ecology: An International Perspective on the Interaction between Humans and Nature, Springer, (2008), 321–338.
- [12] Kutnar, L., Intenzivni monitoring vegetacije gozdnih ekosistemov v Sloveniji: Intensive monitoring of vegetation of forest ecosystems in Slovenia, In: Hladnik, D. (Ed.). Monitoring gospodarjenja z gozdom in gozdnato krajino: Monitoring the management of forests and forested landscapes, Studia forestalia Slovenica, št. 127, Ljubljana, Biotechnical Faculty, Department of Forestry and Renewable Forest Resources, (2006), 277–290. (In Slovenian with English summary).
- [13] Kutnar, L., Kobler, A., Sedanje stanje razširjenosti robinije (*Robinia pseudoacacia* L.) v Sloveniji in napovedi za prihodnost: The current distribution of black locust (*Robinia pseudoacacia* L.) in Slovenia and predictions for the future, Acta Silvae et Ligni 102, (2013), 21–30. (In Slovenian with English summary).
- [14] Kutnar, L., Pisek, R., Non-native and invasive tree species in the Slovenian forests. Gozdarski vestnik, 71, 9, (2013), 402–417. (In Slovenian with English summary).
- [15] Levine, J.M., Species diversity and biological invasions: relating local process to community pattern, Science, 288, 5467, (2000), 852–854.
- [16] Londo, G., The decimale scale for relevées of permanent quadrats, In: Knapp, R. (ed.), Handbook of Vegetation Science, 4, (1975), 45–50.
- [17] Martinčič, A., Wraber, T., Jogan, N., Podobnik, A., Turk, B., Vreš, B., Mala flora Slovenije – Ključ za določanje praprotnic in semenk: Flora of Slovenia – Identification Key for Ferns and Vascular Plants. Tehniška založba Slovenije, Ljubljana, (2007), pag. 968. (In Slovenian).
- [18] McCune, B., Mefford, M.J., PC-ORD. Multivariate Analysis of Ecological Data. Version 6.19, MjM Software, Gleneden Beach, Oregon, USA, (2011).
- [19] Rejmánek, M., Richardson, D.M., Pyšek, P., Plant invasions and invasibility of plant communities, In: van der Maarel, E. (Ed.), Vegetation ecology. Blackwell Publishing, (2005), 332–355.
- [20] Sax, D.F., Native and naturalized plant diversity are positively correlated in scrub communities of California and Chile, Diversity and Distributions, 8, (2002), 193–210.
- [21] Säumel, I., Kowarik, I., Urban rivers as dispersal corridors for primarily wind-dispersed invasive tree species, Landscape and Urban Planning, 94, (2010), 244–249.
- [22] Stohlgren, T.J., Binkley, D., Chong, G.W., Kalkhan, M.A., Schell, L.D., Bull, K.A., Otsuki, Y., Newman, G., Bashkin, M., Son, Y., Exotic plant species invade hot spots of native plant diversity, Ecological Monographs 69, (1999), 25–46.
- [23] Tutin, T.G., Burges, N.A., Chater, A.O., Edmondson, J.R., Heywood, V.H., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A., Flora Europaea, Vol. 1 (*Psilotaceae* to *Platanaceae*). 2nd Edition, Cambridge University Press, Cambridge, MA, USA, (1993), pag. 581.
- [24] Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A., Flora Europaea, Vol. 2–5. Cambridge University Press, Cambridge, MA, USA, (1964–1980), pag. 455, 370, 523, 467.

## FIRST RECORDS FOR SEVEN SPECIES AND ONE HYBRID FOR THE FLORA OF THE REPUBLIC OF MACEDONIA

MELOVSKI Lj., HRISTOVSKI S.

*Institute of Biology, Faculty of Natural Sciences and Mathematics, Ss. Cyril and Methodius University, Skopje, Macedonia*

*Corresponding author e-mail address: melovski@pmf.ukim.mk*

**ABSTRACT:** Contribution to the knowledge of distribution ranges of several plant species is presented in this article. The following plant species were recorded in course of the botanical research during last several years and they represent new findings for the flora of the Republic of Macedonia: *Polypodium interjectum*, *Polypodium x mantoniae*, *Clematis recta*, *Salix repens* subsp. *rosmarinifolia*, *Lathraea rhodopea*, *Cirsium grecescui*, *Eleocharis mamillata* subsp. *austriaca* and *Arabidopsis arenosa*.

**Keywords:** Macedonia, flora, new records.

### 1 INTRODUCTION

The flora of the Republic of Macedonia is extraordinary rich in comparison with the territory of the country and taking into account that it is a landlocked country. According to recent data, 210 families, 920 genera and 3700 species comprise the flora of higher plants, angiosperms being the richest group with about 3200 species [28]. This makes the flora of the Republic of Macedonia one of the richest floras in the context of the whole European continent [43]. Diverse geological, geomorphological and climatic factors, as well as historic circumstances have contributed for such richness and diversity of vascular plants [28, 37]. As a result, the current Macedonian flora represents a mosaic of various floral elements – Tertiary relicts, Mediterranean, Greek-Asia Minor, Caucasian, arcto-alpine, middle-European, Eurasian, Holarctic and cosmopolitan [28]. Endemic taxons (Illirian, Scardo-pindic, Balcan etc.) represent a special characteristic and value of the Macedonian flora. Among them, 114 flowering plant species are Macedonian endemics [28, 40, 41]. Due to these floral values, 42 Important Plant Areas were recently identified across Macedonian territory [37].

The flora of the Republic of Macedonia has been intensively studied for more than one and a half century. The most extensive data can be found in [3, 4, 5, 6, 17, 30, 39]. Floristic contributions with new species and/or findings for the flora of Macedonia are regularly published as well, like for example in [10, 11, 12, 13, 26, 27, 29, 31, 32, 33, 34, 35, 42, 43]. In addition, several new plant species were described in the last ten years from the territory of the Republic of Macedonia: *Helianthemum marmoreum* Stevan., Matevski & Kit Tan [41], *Jurinea micevskii* Stevan., Matevski & Kit Tan [42], *Festuca jakupicensis* Kostadinovski [21], *Viola pseudaeolica* Tomović, Melovski & Niketić [44].

This paper represents the continuation of the floristic studies of vascular flora of the Republic of Macedonia. Data on seven new species and one new hybrid for the flora of Macedonia, collected at different localities throughout the country are presented in this paper.

### 2 METHODS

The study was based on fieldwork in the territory of the Republic of Macedonia during the last two decades and collected and stored plant material in Herbarium Melovskiorum (MKMEL!) [47], as well as in the Biology Students Research Society herbarium of the Faculty of

Natural Sciences and Mathematics in Skopje. Besides the field survey, relevant literature sources were used for supplementation of the distribution records.

The determination of species was done according to standard floras and related recent taxonomic articles. Identification of the two *Polypodium* taxa was performed with microscopic method by analysis of indurated cells per annulus and basal cells per sporangium as well as the proportion of sterile spores in sporangia [22].

We followed the nomenclature of the taxa used in Euro+Med Plant Base (<http://www2.bgbm.org/EuroPlusMed/query.asp>). The quotation of the distribution locations and habitats for all records in this article follows the MAKMEL data base pattern.

The regional division of Melovski *et al.* [38] was used to present the general geographic position of the localities.

### 3 RESULTS AND DISCUSSION

*Polypodium interjectum* Shivas in J. Linn. Soc., Bot. 58: 28 (1961)


#### *First chorological records in Macedonia*

Mt. Belasica: above v. Smolare – Smolarski Vodopad, on the rock in chestnut forest, 550 m a.s.l., 15.09.2009, leg. S. Hristovski, det. Lj. Melovski (MKMEL! 04335-04339); Bistra: Lopušnik – Dolen Lopušnik, on limestone rocks in narrow dale with horse chestnut and beech forest, 900-1000 m a.s.l., 10.08.2010, leg./det. Lj. Melovski (MKMEL! 04341); Bistra: above v. Janče – lower part of the valley of Galička Reka, silicate rocks in mixed broadleaf forest next to the river, 800-900 m a.s.l., 29.07.2010, leg./det. Lj. Melovski (MKMEL! 04343-04347); Plačkovica: Zrnovska Reka – between Orlov Kamen and Ploča peaks, mixed broadleaf forest with linden, beech, *Carpinus betulus* and other trees, 600-700 m a.s.l., 4.09.2014, leg./det. Lj. Melovski (MKMEL! 08189); Plačkovica: v. Laki – Selska Reka valley (between Strumički Rid and Kartal - below and above the inflow of Lomska Reka), eroded stony site next to the road in beech forest, 950-1000 m a.s.l., 24.10.2014, leg./det. Lj. Melovski (MKMEL! 08190); Drimkol: Drenočka Reka – just above the inflow into Crn Drim, limestone stony site in mixed broadleaf forest in mesophilous ravine, 735 m a.s.l., 16.11.2014, leg./det. Lj. Melovski (MKMEL! 08191 and 08192).

*P. interjectum* grows in deep and humid forested dales and ravines and was found in several localities in Macedonia (Fig. 1). It is broadly distributed in Europe

[51] but its distribution in Balkan Peninsula was not well studied until recently when it was recorded in several Balkan countries [7, 21, 22]. This species was not mentioned in the Flora of the Republic of Macedonia Micevski [39]. The ecology of this species and its closely

related *P. vulgare* as well as the main taxonomic characteristics is described in Ivanova [22].



**Figure 1:** Distribution map of the first records of seven plant species and one hybrid in the Republic of Macedonia

***Polypodium x mantoniae* (Rothm.) Shivas** in Rothm. & U. Schneid., Kulturpflanze Beih. 3: 245 (1962) [*P. interjectum* Shivas x *P. vulgare* L.]

#### First chorological records in Macedonia

Treska canyon: Matka – near the cave "Krštalna", limestone stony ground, 320 m a.s.l., 04.09.2009, leg. S. Hristovski, det. Lj. Melovski (MKMEL! 04329-04333); Dren: below Ligurasa – toward Dlaboki Dol, on silicate rocks in shady and mesophilous dale, 950 m a.s.l., 28.04.2012, leg./det. Lj. Melovski (MKMEL! 04348); Šar Planina: Podgora – Petočko Vodiče (above v. Belovište), Silicate rocks next to the brook in beech forest, 920 m a.s.l., 12.09.2012, leg./det. Lj. Melovski (MKMEL! 04349-04351); Plačkovica: above v. Gradec – Next to Gradečka Reka (below Garvan), on silicate rocks in beech forest - mesophilous site, 870 m a.s.l., 30.11.2012, leg./det. Lj. Melovski (MKMEL! 04352).

This taxon was described in 1960-ties but its distribution is not well known. Only recently some papers appeared confirming its distribution in Balkan Peninsula [7, 19, 21, 22]. *P. x mantoniae* can be found together with the populations of its parent plant species but it also has well established populations in sites where one parent

plant is absent (Fig. 1). This may be result of lack of complete distribution data for both parent plants. It is also evident that *P. x mantoniae* populations are vigorous and effectively spread vegetatively [22].

***Salix repens* subsp. *rosmarinifolia* (L.) Andersson** in Sp. Pl. 1020 (1753) [syn.: *Salix rosmarinifolia* L.]

#### First chorological records in Macedonia

Šar Planina: Dolno Lukovo Pole – upper part, peat-bog/mire, 1730 m a.s.l., 24.05.2010, leg. Lj. Melovski & S. Hristovski, det. Lj. Melovski (MKMEL! 04313-04315); Šar Planina: Dolno Lukovo Pole – upper part, peat-bog/mire, 1730 m a.s.l., 10.07.2011, leg./det. Lj. Melovski (MKMEL! 04311 and 04312).

Only one locality of this willow species in north-western Macedonia (in the southern part of Šar Planina Mt.) is known so far (Fig. 1). Only one cluster of *Salix repens* subsp. *rosmarinifolia* (Fig. 2) was discovered in Dolno Lukovo Pole peat land (Fig. 3). This cluster of shrubs was partially burnt at the end of 2011. In the south and central Balkans it is known from Bulgaria [18] and Serbia [24].





**Figure 2:** *Salix repens* subsp. *rosmarinifolia* – habitus; (a) the whole cluster, (b) branches with leaves, (c) young branch with female flowers (Photo: S. Hristovski)



**Figure 3:** Dolno Lukovo Pole mires – *Salix repens* subsp. *rosmarinifolia* habitat (Photo: Lj. Stefanov)

***Clematis recta* L.** in Sp. Pl. 554 (1753)

*First chorological records in Macedonia*

Rožden region (Mariovo): v. Mrežičko – below Glavite, mixed broadleaf forest in mesophilous ravine on limestone ground – next to the stream, 600 m a.s.l., 30.05.2010, leg. Lj. Melovski & S. Hristovski, det. Lj. Melovski (MKMEL! 02961).

*Clematis recta* was found only on one locality close to v. Mrežichko in Mariovo region (Fig. 1). This species is common in Central, South and East Europe [50].

***Arabidopsis arenosa* (L.) Lawalrée** in Bull. Soc. Rov. Bot. Belg. 42: 242 (1969). 1969 [syn.: *Cardaminopsis arenosa* (L.) Hayek]

*First chorological records in Macedonia*

Bilina Planina: v. Podržikonj, Zli Dol – below Mišinci, beech forest, 1000 m a.s.l., 15.05.2005, leg./det. Lj. Melovski (MKMEL! 05154); Osogovski Planini: Kriva Palanka – Varoški Rid, hornbeam forest with sessile oak, 700-750 m a.s.l., 15.05.2011, leg./det. Lj. Melovski (MKMEL! 05155); Osogovski Planini: Emirica – above the inflow of Emirička Reka into Zletovska Reka, eroded sandy site next to the road, 850 m a.s.l., 03.05.2008, leg./det. Lj. Melovski (MKMEL! 05156);

Osogovski Planini: Ruen – above Toranica mine, beech forest – close to the road, 1430 m a.s.l., 23.05.2009, leg./det. Lj. Melovski (MKMEL! 05157); Osogovski Planini: v. Stanci – in the village, grassy site next to the road and houses, 850 m a.s.l., 13.05.2007, leg. IDSB '07, det. Lj. Melovski (MKMEL! 05158); Osogovski Planini: v. Duračka Reka – next to Stanečka Reka, next to the road and houses, 700-750 m a.s.l., 13.05.2007, leg. IDSB '07, det. Lj. Melovski (MKMEL! 05159).

*Arabidopsis arenosa* is widespread species in Europe and in the neighbouring countries [2, 23, 27]. The species was cited by Hayek [20] for Ma (=Mazedonien) but it was not included in the flora of Macedonia by Micevski [39]. The term “Mazedonien” in Hayek [20] does not coincide with the present administrative territory of the Republic of Macedonia.

All of the presented records from Macedonia are from the north-east part of the country (Fig. 1).

***Lathraea rhodopea* Dingler** in Bot. Zeitung (Berlin) 35: 74 (1877)

*First chorological records in Macedonia*

Belasica: above v. Smolare – below Šarena Češma, mixed broadleaf forest with beech, 800-1100 m a.s.l., 17.05.2010, leg. Lj. Melovski & S. Hristovski, det. S. Hristovski & Lj. Melovski (MKMEL! 04316).

*Lathraea rhodopea* (Fig. 4) was found on Belasica Mt. (Fig. 1), which actually represents a continuation of the Rhodope massif and Slavyanka Mt. toward Macedonia. Up to now, this species was known from south Bulgaria and north-east Greece [1, 14].12]. The new finding extends its distribution range toward west.



**Figure 4:** *Lathraea rhodopea* from Belasica Mt. (Photo: S. Hristovski)

***Cirsium grecescui* Rouy** in Bull. Soc. Bot. France 37: 164 (1890)

*First chorological records in Macedonia*

Bistra-Šar Planina: Children's resort "Bunec" – below the resort, marshy site next to a small stream, 1250 m a.s.l., 25.08.2004, leg. Lj. Melovski & N. Angelova, det. Lj. Melovski (MKMEL! 00180).

*Cirsium grecescui* (Fig. 5) was found on the border between Shara Mt. and Bistra next to the wetland at the Bunec resort (Fig. 1). This record was already announced in Melovski & Matevski [36]. It was known from south and east Romania and north-east Serbia [8] although in the Flora of Serbia it is quoted as common plant, but without exact distribution localities [15]. However, it was recently reported from Sokolovica [49] and Rogozna mountains [44] in central and southwest Serbia which is the closest finding to the locality from Macedonia.



**Figure 5:** *Cirsium grecescui* from Bunec wetland – habitus (a) and inflorescence (b) (Photo: Lj. Melovski)

***Eleocharis mamillata* subsp. *austriaca* (Hayek) Strandh.** in Opera Bot. 9(2): 9 (1965) [Basionym: *Eleocharis austriaca* Hayek]

*First chorological records in Macedonia*

Šar Planina: Borislovec-Mala Smreka – Bogovinsko Ezero, sandy-gravelly place next to the lake, 1935 m a.s.l.,



08.09.2006, leg./det. Lj. Melovski (MKMEL! 01588); Šar Planina: Rudoka – above Crno Ezero, small pond, 2250 m a.s.l., 30.07.2008, leg. Lj. Melovski & N. Angelova, det. Lj. Melovski (MKMEL! 01590).

*Eleocharis mamillata* subsp. *austriaca* grows on Shar Planina next to the alpine lake Bogovinsko Ezero and above Crno Ezero (Fig. 1). This subspecies is known from Europe and Asia. Most of the European records come from the Alps but it is also known from Pyrenees, Carpathians and Pirin mountain in Bulgaria [16] and Tara Mt. in Serbia [52].

#### 4 CONCLUSIONS

Seven plant species and one hybrid were recorded for the first time in the flora of the Republic of Macedonia: *Polypodium interjectum* Shivas, *Clematis recta* L., *Salix repens* subsp. *rosmarinifolia* (L.) Andersson, *Lathraea rhodopea* Dingler, *Cirsium grecescui* Rouy, *Eleocharis mamillata* (H. Lindb.) H. Lindb. subsp. *austriaca* (Hayek) Strandh., *Arabidopsis arenosa* (L.) Lawalrée and *Polypodium x mantoniae*. The discovery of these species was well expected, or not surprising for some of them, since they are known from neighbouring countries. The records of *Salix repens* subsp. *rosmarinifolia*, *Cirsium grecescui* and *Eleocharis mamillata* subsp. *austriaca* represent one of the southernmost localities in Europe. These three species were found in wetlands on the high mountains in west Macedonia. The Balkan endemic species *Lathraea rhodopea* was found in south-east Macedonia and this record extends its distribution range westwards.

#### 5 ACKNOWLEDGEMENTS

We are grateful to Ms. Daniela Jovanovska for preparation of the distribution map and to Macedonian Ecological Society for supporting part of the field research in the scope of its regular activities.

#### 6 REFERENCES

- [1] Aneva, I. Y., Dimitrov, D. ST. and V. M. Vutov (2015). Flora and vegetation of Slavyanka Mountain. *Bulg. J. Agric. Sci.*, 21: 926–934.
- [2] Asenov, I. (1970). Rod Cardaminopsis (C. A. Mayer) Hayek. p. 447–448. In: Jordanov, D. (ed.). Flora Republicae Popularis Bulgaricae. Vol. IV. Bulgarian Academy of Sciences, Sofia, 748 pp.
- [3] Bornmüller, J. (1925). Beiträge zur Flora Mazedoniens I. *Bot. Jahrb.* **59**: 293–504.
- [4] Bornmüller, J. (1926). Beiträge zur Flora Mazedoniens II. *Bot. Jahrb.* **40(136)**: 1–125.
- [5] Bornmüller, J. (1928). Beiträge zur Flora Mazedoniens III. *Bot. Jahrb.* **41(140)**: 1–196.
- [6] Bornmüller, J. (1937). Zur Flora Mazedoniens. *Feddes Repert. Spec. Nov. Regni Veg.* **42(1–10)**: 126–142.
- [7] Bureš, P., Tichý, L., Wang, Y.-F. & Bartoš, J. (2003). Occurrence of *Polypodium x mantoniae* and new localities for *P. interjectum* in the Czech Republic confirmed using flow cytometry. *Preslia* **75**: 293–310.
- [8] Carucci, F. (2011). Filogenesi molecolare del genere *Cirsium* Mill. sect. *Eriolepis* (Cass.) Dumort. sect. *Eriolepis* (Cass.) Dumort. Tesi di Dottorato di ricerca in Biologia Avanzata (Sistematica Molecolare). Università degli studi di Napoli Federico II. Facoltà di Scienze Matematiche, Fisiche e Naturali, 149 pp.
- [9] Diklić, N. (1972). Rod Cardaminopsis. In: Josifović, M. (ed.). *Flora de la Republique Socialiste de Serbie. III. Academie Serbe des sciences et des arts. Classe des sciences naturelles et mathematiques*, Belgrade, p. 265–270.
- [10] Dimitrov, D. S. (2007a). Reports 41–58. In: Vladimirov V, Dane F, Matevski V & Tan K. (eds.), New floristic records in the Balkans 4. *Phytologia Balcanica* **13**: 113–114.
- [11] Dimitrov, D. S. (2007b). Reports 20–27. In: Vladimirov V, Dane F, Matevski V & Tan K. (eds.), New floristic records in the Balkans 5. *Phytologia Balcanica* **13**: 266.
- [12] Dimitrov, D. S. (2011). Reports 46–48. In: Vladimirov V, Dane F, Matevski V, Stevanović V & Tan K. (eds.), New floristic records in the Balkans 15. *Phytologia Balcanica* **17**: 135–136.
- [13] Dimitrov, D. S. (2012). Reports 55–77. In: Vladimirov V, Dane F, Matevski V & Tan K. (eds.), New floristic records in the Balkans 4. *Phytologia Balcanica* **18**: 76–78.
- [14] Eleftheriadou, E., Athanasiadis, N., Theodoropoulos, K. & Tsiropidis, I. (1998). Geographical distribution of *Arabis procurrens* Waldst. & Kit., *Lilium rhodopaeum* Delip. and *Lathraea rhodopaea* Dingler in Greece. pp. 57–61. In: Tsekos, I. & Moustakas, M. (Ed.) *Progress in Botanical Research: Proceedings of the 1st Balkan Botanical Congress*. Springer Science+Business Media, B.V.
- [15] Gajić, M. (1975). Fam. Asteraceae. In: Josifović, M. (ed.). *Flora de la Republique Socialiste de Serbie. VII. Academie Serbe des sciences et des arts. Classe des sciences naturelles et mathematiques*, Belgrade, p. 1–465.
- [16] Gregor, T. (2003). *Eleocharis mamillata* - distribution and infraspecific differentiation. *Folia Geobotanica* **38**: 49–64.
- [17] Grisebach, A. (1843). *Spicilegium Florae rumelicae et bithynicae exhibens synopsis plantarum quas in aest. 1839 legit auctor A. Grisebach* 1. Fridericus Vieweg et filius, Brunsvigae [Braunschweig].
- [18] Hájek, M., Hájková, P., Apostolova, I., Sopotlieva, D., Velez, N. (2006). Report 49–52. In: Vladimirov, V., Dane, F., Nikolić, T., Stevanović, V., Tan, K. (comp.). New floristic records in the Balkans: 2. *Phytologia balcanica* **12 (2)**: 279–301.
- [19] Helánová, K., Bureš, P., Šmarda, P., Horová, L. (2004). *Polypodium x mantoniae* (*P. interjectum* x *P. vulgare*) new hybrid in Romania, confirmed using flow cytometry. *Contribuții Botanice*, **39**: 7–11.
- [20] Hayek, A. (1927). *Prodromus Florae peninsulae Balcanicae*, 1. Band. Dahlembei Berlin, 1927.
- [21] Hršak, V. (2000). Two fern taxa from the *Polypodium vulgare* L. complex - new for Croatia. *Acta botanica croatica* **59 (2)**: 375–381.
- [22] Ivanova, D. (2006). *Polypodium interjectum* and *P. x mantoniae* (Polypodiaceae: Pteridophyta), new to the Bulgarian flora. *Phytologia Balcanica* **12 (2)**: 191–202.
- [23] Jones, B. M. G. (Edition 1), revised by Akeroyd, J. R. (Edition 2) (1993). *Cardaminopsis* (C.A. Meyer)

- Hayek. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Valentine, D. H., Walters, S. M., Weeb, D. A. (Eds.) *Flora Europaea, Volume 1 (second edition): Lycopodiaceae to Platanaceae*. Cambridge University Press, Cambridge, pp.351-352.
- [24] Jovanović, B., Tucović, A. (1972). Fam. SALICACEAE Lindl. In: Josifović, M. (ed.). *Flora de la Republique Socialiste de Serbie. III*. Academie Serbe des sciences et des arts. Classe des sciences naturelles et mathematiques, Belgrade, p. 405-457.
- [25] Kostadinovski, M. (2007). *Festuca jakupicensis* spec. nova (Poaceae) in the flora of the Republic of Macedonia. In: Filipovski, Gj., Lozanovski, R., Matevski, V. (eds.). *Collection of papers devoted to Academician Kiril Micevski. On the occasion of the 80 years of his birth*. Macedonian Academy of Sciences and Arts, Skopje, pp. 209-222.
- [26] Matevski, V. (2002). New data regarding the flora in Republic of Macedonia. In: Proceedings of 7th Symposium on Flora of Southeastern Serbia and Neighboring Regions, Dimitrovgrad, 9–13.
- [27] Matevski, V. (2002-2003). New data regarding the Flora in the Republic of Macedonia. *God. zb. Biol.* **55–56**: 1–10, Skopje.
- [28] Matevski, V. (2013). Diversity and origin of the flora of the Republic of Macedonia. Induction speech, contributions and bibliography of the new member of the Macedonian Academy of Sciences and Arts. Macedonian Academy of Sciences and Arts, Skopje, pp. 125-186 [in Macedonian].
- [29] Matevski, V. & Teofilovski, A. (2004). New contributions for the flora of the Republic of Macedonia. In: *Proceedings of the 2<sup>nd</sup> Congress of Ecologists of the Republic of Macedonia, 25-29.10.2004, Ohrid. Special issues of Macedonian Ecological Society* **6**, pp. 384–389.
- [30] Matevski, V. (ed.) (2010). *The Flora of the Republic of Macedonia II(1)*. Macedonian Academy of Sciences and Arts, Skopje, pp. 153-394.
- [31] Matevski, V. & Kostadinovski, M. (2009). New locality of species *Limosella aquatica* L. in the Flora of the Republic of Macedonia. *Hladnikia* **23**: 77-80.
- [32] Matevski, V. & Teofilovski, A. (2011). New species in the Flora of the Republic of Macedonia. *Biol. Macedonica* **62**: 49-54.
- [33] Matevski, V., Čarni, A. & Kostadinovski, M. (2005). *New data regarding the flora in Republic of Macedonia*. In: Proceedings of the 8th Symposium on the Flora of Southeastern Serbia and Neighbouring Regions, pp. 25-28.
- [34] Matevski, V., Čarni, A., Kostadinovski, M., Košir, P., Šilc, U., Zelnik, I. (2008). *Flora and vegetation of the Macedonian steppe*. Biol. Inst. ZRC, SAZU, Ljubljana.
- [35] Matevski, V., Čarni, A., Avramoski, O., Juvan, N., Kostadinovski, M., Košir, P., Marinšek, A., Paušič, A., Šilc, U. (2011). Forest vegetation of the Galičica mountain range in Macedonia. Založba ZRC, ZRC SAZU, Ljubljana.
- [36] Melovski, Lj., Matevski, V. (2008). Development of criteria for the selection of wetland habitats for strict protection - case study of Bunec site in the National Park "Mavrovo". Proceedings of the III Congress of Ecologists of Macedonia with International Participation. Special issues of Macedonian Ecological Society, Vol. 8, Skopje, p. 157-166.
- [37] Melovski, Lj., Matevski, V., Kostadinovski, M., Karadelev, M., Angelova, N. & Radford, E. A. (2010). Important Plant Areas in the Republic of Macedonia. Special issue of the Macedonian Ecological Society **9**, Skopje, 128 pp [in Macedonian].
- [38] Melovski, L., Markovski, B., Hristovski, S., Jovanovska, D., Anastasovski, V., Klincharov, S., Veleviski, M., Velkovski, N., Trendafilov, A., Matevski, V. and Kostadinovski, M. (2013). Regional division of the Republic of Macedonia for the needs of biological databases. *Macedonian Journal of Ecology and Environment*, 15(2), pp.81-111.
- [39] Micevski, K. (1985-2005). *The Flora of the Republic of Macedonia. Vol. I, Issues 1-6*: 1-1716. Macedonian Academy of Sciences and Arts, Skopje
- [40] Micevski, K. & Matevski, V. (1987). Die territoriale Verteilung der Endemiten in der SR Makedonien und das Problem ihrer Gefährdung. Symposium on: Protection of endemics in the living world of Yugoslavia (Sarajevo, May 15-16, 1986). Academy of Sciences and Arts of Bosnia and Herzegovina, Department of natural sciences and mathematics. *Special editions* **14**: 199–207.
- [41] MoEPP (2003). Country Study for Biodiversity of the Republic of Macedonia (First National Report). Ministry of Environment and Physical Planning, Skopje, 217 pp.
- [42] Niketić, M., Melovski, Lj. & Tomović, G. (2007). Reports 59–68. In: Vladimirov V, Dane F, Matevski V & Tan K. (eds.), New floristic records in the Balkans 4. *Phytologia Balcanica* **13(1)**: 114–116.
- [43] Niketić, M., Tomović, G., Melovski, Lj., Stevanović, V. and Matevski, V. (2014). New species for the vascular flora of Republic of Macedonia and their distribution in the Balkan Peninsula. *Botanica Serbica*, **38(1)**: 57-67.
- [44] Papović, O., Miljković, M., Randelović, N., Randelović, V. (2014). Analysis of the flora of Rogozna Mountain in Southwestern Serbia. *Biologica Nyssana*, 5 (1): 17-30.
- [45] Stevanović, V., Matevski, V., Tan, K. (2009). *Helianthemum marmoreum* (Cistaceae), a new species from the Central Balkan. *Botanica Serbica* **33(1)**: 13-19.
- [46] Stevanović, V., Matevski, V., Tan, K. (2010). *Jurinea micevskii* (Asteraceae), a new species from the Republic of Macedonia. *Phytologia Balcanica* **16(2)**: 249-254.
- [47] Thiers, B. (2015). *Index Herbariorum: A global directory of public herbaria and associated staff*. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/ih/>[continuously updated]
- [48] Tomović, G., Niketić, M., Lazarević, M. & Melovski, Lj. (2016). Taxonomic reassessment of *Viola aetolica* and *Viola elegantula* (V. sect. *Melanium*, Violaceae), with descriptions of two new species from the Balkan Peninsula. *Phytotaxa* **253 (4)**: 237–265.
- [49] Tomović, G., Niketić, M., M., Randelović, V. & Stevanović, V. (2005). The vascular flora of

- Mountain Sokolovica in Central Serbia (Serbia and Montenegro). *Fl. Medit.* **15**: 9-55.
- [50] Tutin, T. G. (Edition 1), revised by Akeroyd, J. R. (Edition 2) (1993). *Clematis* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Valentine, D. H., Walters, S. M., Weeb, D. A. (Eds.) *Flora Europaea, Volume 1 (second edition): Lycopodiaceae to Platanaceae*. Cambridge University Press, Cambridge, pp. 266-267.
- [51] Valentine, D. H. (Edition 1), revised by Akeroyd, J. R. and Jermt, A. C. (Edition 2) (1993). *Polypodium* L. In: Tutin, T. G., Heywood, V. H., Burges, N. A., Valentine, D. H., Walters, S. M., Weeb, D. A. (Eds.) *Flora Europaea, Volume 1 (second edition): Lycopodiaceae to Platanaceae*. Cambridge University Press, Cambridge, pp.15-16.
- [52] Zlatković, B., Tomović, G., Randelović, V., Vukojičić, S. & Niketić, M. (2009). Distribution and conservation status of several new and neglected vascular plants in Serbia. *Phytologia balcanica* **15** (1): 95 – 105, Sofia, 2009.

**BOREO-MONTANE FOREST PHYTOCOENOSES IN CENTRAL STARA PLANINA MTS.**<sup>1</sup>NIKOLOV I., <sup>2</sup>DIMITROV M.<sup>1</sup>Central Balkan National Park Directorate, Gabrovo, Bulgaria<sup>2</sup>University of Forestry, Department of Dendrology, Sofia, Bulgaria

Corresponding author e-mail address: mariusdimitrov@abv.bg

**ABSTRACT:** This study deals with the subalpine conifers and related birch forests on the territory of National Park "Central Balkan" in Bulgaria. It is a part of a comprehensive study on the forest vegetation in the Park. 30 relevés were collected during the field research. JUICE, TWINSpan and STATISTICA were applied for classification and statistical analysis. The studied forests are fragmented and do not form a continuous altitudinal belt within the park. The species composition, ecology and distribution determine their differentiation into four groups of phytocoenoses. Most widespread are the communities dominated by Norway spruce, which combine mesohygrophitic and eutrophic phytocoenoses, forming the timberline. Community of *Picea abies* and *Luzula sylvatica* includes woods, tied to acidophilous and relatively less humid and poor soil. Mesophytic and mesotrophic community of *Betula pendula* develops in areas where the timberline is composed of spruce forests, mostly southward from the main ridge of the mountain. The natural habitat of the Macedonian pine in the study area represents northernmost locality of its natural area of distribution and is the only one in Stara Planina Mts. Two associations were described: *Adenostylo alliariae-Pinetum peucis* and *Ranunculo carthusianae-Piceetum abietis*. They include hygromesophytic and mesotrophic forests, forming some parts of the alpine timberline.

**Keywords:** subalpine forests, classification of vegetation, diagnostic species, environmental variables, *Pinus peuce*, phytocoenoses.

**1 INTRODUCTION**

Holarctic disjunctions are biogeographical phenomena formed after cooling of the climate during the Pliocene and are expressed in both distribution of plant species and communities in areas with different latitude and different altitude [1]. Boreal mountain disjunction includes coniferous forests growing north in the taiga and their vicariant phytocoenoses that develop in the high mountains in the central and southern parts of the continent. This is determined by the temperature gradient that changes analogously from south to north in the plains and from lower to higher altitudes in the mountains. According to the proposed altitudinal stratification of mountain systems depending on environmental, floristic and vegetation criteria, the coniferous forests grow in the lower subalpine belt [2]. According to the traditions concerning the zone differentiation of vegetation in Bulgaria [3, 4, 5] the boreal coniferous forests grow in the high mountain belt, which is equal to the lower subalpine belt, according to the traditional European stratification [2, 6].

Boreal forests include monodominant and polydominant coniferous and deciduous-coniferous forest phytocoenoses developing on diverse habitats. In southeastern Europe this vegetation is defined as a submediterranean and subcontinental mountain vegetation [7]. Besides typical boreal elements it is constituted of part European and endemic elements, which makes it somewhat different from the typical boreal vegetation. This research confines to the approach of boreal vegetation [8] and the visions for the hemiboreal vegetation in Southeast Europe [9, 10, 11, 12]. Similar mountain vegetation in Europe is found only on higher mountain ridges - the Pyrenees, the Alps, Jura, Black Forest, Hercynian hills, Thuringia, Carpathians, Dinaric Alps, Rhodope, Rila, Pirin, Stara Planina, and the Caucasus. Regional features are the main reason for the

formation of a large number of regional plant associations.

Central Stara Planina is the northernmost mountain in Bulgaria with occurrence of vegetation types, common to the mountains in Central Europe [2] (between the boreal and mediterranean zone).

Forest communities in Central Stara Planina had been partly studied by different methodological approaches for collection, analysis and aggregation of phytocoenotic information [13, 14, 15, 16, 17, 18, 19, 20, 21, 22]. Currently there is not a comprehensive classification scheme providing an idea of the syntaxonomic diversity. The beech forests were relatively well studied out of the forest habitats and plant communities [23]. There is also some information about syntaxa in the forests in some of the reserves located within Central Stara Planina, differentiated using the Broun-Blanquet's approach [17, 18, 22].

This research is part of a comprehensive study of the forest vegetation within the "Central Balkan" National Park, which covers the highest part of the Central Stara Planina. Main purpose is to study the boreo-montane forest phytocoenoses, which includes subalpine coniferous and related birch forests.

**2 MATERIALS AND METHODS**

Central Stara Planina is the highest part of the Stara Planina Ridge, located between Zlatishki and Shipchenski passes (Fig. 1). Well shaped are Zlatishko-Tetevenski parts (Vejen peak – 2198 m.a.s.l., Troyanski (Kupena peak – 2169 m.a.s.l.) and Kaloferski (Botev peak – 2376 m.a.s.l.). The studied area falls within the Meridional zone, Submediterranean region, Balkan province [24]. The climatic conditions in this area are diverse and with dynamic changes in their values. This is due to the rough terrain, deep river valleys and hollows. The climate in the northern parts of the territory is temperate continental and

in the southern parts is subcontinental. The thermal conditions are characterized by well-defined seasons, depending on the topography of the region [25]. This study covers the areas in the mountain belt (1500 - 2000 m.a.s.l.), characterized by a short vegetation period, an average annual temperature between 3,3°C and 5,1°C and annual rainfall sum from 860 to 1300 mm/m<sup>2</sup> [25]. Hydrographic system has a very high overall density – 3,03 km/km<sup>2</sup>.



**Figure 1:** Location of Central Stara Planina Mts.

Soil variability is represented by zonal soils of the classes Cambisols – Dystric-Eutric Cambisols, Umbric Cambisols, Modic Cambisols [26].

Coniferous forests occupy a limited area primarily in the higher western and central parts of Central Stara Planina.

For purposes of this study in the period May 2011 – July 2013 30 relevés were established in the range of 1450-1850 m.a.s.l. The combined (cover-abundance) scale of Braun-Blanquet was used [27, 28].

When drawing up the environmental characteristics and while performing the gradient analysis the following factors and indicators were taken into account: exposition, slope, altitude, soil type, bedrock, stoniness, mechanical structure, morphological complexity, continentality, humidity, precipitation, light, and temperature. Environmental groups of plants were determined according to Ellenberg [29].

The nomenclature of higher plants is according to Delipavlov and Cheshmedjiev [30]. Floral elements were determined using the system of Walter [31].

Phytocoenological nomenclature is according to ICPN [32] and the referenced literature. Syntaxa to alliance level are in accordance with the concepts of Rodwell et al. [33] and Mucina et al. [34]. Natural habitats are according to Kavrakova et al. [35] and Biserkov et al. [36].

Phytocoenological analysis was conducted according to the Braun-Blanquet approach [27]. The classification analysis was performed by using the software product JUICE [37]. Cluster analysis was done using Modified TWINSpan [38] and additional use of control clustering tools TWINSpan 1979 and K-means [39]. Sorensen's dissimilarity index was used.

Phi-coefficient (fidelity – association ratio between species and plant units) was used to determine the diagnostic species [40]. Preliminary thresholds for

fidelity are  $\Phi \geq 20$  Lower,  $\geq 60$  Higher [41, 42, 43]. The values of the constancy of species were taken into consideration [41]. The thresholds accepted for constant species are those with 'Frequency threshold' –  $\geq 50$  Lower,  $\geq 60$  Higher. Dominant species have cover higher than the upper limit provided in the methodology  $\geq 20$  Lower,  $\geq 60$  Higher. For the purpose of comprehensive treatment of the database the program STATISTICA [44] was used.

Classification of species was done using the scale of Ellenberg [29]. To assess the influence of several factors (variables) we used simultaneously the multidimensional variance analysis. Also, descriptive analysis of the environmental variables was performed (Descriptive statistics). 26 environmental variables were compared in the range of 30 relevés. A separate grouping of environmental factors equal to the grouping of the phytocoenotic descriptions was achieved as a result. Principal Components Analysis (PCA) for ordinates of vegetation to environmental variables was put into practice [29].

### 3 RESULTS AND DISCUSSION

As a result of the implemented classification and coordination procedures 4 basic syntaxa were identified.

They are included in the following syntaxonomic scheme:

Class *Vaccinio-Piceetea* Br.-Bl. in Br.-Bl. et al. 1939

Order *Piceetalia excelsae* Pawłowski et al. 1928

Alliance *Piceion excelsae* Pawłowski et al. 1928

*Eu-Vaccinio-Piceion* Oberdorfer 1957


Association *Ranunculo carthusianae-Piceetum abietis* ass. nova (holotypus hoc loco, Annex I, rel. 8)

Community of *Picea abies* and *Luzula sylvatica*

Community of *Betula pendula*

Alliance *Pinion peucis* Horvat 1950

Association *Adenostylo alliariae-Pinetum peucis* ass. nova (holotypus hoc loco, Annex I, rel. 28)



**Figure 2:** PCA of the boreal mountain forests on the territory of Central Stara Planina. 1 – Assoc. *Ranunculo carthusianae-Piceetum abietis*; 2 – Comm. of *Picea abies* and *Luzula sylvatica*; 3 – Comm. of *Betula pendula*; 4 – Assoc. *Adenostylo alliariae-Pinetum peucis*.

Fig. 2 illustrates the spatial arrangement of the isolated phytocoenoses in relation to the floristic similarity and the main environmental factors. Phytocoenoses from group № 1 develop on moderately

moist acidic soils at low temperatures and sufficient light. Group № 2 prefers less acidic substrates and relatively higher temperatures. Plant communities of Group № 3 develop at relatively low humidity and on poorer soils. Group № 4 combines hydrophilic and eutrophic communities.

### 3.1 Forests of *Picea abies*

Total 18 phytocoenotic relevés were done in this type of forest. Group №1 includes monodominant spruce forests. Group №2 unites forests with domination or high participation of *Picea abies*, in some places dominated by *Pinus peuce* and rarely (one time each) by *Abies alba* and *Fagus sylvatica*. The phytocoenoses dominated by or with participation of *Betula pendula* - group № 3 – possess high floristic and ecological similarity with forests of *Picea abies*. Species composition and location of these forests show that they are part of the succession line, developing on the site where coniferous forests were destroyed in the past. That is why they are referred to the alliance *Piceion excelsae* and are considered simultaneously as related to climax spruce forests.

High-mountain forests of *Picea abies*, monodominant or involving other conifers, form associations related to suballiance *Eu-Vaccinio-Piceenion* Oberd. 1957 and alliance *Piceion excelsae* Pawłowski et al. 1928. In Central Europe these communities often include *Larix europea*, *Pinus cembra* and *Pinus uncinata*, which sometimes form also monodominant phytocoenoses. The spruce forests in Southeastern Europe are distinctive and with regional characteristics derived from both their history and the zonal location of mountains where they develop. Some authors accept the differentiation of regional suballiances, while others describe the regional associations. Based on some floristic characteristics in Romania the suballiance *Soldanello majori-Piceenion* Coldea 1991 is described [45, 46] as synvicariant of *Eu-Vaccinio-Piceenion* Oberd. 1957. The following associations are related to it: *Hieracio rotundati-Piceetum* Pawl. et Br.-Bl. 1939, *Luzulo sylvaticae-Piceetum* Wraber 1953, *Bruckenthalio-Piceetum* Borhidi 1969, *Vaccinio-Piceetum* Brez. et Had. 1962, *Sphagno-Piceetum* Brez. et Had. 1962, *Soldanello majori-Piceetum* Coldea et Wagner 1998 [45, 46, 47, 48, 49, 50, 51].

Phytosociologists in Serbia [52, 53, 54, 55, 56, 57, 58] point out a series of associations related to *Eu-Vaccinio-Piceenion* – *Junipero sibiricae-Piceetum* (Rudski 1947) Mišić et Popović 1980, *Oxali acetosellae-Piceetum abietis* (Rudski 1947) Mišić et Popović 1980, *Arctostaphyllo-Piceetum* (Jov. 1953) Mišić et Popović 1954.

The association *Piceetum subalpinum scardicum* Em (1962) 1986 was described in Macedonia (Jovanović et al. 1986). In Slovenia Zupančič (1980, 1999) includes spruce phytocoenoses in *Luzulo sylvaticae-Piceetum* Wraber 1963 em. Zupančič 1999 and *Luzulo albidiae-Piceetum* Zupančič 1980. For Western Balkans the following associations were reported: Bosnia and Herzegovina - *Sorbo-Piceetum* Fukarek 1964; Croatia – *Piceetum croaticum subalpinum* Ht. (1950) 1967, *Helleboro nigrae-Piceetum* (Ht. 1958) Trinajstić et Pelcer

2005; Montenegro - *Piceetum abietis bertisceum subalpinum* Blečić (1961) 1964 [54]. In Greece the spruce forests have limited distribution. Dafis [59] refers them to the association *Piceetum abietis* s.l. and alliance *Piceion excelsae*.

In Bulgaria the spruce phytocoenoses are not enough studied. Most frequently mentioned in the phytocoenotic is the association *Piceetum myrtillosum*, distinguished by using the dominant method of Russian-Scandinavian phytocoenological school [6, 60]. Michalik [18] mentions the association *Piceetum excelsae balcanicum* (nom. illeg. – ICPN Art. 2b, 5, 34a) for the territory of the reserve "Boatin" (Central Stara Planina) [32]. An attempt to clarify the syntaxonomical status of spruce phytocoenoses in Rila was done by Rusakova & Valchev [61]. They relate the studied communities to alliance *Piceion excelsae* without any associations. Later, some of their relevés were included in the diagnosis of association *Moehringio pendulae-Piceetum* Roussakova et Dimitrov 2005, described for the territory of Rila and Western Rhodopes [62]. Pine species like *Pinus peuce* (as ecological vicariant of *Pinus cembra*) and *Pinus sylvestris*, and sometimes (in Rila and Pirin) *Pinus mugo* as undergrowth element, are common in the alpine spruce forests in Bulgaria [63]. Spruce forests in the Central Stara Planina occupy separate areas – Dobrila, Vezhen, Bashitite, Cartula, Boatin, Cherni vrh, Rozinska ravna and others. Most of these forests are older than 200 years. *Picea abies* forms mixed communities with *Pinus peuce* near Vezhen massif and partly with *Pinus sylvestris* north of Vezhen peak.

#### 3.1.1 Ass. *Ranunculo carthusianae-Piceetum abietis* ass. nova hoc loco (Group № 1)

**Diagnostic species:** *Picea abies*, *Ranunculus carthusianus*, *Ribes alpinum*, *Rumex alpinus*, *Juniperus sibirica*, *Potentilla haynaldiana*. **Constant species:** *Picea abies*, *Juniperus sibirica*, *Ajuga reptans*, *Dryopteris filix-mas*, *Fagus sylvatica* ssp. *sylvatica*, *Myosotis sylvatica*, *Rubus idaeus*, *Scilla bifolia*, *Senecio nemorensis*, *Sorbus aucuparia*, *Vaccinium myrtillus*, *Vaccinium vitis-idaea*. **Dominant species:** *Picea abies*, *Juniperus sibirica*, *Vaccinium myrtillus*. **Nomenclature type (holotypus):** Relève 8, Annex I. The specific composition of species and environmental conditions are the reason for diagnosing the new association *Ranunculo carthusianae-Piceetum abietis*, which unites 11 phytocoenotic relevés.

The communities of the association *Ranunculo carthusianae-Piceetum abietis* are distributed in the range of 1600 – 1800 m.a.s.l. on slopes an average 8-10 degrees. (Fig. 3). The expositions vary, but those with western and northern components dominate, and there are also some with southern component (below Karatepe peak). The rocks are silicate – south Bulgarian granite, slate, and diorite. The soils are Umbric, rarely Dystric or Modic Cambisols – moderately rich, medium deep, sandy and clay soils. The average temperature is about 3 °C.

The tree layer covers about 60% in average. Dominant species is *Picea abies* (Annex I). The cover of shrub layer ranges from 0 to 60%. *Picea abies*, *Juniperus sibirica*, *Ribes alpinum*, *Sorbus aucuparia*, *Fagus sylvatica* ssp. *sylvatica*, *Rubus idaeus*, *Rumex alpinus* are



the most common species. The herbaceous layer has a cover of an average 40%. *Picea abies*, *Juniperus sibirica*, *Ranunculus carthusianus*, *Rumex alpinus*, *Vaccinium vitis-idaea*, *Veronica alpina* and others are with high constancy. Typical physiognomic feature for these communities is the low species diversity. These are light, subalpine forests, in certain areas changing into sparse early stage of forests involving *Juniperus sibirica*.



**Figure 3:** Ass. *Ranunculo carthusianae-Piceetum abietis*

Given the location of the forests at higher parts of the mountain and the climatic conditions, the boreal elements dominate (50%) in the phytogeographic spectrum, followed by the european (17%), arctic-alpine (13%), balkan, eurasian, euro-mediterranean, pontic and cosmopolitans (4% each).

Such types of forest are common in the Bashitite area, Karatepe, Dobrila, Gerdeka, Kumanitsa, and Harmana.

The combination of diagnostic species *Ranunculus carthusiana*, *Ribes alpinum*, *Rumex alpinus*, *Juniperus sibirica*, *Potentilla haynaldiana* reflects the highland character of the communities. These species are differential compared to other known associations of ordinary spruce on the Balkan Peninsula. The communities of the association *Ranunculo carthusianae-Piceetum abietis* are isolated from the similar forests, located in the western part of Stara Planina where the association *Junipero sibiricae-Piceetum* was established (Rudski 1947) Mišić et Popović 1980. The latter is distinguished by its open nature and the constant participation of species such as *Poa violacea*, *Potentilla ternata*, *Bruckenthalia spiculifolia*, *Nardus stricta*, *Polygonum bistorta*, *Hypericum alpigenum*, *Agrostis capillaris*, *Thymus glabrescens*, *Alchemilla pubescens*, *Polygala comosa*, etc. [52, 58]. Furthermore *Ranunculus carthusianus*, *Ribes alpinum* and *Potentilla haynaldiana* are missing.

Association *Ranunculo carthusianae-Piceetum abietis* differentiates from the spruce forests described in Rila and Rhodope Mountains in Bulgaria, where participate species with high constancy as *Moehringia pendula*, *Melampyrum sylvaticum*, *Campanula sparsa*, *Lerchenfeldia flexuosa*, *Dicranum scoparium*, *Calamagrostis arundinacea*, *Orthilia secunda*, *Valeriana tripteris* [62]. This research confirms the findings of

Roussakova and Dimitrov [62] that the spruce forests in Bulgaria differ from those located in neighboring countries which, although close as physiognomy and species composition, are characterized by the presence of species related to the specific historical development of the flora and vegetation. For example, the spruce communities in the Carpathian Mts are differentiated by a number of carpathian elements such as *Hieracium rotundatum*, *Syphythum cordatum*, *Dentaria glandulosa*, *Euphorbia carniolica*, *Soldanella major*, *Leucanthemum waldsteinii*, *Aconitum moldavicum*, etc. [48, 64]. The presence of species unusual for Central Stara Planina is typical in western Balkans - *Astrantia carniolica*, *Homogyne sylvestris*, *Adenostyles glabra*, *Lonicera alpigena*, *Sorbus mougeotii*, and *Helleborus niger* [6, 58].

### 3.1.2 Community of *Picea abies* and *Luzula sylvatica* (group №2)

**Diagnostic species:** *Picea abies*, *Pinus peuce*, *Dryopteris expansa*, *Laserpitium krapfii*, *Galium rotundifolium*, *Cardamine pectinata*, *Fragaria vesca*, *Geranium pyrenaicum*, *Galium odoratum*. **Constant species:** *Picea abies*, *Dryopteris filix-mas*, *Luzula luzuloides*, *Luzula sylvatica*, *Senecio nemorensis*, *Vaccinium myrtillus*. **Dominant species:** *Picea abies*, *Pinus peuce*.

The communities of this group are mainly located in the area between Kositsa peak and Yumruka peak (Fig. 4). They are distributed in the range of 1400 – 1600 m.a.s.l. The slopes are about 20 degrees. The exposures vary, but are dominated by those with northern and eastern components. The rocks are south Bulgarian granites, crystalline schists and others. Almost everywhere the soils are Umbric Cambisols, but in the lower parts are Dystric-Eutric Cambisols, medium deep, sandy and stony. The average temperature is above 4 °C. The environmental analysis shows that these are alpine forests, growing in places rich with nitrogen and water, shady and dark.



**Figure 4:** Community of *Picea abies* and *Luzula sylvatica*

These forests accomplish the transition from the mixed coniferous-deciduous forests to macedonian pine forests. The cover of the tree layer is 60-90%. Dominant and subdominant species is *Picea abies*, in certain communities *Pinus peuce* dominates, while in others

there is participation of *Fagus sylvatica* and *Abies alba*. In the shrub layer the coverage ranges from 0 to 20%, involving mainly undergrowth of *Picea abies* and *Fagus sylvatica*. The herbaceous layer has a cover an average of 30%. The most commonly found species are *Dryopteris filix-mas*, *Luzula luzuloides*, *Luzula sylvatica*, *Senecio nemorensis*, *Vaccinium myrtillus* (Annex I).

Boreal elements dominate in those communities too (40%) although their quantity is less compared to the monodominant spruce and macedonian pine forests that grow at high altitudes and in extreme ecological conditions. There is a relatively high participation of the sub-mediterranean elements (17%), followed by european (13%), mediterranean (9%) balkan (9%). Euro-mediterranean, euro-asian and cosmopolitan elements are involved with 4% each.

Typical physiognomic feature of these forests is the big cover of the tree layer, the lack of shrub layer at most places and poor grass layer with low coverage. The grass cover is formed only near small streams, bright open "windows" when the horizontal structure is disrupted, around springs. These phytocoenoses do not border the timberline zone, as the belt over them are common to the forests of *Pinus peuce*. The facies of *Luzula sylvatica* and *Senecio nemorensis* which develop in the lighter areas are typical.

### 3.1.3 Community of *Betula pendula* (group № 3)

**Diagnostic species:** *Betula pendula*. **Constant species:** *Fagus sylvatica* ssp. *sylvatica*, *Betula pendula*, *Luzula luzuloides*, *Vaccinium myrtillus*.

**Dominant species:** *Betula pendula*, *Juniperus sibirica*, *Vaccinium myrtillus*.

The communities of this group develop around the timberline in the areas where this line is composed of spruce forests, mostly south of the main ridge of the mountain. They are distributed in the range of 1400 – 1600 m.a.s.l. Sloppes are about 30-35 degrees. The expositions are diverse, but dominated by those with a southern component. The rocks are silicate – south Bulgarian granite, slate, diorite and some other less represented types. The soils are Umbric Cambisols, and in the lower parts Dystric Cambisols, moderately rich, medium deep, sandy, mostly stony. The average annual temperature is above 4 °C.



**Figure 5:** Community of *Betula pendula*

The cover of the tree layer is 50% on average. *Betula pendula* is the main dominant, but *Picea abies* dominates in some localities (Fig. 5). In the shrub layer *Fagus sylvatica* ssp. *sylvatica* and *Betula pendula* have sporadic occurrence. The grass layer has cover an average of 30%. Most commonly are found *Betula pendula*, *Luzula luzuloides* and *Vaccinium myrtillus* (Annex I).

The boreal floristic elements (36%) have the lowest relative participation in these forests. The higher temperatures and the open character of some of the communities are the major cause of strong presence of euro-mediterranean (25%) and mediterranean elements (7%), followed by euro-asian (14%), balkan (11%) and european (7%).

Characteristics of these forests are their fragmented distribution, their great lighting, the presence of *Betula pendula* in all layers, the acidophilous and oligotrophic character, and the low soil moisture. During the last 10–20 years there has been an expanding of their area and occupation of territories in the subalpine treeless zone in many places in the mountains, mainly in the high southern slopes – Ravnets massif over Vasil Levski, Karlovo and Sushitca villages, above Sopot, Anevo, Karnare, Iganovo and in several other places in the western part of the mountain.

### 3.2 Forests of *Pinus peuce*

12 relevés dominated by *Pinus peuce* which form a separate cluster of high floristic similarity were made during this research (group №4).

The forests of *Pinus peuce* in Europe belong in the mountainous and sub-alpine nemoral coniferous vegetation. These forests were much or less formed under the Mediterranean influence. Therefore, and due to historical reasons associated with the last icing, that vegetation can be considered endemic. The spread of the forests of macedonian pine is like insularity. They can be found in the highest mountain systems of Montenegro, Kosovo, Albania, Macedonia, and Bulgaria. The data on the natural habitat of macedonian pine in Bosnia and Herzegovina needs confirmation through genetic studies [65]. Vertically and environmentally the macedonian pine forests occupy mostly areas above beech and spruce forests and form the upper timberline, mainly on acid substrates, and less frequently on basic rocks.

The areal of macedonian pine forests include the Rila-Rhodope massif, Stara Planina, Voras, Shar Planina, Mokra gora and others. Their total area is about 96 000 ha [66], of which about 250 ha is territory of Central Stara Planina.

Various of authors have contributed in the syntaxonomy of the forests of *Pinus peuce* in the Balkans [6, 67, 68, 69, 70, 71, 72, 73, etc.]. The syntaxonomy of phytocoenoses of macedonian pine in Bulgaria is developed using only the dominant method of Russian-Scandinavian Phytocoenological School [14, 16, 74, 75]. The presence of the typical species *Gentiana lutea*, *Knautia midzorensis*, *Verbascum longifolium* ssp. *Pannosum* and the floristic composition of some phytocoenoses are a reason to assume their distribution in this country, at least partly in Rila and Pirin. The association *Gentiano luteae-Pinetum peucis* described by Pelister in Macedonia and the typical species of the endemic for the Balkan alliance *Pinion peucis* are found



everywhere in phytocoenoses of *Pinus peuce* in Bulgaria [76].

Studies show that since 1970 there is a remarkable ascent of the macedonian pine in Rila and Stara Planina, which is mainly associated with climate changes and the increasing temperatures and weakened anthropogenic pressure [16, 77].

3.2.1 Ass. *Adenostylo alliariae-Pinetum peucis* ass. nova hoc loco (group № 4)


**Diagnostic species:** *Pinus peuce*, *Adenostyles alliariae*, *Calamagrostis arundinacea*, *Homogyne alpina*.

**Constant species:** *Picea abies*, *Athyrium filix-femina*, *Dryopteris filix-mas*, *Hieracium murorum* gr., *Luzula luzuloides*, *Luzula sylvatica*, *Mycelis muralis*, *Prenanthes purpurea*, *Rubus idaeus*, *Senecio nemorensis*, *Sorbus aucuparia*, *Stellaria nemorum*, *Vaccinium myrtillus*.

**Dominant species:** *Pinus peuce*, *Picea abies*, *Pinus sylvestris*, *Sorbus aucuparia*, *Festuca drymeja*, *Luzula sylvatica*. **Nomenclature type (holotypus):** Relève 28, Annex I.

The comparative analysis of the total species composition, of the diagnostic species and of the environmental conditions in relation to the previously described syntaxons allows us to describe a new association - *Adenostylo alliariae-Pinetum peucis*.

The distribution of its communities is in the range of 1600-1900 m.a.s.l., but mostly at about 1700 m.a.s.l. The slopes are an average 35 degrees. The exposures are varied, but dominated by those with westerly and northerly component. South of Vezhen peak exposures are with southern components (Fig. 6). The rocks are acidic – south bulgarian granite, diorite and others. The soils are mainly Umbric Cambisols, medium rich to rich, medium deep, sandy, stony, wet. Compared with other boreal mountain forests in Central Stara Planina, the communities of *Pinus peuce* grow in the most humid and rich soils (Group № 4, Fig.2). The average annual temperature is about 3 degrees.



**Figure 6:** Ass. *Adenostylo alliariae-Pinetum peucis*

The tree layer cover is an average 60%. Dominants in various ratios, mostly prevailed by *Pinus peuce* are *Pinus peuce*, *Picea abies*, *Pinus sylvestris* and *Abies alba*. The cover of the shrub layer is generally low (below 10%). *Sorbus aucuparia* and *Rubus idaeus* are most common. The grass layer has a cover of on average 40%. Dominant

and constant species most often are *Athyrium filix-femina*, *Adenostyles alliariae*, *Diphyscium foliosum*, *Gentiana asclepiadea*, *Gymnocarpium dryopteris*, *Homogyne alpina*, *Lamium garganicum*, *Calamagrostis arundinacea*, *Stellaria nemorum* (Annex I).

Typical physiognomic feature of these forests are dense facies of *Adenostyles alliariae* in the grass layer, especially on a wet places and near streams.

In phytogeographical spectrum the boreal elements dominate (56%), which is determined by the environmental conditions, especially by the acidic substrates and the altitude. Follow by european and balkan (12%), mediterranean (8%), arctic-alpine, euro-mediterranean and cosmopolitans (4% each).

The association *Adenostyles alliariae-Pinetum peucis* has a relatively hygrophylous character and species that are not found in other known associations are present here - *Adenostyles alliariae*, *Cirsium appendiculatum*, *Heracleum verticillatum*, *Veronica beccabunga*, *Stellaria nemorum*, *Gymnocarpium dryopteris*, *Dryopteris dilatata*, *Senecio nemorensis*, *Diphyscium foliosum*. Moreover, other species, described in other associations for different areas, cannot be found here: *Gentiana lutea*, *Lilium carnolicum*, *Doronicum columnae*, *Bruckenthalia spiculifolia* and *Knautia midzoensis* for association *Gentiano luteae-Pinetum peucis* Em 1960; *Alyssum bertolonii*, *Dactylorhiza saccifera*, *Daphne blagayana*, *Minuartia baldaccii* and *Thymus boissieri* for association *Pinetum peucis* Janković 1959; *Digitalis viridiflora*, *Vaccinium uliginosum*, *Helleborus cyclophyllus* and *Astragalus glycyphyllos* for association *Digitali viridiflorae-Pinetum peucis* Em 1960; *Wulfenia carinthiaca*, *Potentilla ternata*, *Geum montanum* for association *Wulfenio carinthiaca-Pinetum peucis* Blečić et Tatić 1957; *Ajuga pyramidalis*, *Galium rotundifolium*, *Briza media*, *Platanthera bifolia* for Association *Ajugo pyramidalis-Pinetum peucis* Janković et R. Bogojević 1962, etc. [6, 58, 78].

The monodominant and mixed communities of macedonian pine in Central Stara Planina are located in separate mountain localities – north of Vezhen peak and Kamenitza peak. Scattered small groups are developing around Yumruka peak. In the recent past they had been developing into lower altitudes on inaccessible steep terrains [16]. In the recent years the lower limit has risen, where the macedonian pine is replaced primarily by beech. The upper limite has been higher as well and it corresponds to the upper timberline of the forest.

### 3.3 Conservation

Boreal mountain forests in Central Stara Planina are located within the National Park "Central Balkan". This is the second largest national park in Bulgaria and one of the most valuable park and one of the largest protected areas in Europe – second category according to the International Union for Conservation of Nature (IUCN). The territory of the park overlaps with the territory of the Special Protection Area – BG 0000494 Central Balkan (NATURA 2000), and about 45 000 ha of forest habitats are preserved there. The studied boreo-mountain forest communities belong to three habitat types listed in Annex I of the Habitats Directive 92/43/EEC and to 3 types of

habitats of the Red Book of Bulgaria – section 3 Natural habitats [36].

The spruce forests (ass. *Ranunculo carthusianae-Piceetum abietis*, comm. of *Picea abies* and *Luzula sylvatica*) are related to habitat 9410 Acidophilous *Picea* forests of the montane to alpine levels (*Vaccinio-Piceetea*) and habitat 34G3 Norway spruce (*Picea abies*) forests with conservation category Nearly threatened [63].

The forests dominated by common birch (comm. of *Betula pendula*) are related to habitat 25G1 Birch (*Betula pendula*) forests with conservation category Nearly threatened [79].

Macedonian forests (ass. *Adenostylo alliariae-Pinetum peucis*) are identified with habitat 95A0 High oro-Mediterranean pine forests and 38G3 Macedonian pine (*Pinus peuce*) forests – Endangered habitat [77].

#### 4 CONCLUSION

The boreo-mountain forests on the territory of Central Stara Planina are fragmented, forming a separate belt only at the highest parts. The described new associations with specific species composition show their relict character, their geographical isolation and the specific environmental conditions.

Early stages of forest dominated by spruce, macedonian pine, silver birch and scots pine, which displace mainly heaths and bushes of bilberries and siberian juniper are developing in the upper timberline. Particularly active in this process is the macedonian pine that is successfully spreading on mountain scree, rock landslips and steeper slopes, on places where spruce forests cannot grow.

The results obtained from this research will be a sound basis for making correct management decisions. They will be of great help at monitoring the forest succession in the highest part of the mountain.

#### 5 REFERENCES

- [1] Pavlov, D., Dimitrov, M., Phytocoenology. Univ. of Forestry Press, Sofia, (2012), 283. (in Bulgarian)
- [2] Ozenda, P., Perspectives pour une Geobiologie des montagnes. Lausanne, PPUR, (2002), 195.
- [3] Zahariev, B., Donov, V., Petrunov, K., Masarov, S., Forest vegetation zoning of Bulgaria. Zemizdat, (1979), 197. (in Bulgarian)
- [4] Pavlov, D., Phytocenological basics of forest typology In Bulgaria. DSc. thesis, Univ. of Forestry, Sofia, (1998), 298. (in Bulgarian)
- [5] Roussakova, V., Vegetation belts in Rila Mts and their place in a general Alpine system. Problems of Geography, 3-4, (2001), 75-93.
- [6] Horvat, I., Glavac, V., Ellenberg, H., Vegetation Südosteuropas. Geobotanica lecta. Fischer Verlag, Stuttgart, (1974), 768.
- [7] Vassilevič, V., Plant associations of spruce forests of the North-West. Bot. Zhurn. U.S.S.R. 68, 12, (1983), 1604–1612. (in Russian)
- [8] Oberdorfer, E., Pflanzensoziologische Exkursionsflora für Deutschland und angrenzende Gebiete. 8. Aufl. Stuttgart (Ulmer) Stuttgart, (2001), 1051
- [9] Sochava, V., Model of the new vegetation map of the world. In: Geobot. Mapp. M.–L., (1964), 3–16. (in Russian)
- [10] Ellenberg, H., Kloetzli, F., Waldgesellschaften und Waldstandorte der Schweiz. Mitteilungen. Bd./Vol. 48, Heft/Fasc.4, Zurich, (1972), 930.
- [11] Zukrigl, K., Montane und subalpine Waldgesellschaften am Alpenstrand. Mitteilungen der forstlichen Bundes Versuchsamtes, Wien (1973), 101.
- [12] Ewald, J., Die Bergmischwälder der Bayerischen Alpen - Soziologie, Standortbindung und Verbreitung. Diss. Bot. 290, (1997), 234.
- [13] Bonchev, S., Localities of coniferous phytocoenoses in Teteven area. Forestry ideas, 1-6, (1939). (in Bulgarian)
- [14] Penev, N., Georgiev, A., Macedonian pine (*Pinus peuce*) in Central Stara Planina Mountain. Izv. Bot. Inst. Sofia, 6, (1958), 215-282. (in Bulgarian)
- [15] Kochev, H., Forest phytocoenoses in the valley of Cherni Osam River (Central Balkan mountain range). Bull. Inst. Bot. Sofia, 19, (1969), 9-62. (in Bulgarian)
- [16] Velčev, V., Phytozoölogisches untersuchungen des *Pinus peuce* Griseb. in der Mittleren Stara planina. Act. Bot. Inst. 24, (1973), 193-198. (in Bulgarian)
- [17] Michalik, S., La repartition des hetraies suivant le relief du terrain sur les pentes du Nord de Stara Planina dans le reserve Boatin. Colloques phytosociologiques, 13, (1985), 867-874.
- [18] Michalik, S., Plant communities in the Boatin biosphere reserve on the northeren slopes of Stara Planina Mts (Central Bulgaria). Ochrony przyrody, 47, (1990), 9-36.
- [19] Michalik, S., The upper beech forest line in the Steneto Biosphere Reserve on the northern slopes of the Stara Planina Mts. (Central Bulgaria). Acta Soc. Bot. Pol. 61, (1992), 273-280.
- [20] Michalik, S., Badania geobotaniczne i ekologiczne w gorach Starej Planiny (Bulgaria). Wiadomosci Botaniczne, 37 (3/4), (1993), 73-75.
- [21] Apostolova, I., Meshinev, T., Biodiversity of plant communities in representative sections of forest area in the National Park "Central Balkan". In: Sakalyan, M. (ed.), Biodiversity of National Park "Central Balkan" PENSOFT, (1999), 142-168.
- [22] Pachedjieva K., Dimitrov D., Lyubenova M., Floristic investigation of plant communities in the reserve "Kamenshtitsa", Central Stara Planina Mountain. In: Proceedings of the 2<sup>nd</sup> Congress of Ecologists of the Republic of Macedonia with International Participation, Ohrid, Macedonia, 25 – 29.10.2003, (2003), 458 – 466.
- [23] Tzonev, R., Dimitrov, M., Chytry, M., Roussakova, V., Dimova, D., Gushev, C., Pavlov, D., Vulchev, V., Vitkova, A., Gogushev, G., Nikolov, I., Borisova, D., Ganeva, A., Beech forests communities in Bulgaria. Phytocoenologia, Stuttgart, 36 (2), (2006), 247-279.
- [24] Meusel, H., Jager, E., Vergleichende Chorologie der Zentraleuropäischen Flora, Bd. III. Gustav Fischer Verlag, Jena-Stuttgart-New York, (1992).
- [25] Mirchev, S., Climate. In: Management Plan for the National Park "Central Balkan" 2016–2025 Project № DIR-593212-1-8, (2016), 31-33. (in Bulgarian)

- [26] Petrova, R., Soils. In: Management Plan for the National Park "Central Balkan" 2016–2025 Project № DIR-593212-1-8, (2016), 37-39. (in Bulgarian)
- [27] Braun–Blanquet, J., Pflanzensoziologie – Grundzuege der Vegetationskunde. Wien, N.Y., 3 Aufl., Springer-Verlag (1964), 865.
- [28] Maarel, E., Transformation of cover-abundance in phytosociology and its effects on community similarity. *Vegetatio*, 39-2, (1979), 97-114.
- [29] Ellenberg H., Weber H.E., Düll R., Wirth V., Werner W., Paulissen D., Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobot.*, 18 (1991), 1-248.
- [30] Delipavlov, D., Cheshmedjiev I. (eds.), Manual to the Plants in Bulgaria. Plovdiv, Akad. Ed. Agricultural University, (2003), 590. (in Bulgarian)
- [31] Assyov, B., Petrova, A., Dimitrov, R., Vassilev, R., Conspectus of the Bulgarian vascular flora. Distribution maps and floristic elements. *BBF*, 4<sup>th</sup> ed., (2012), 488. (in Bulgarian)
- [32] Weber, H., Moravec, J., Theurillat, J.-P., International Code of Phytosociological Nomenclature. 3<sup>rd</sup> edition. *J. Veg. Sci.*, 11, (2000), 739-768.
- [33] Rodwell, J.S., Schaminée, J.H.J., Mucina, L., Pignatti, S., Dring, J., Moss, D., The diversity of European Vegetation. An overview of phytosociological alliances and their relationship to EUNIS habitats. Wageningen, NL, Report EC-LNV 2002/054, (2002), 168.
- [34] Mucina, L., Bültmann, H., Dierßen, K., Theurillat, J.-P., Raus, T., Čarni, A., Šumberová, K., Willner, W., Dengler, J., Gavilán García, R., Chytrý, M., Hájek, M., Di Pietro, R., Iakushenko, D., Pallas, J., Daniëls, F. J. A., Bergmeier, E., Santos Guerra, A., Ermakov, N., Valachovič, M., Schaminée, J. H. J., Lysenko, T., Didukh, J.P., Pignatti, S., Rodwell, J. S., Capelo, J., Weber, H. E., Solomeshch, A., Dimopoulos, P., Aguiar, C., Freitag, H., Hennekens, S. M., Tichý, L., Vegetation of Europe Hierarchical floristic classification system of vascular plant, bryophyte, lichen, and algal communities. *Applied Vegetation Science*, Volume 19, (Suppl. 1), (2016), 264.
- [35] Kavrakova V., Dimitrov M. Dimova D., Tsonev R., Belev T., Rakovska R. (eds.), Guidance for identifying habitats of European importance in Bulgaria. Second revised edition. Sofia, WWF–DCP and Fed "Green. Balkans", (2009), 131. (in Bulgarian)
- [36] Biserkov, V., Gussev, Ch., Popov, V., Hibaum, G., Roussakova, V., Pandurski, I., Uzunov, Y., Dimitrov, M., Tzonev, R., Tsoneva, S., (eds.), Red Data Book of the Republic of Bulgaria. Volume 3, Natural habitats, (2015), 422.
- [37] Tichý, L., JUICE, software for vegetation classification. *Journal of Vegetation Science*, 13, (2002), 451-453.
- [38] Roleček, J., Tichý, L., Zelený, D., Chytrý, M., Modified TWINSpan classification in which the hierarchy respect cluster heterogeneity. *J. Veg. Sci.*, 20, (2009), 596–602.
- [39] MacQueen, J., Some Methods for classification and Analysis of Multivariate Observation. In: 5<sup>th</sup> Berkeley Symposium on Mathematical Statistics and Probability, Berkeley University of California Press, (1967), 281-297.
- [40] Tichý L., Holt J., JUICE program for management, analysis and classification on ecological data. Vegetation science group, Masaryk University, Brno, (2006).
- [41] Dengler, J., Berg, C., Jansen, F., New ideas for modern phytosociological monographs. *Ann. Bot.*, 5, (2005), 193-210.
- [42] Chytrý, M., Tichý, L., Holt, J., Botta-Dukát, Z., Determination of diagnostic species with statistical fidelity measures. *J. Veg. Sci.*, 13 (2002), 79–90.
- [43] Chytrý, M., Tichý, L., Diagnostic, constant and dominant species of vegetation classes and alliances of the Czech Republic a statistical revision. *Folia Facultatis Scientiarum Naturalium Universitatis Masarykianae Brunensis*, 108, (2003), 1-231.
- [44] StatSoft, Inc. STATISTICA (data analysis software system), version 10.0. [www.statsoft.com](http://www.statsoft.com), (2011).
- [45] Coldea, G., Muntii Rodnei. *Studiu geobotanic*. Bucuresti, Ed.Ac.R., (1990), 172.
- [46] Coldea, G., Prodrome des associations végétales des Carpates du Sud-Est (Carpates Roumaines). *Doc. Phytosociol.*, 13, (1991), 317–539.
- [47] Ivan, D. (ed.), *Vegetatia Romaniei*. Bucuresti, Ed. Tehnica Agricola, (1992), 410.
- [48] Ivan, D., Doniță, N., Coldea, G., Sanda, V., Popescu, A., Chifu, T., Boșcaiu, N., Mititelu, D., Pauca-Comanescu, M., *Vegetation potentielle de la Roumanie*. *Braun-Blanquetia*, 9, (1993), 78.
- [49] Coldea, G., Wagner, I., *Beitrag zum studium der vegetation Rumaniens (III)*. Cluj, *Contributii Botanice*, (1997-1998), 81-87.
- [50] Kato, Z., Vegetation of the Fagul peatbog and the neighbouring swamps the Apa Rosie basin (Covasna country, Romania). *Contributii Botanice*, I, (1999-2000), 65-80.
- [51] Pop, I., Cristea, V., Hodișan, I., *Vegetatia judetului Cluj (Studiu fotocenologic, ecologic, biocenologic, bioeconomic si eco-protectiv)*. *Contributii Botanice*, (1999-2000), 35, 2, (2002), 5-254.
- [52] Mišić, V., Jovanović-Dunjić, R., Popović, M., Borisavljević, Lj., Antić, M., Dinić, A., Danon, J., Blaženčić, Ž., Biljne zajednice i staništa Stare planine [Plant communities and habitats of the Stara planina Mt]. – Posebna izdanja 511, odeljenje prirodno-matematičkih nauka 49, SANU, Beograd, (1978), 389.
- [53] Jovanović, B., Sumske fitocenoze i stanista Suve Planine. *Glasnik Sumarskog F-ta, ser. Sumarstvo*, posebno izdanje 55, (1980), 216.
- [54] Jovanović, B., Lakušić, R., Rizovski, R., Trinjastić, I., Zupančić, M., *Prodromus phytocoenosum Jugoslaviae. Ad mappam vegetationis M 1200 000. Bribir-Ilok*, (1986), 46.
- [55] Dzekov, B., Jovanović, B., Jovanović, R., Lakušić, R., Mišić, V., Parabuški, I., Puncer, I., Rexhepi, F., Rizovski, R., Srefanović, V., Šugar, I., Trinajstić, I., Vučković, M. & Zupančić, M. (eds.), *Natural potential vegetation of Yugoslavia*. Ljubljana, *Do paralele Publ.*, (1986), 122.
- [56] Gajić, M., *Flora i vegetacija Golije i Javora*. Šumarski fakultet, Beograd, (1989), 520.
- [57] Kojić, M., Popović, R., Karadžić, B., *Sintaksonomski pregled vegetacije Srbije*. Beograd. Inst. za biol. istraž. "S. Stanković" Publ., (1998), 218.
- [58] Tomić, Z., Rakonjac, Lj., Šumske fitocenoze Srbije. Institut za šumarstvo i Univerzitet Singidunum,

- Fakultet za primenjenu ekologiju Futura, Beograd, (2013), 1-177.
- [59] Dafis, S., Classification of forest vegetation of Greece. Min.of Agriculture - Gen. Direct. of Forests, Publ. No 36, Athens, (1976), 22.
- [60] Dimitrov M., Floristic classification of forest vegetation in the Yundola Forestry Experimental Station. Forestry ideas, 1-4 (2003), 23-46. (in Bulgarian).
- [61] Roussakova, V., Valchev, V., Biodiversity of plant communities in the National Park "Rila" to the west of the Maliovitza River. In: Sakalyan, M. (ed.), Biodiversity National Park "Rila", PENSOFT, (1999), 194-231.
- [62] Roussakova, V., Dimitrov, M., Caractéristiques phytosociologiques des forêts à *Picea abies* et *Abies alba* de la montagne de Rila (Bulgarie). Acta Bot. Gallica, 152, 4, (2005), 563-571.
- [63] Roussakova, V. & Dimitrov, M., 34G3 Norway spruce (*Picea abies*) forests. In: Biserkov, V., Gussev, Ch., Popov, V., Hibaum, G., Roussakova, V., Pandurski, I., Uzunov, Y., Dimitrov, M., Tzonev, R., Tsoneva, S. (eds.), Red Data Book of the Republic of Bulgaria. Volume 3, Natural habitats, (2015), 347-349.
- [64] Doniță, N., Ivan, D., Coldea, G., Sanda, V., Popescu, A., Chifu, T., Paucă-Comănescu, M., Mititelu, D., Boșcaiu, N., Vegetation of Romania. Tehnică Agricolă, Bucharest, (1992). (in Romanian).
- [65] Bucalo, V., Stupar, V., Milanović, D., Characteristics and origin of the population of Macedonian pine (*Pinus peuce* Griseb.) at Jadovnik Mt in Western Bosnia. Bull. Fac. of For. Univ. of Banja Luka, 16, (2012), 7-29. (in Serbian)
- [66] Bohn, U., Neuhausl, R., Gollub, G., Hettwer, C., Neuhauslová, Z., Raus, Th., Schlüter, H., Weber, H. 2000/2003, 2012 Karte der natürlichen Vegetation Europas /Map of the Natural Vegetation of Europe. Maßstab/ Scale 1:2500000. Münster (Landwirtschaftsverlag)
- [67] Blečić, V., Prilog poznavanju šumske vegetacije planine Ljubišnje, (1957) 10, 25-42.
- [68] Blečić, V., Tatić, B., Šuma molike u Crnoj Gori (*Pinetum peucis montenegrinum*), (1957), 10, 43-53.
- [69] Bondev I., The vegetation of Bulgaria. Map M 1600 000 with explanatory text. Sofia Univ. Press, Sofia, (1991). (in Bulgarian)
- [70] Dimitrov, D., Molika, *Pinus peuce* Griseb. Šumska monografija; Balk. Privr. Bibl., Sv. 1, Graficko-industrijsko preduzeće Krajnicanac, Skopje, (1933).
- [71] Em, H., Šumske zajednice četinaru u NR Makedoniji. [Pflanzengesellschaften der Nadelwälder in der V. R. Mazedonien] Biol. Glas., 15, (1962), 1-38.
- [72] Fukarek, P., Podaci o rasirenju molike (*Pinus peuce* Griseb.) na Balkanskom poluostrvu. Godisnjak Istituta Univerziteta u Sarajevu 1/2, (1997), 43-50. (in Serbian)
- [73] Janković, M., Betrachtungen Über die gegenseitigen Beziehungen der Molika (*Pinus peuce*) und Panzerkiefer (*Pinus heldreichii*) sowie auch Über ihre Ökologischen Eigenschaften, besonders in Bezug auf ihre geologische Grundlage. Glasnik Botanickog Zavoda Baste Univerziteta u Beogradu 1/2, (1960), 166-180.
- [74] Velchev, V., Phytocoenological study of overhead phytomass in two communities of *Pinus peuce* Griseb. in the Central Stara Planina. Bull. Inst. Bot. Sofia, 24, (1981), 169-193. (in Bulgarian)
- [75] Velchev, V., Roussakova, V., Ecological peculiarities and phytocoenological characteristics of *Pinus peuce* Griseb. in the Pirin and Rila Mountains. God. Sofiisk. Univ. Biol. Fak., 80, 2, (1990), 58-93. (in Bulgarian)
- [76] Roussakova, V., 38G3 Macedonian pine (*Pinus peuce*) forests. In: Biserkov, V., Gussev, Ch., Popov, V., Hibaum, G., Roussakova, V., Pandurski, I., Uzunov, Y., Dimitrov, M., Tzonev, R., Tsoneva, S. (eds.), Red Data Book of the Republic of Bulgaria. Volume 3, Natural habitats, (2015), 358-360.
- [77] Meshinev, T., Apostolova, I., Koleva, E., Influence of warming on timberline rising a case study on *Pinus peuce* Griseb. in Bulgaria. Phytocoenologia, 30, 3-4, (2000), 431-438.
- [78] Dinić, A., Janković, M., The acidophilous conifer forest. In: Škorić M. D., Vasić O. (ed.), Vegetacija Srbije 2 (2). SANU, Belgrade, (2006), 201-212. (in Serbian).
- [79] Dimitrov, M., 25G1 Birch (*Betula pendula*) forests. In: Biserkov, V., Gussev, Ch., Popov, V., Hibaum, G., Roussakova, V., Pandurski, I., Uzunov, Y., Dimitrov, M., Tzonev, R., Tsoneva, S. (eds.), Red Data Book of the Republic of Bulgaria. Volume 3, Natural habitats, (2015), 330-331.

## 6 ACKNOWLEDGEMENTS

We wish to express our thanks to the University of Forestry – Sofia for the financial support (project BG051PO001-3.3.06-0056). We are also very grateful to Dejan Mandzukovski for providing us some literature sources and to Peter Zhelev for the language revision.



**Annex I:** Syntaxonomical table of boreo-montane forest phytocoenoses in Central Stara Planina Mts.

1 - Assoc. *Ranunculo carthusianae-Piceetum abietis* ass. nova, holotypus hoc loco, rel. 8; 2 - Comm. of *Picea abies* and *Luzula sylvatica*; 3 - Comm. of *Betula pendula*; 4 - Assoc. *Adenostylo alliariae-Pinetum peucis* ass. nova, holotypus hoc loco, rel. 28 (Fidelity >20)

Syntaxa	1											2				3			4																
Relève number	1	2	3	4	5	6	7	8	9	10	11	Fidelity	12	13	14	15	Fidelity	16	17	18	Fidelity	19	20	21	22	23	24	25	26	27	28	29	30	Fidelity	
Altitude (m a.s.l.) (x 10)	169	169	178	172	170	167	169	170	168	168	169		147	160	159	161		151	145	144		162	173	164	168	175	161	170	181	183	183	179	168		
Exposition	W	W	SE	N	S	SW	W	W	S	W	SW		N	NE	NE	NE		N	SE	S		NW	NE	N	N	NE	W	N	N	N	N	NE	N		
Slope (degree)	1	10	2	15	20	10	1	3	8	1	15		50	20	20	10		2	35	40		20	35	45	35	45	35	10	30	50	45	50	10		
Sample size (m <sup>2</sup> )	400	400	400	400	400	400	400	400	400	400	400		400	400	400	400		225	225	225		400	400	400	400	400	400	400	400	400	400	400	400		400
Cover of trees (%)	45	40	45	60	60	70	90	80	60	65	70		60	90	90	90		40	75	50		55	55	70	70	60	65	65	50	40	55	70	65		
Cover of bushes (%)	60	60	4	2	3	4	20	15	10	20	4		10	0	20	5		1	0	40		10	1	2	2	3	5	10	3	3	3	1	40		
Cover of grass (%)	20	20	50	45	60	40	30	35	45	35	45		50	30	30	20		55	30	20		40	60	40	40	55	40	40	40	75	45	40	30		
Soil richness	7	7	7	5	5	5	5	5	5	5	5		5	7	7	7		5	5	3		5	7	5	7	7	5	7	5	5	5	7	7		
Soil depth	4	4	4	3	3	3	3	3	3	3	3		3	4	4	4		3	3	2		3	4	3	4	4	3	4	3	3	3	4	4		
Soil stoniness	2	2	2	2	2	2	3	2	2	3	2		3	2	3	2		3	3	4		2	3	2	3	3	2	3	3	3	3	3	3		
Soil composition	4	4	4	4	4	4	4	4	4	4	4		4	4	4	2		3	4	4		4	4	4	4	4	4	4	4	4	4	4	4		4
Soil reaction	4.5	4.7	4.9	5.1	5.2	4.9	5.2	5.8	5.6	5.1	5.2		4.4	4.2	4.9	4.6		3.5	6.1	4.4		4.4	5	4.7	4.7	5.1	4.5	4.3	4.6	4.6	4.7	4.9	4.2		
Nutrients	4.7	4.6	5	5.2	5.5	5.4	5.3	5.2	5.2	5.2	5.2		5	5	4.7	5.2		2.2	4.7	4.6		5.1	5.3	5.2	5.2	5.3	4.8	5	5.1	5.3	5.2	5.4	5.2		
Moisture	5.2	5.2	5.2	5.2	5.2	5.3	5.6	4.9	4.9	5.7	4.9		5.1	5	5.1	4.9		4.2	5.1	5		5.3	5.5	5.4	5.4	5.6	5.3	5.4	5.5	5.6	5.5	5.6	5.3		
Accessibility Q	500	500	500	500	500	500	500	500	500	500	500		650	650	650	650		700	600	700		700	700	700	700	700	700	700	700	700	700	700	700		700
Air humidity Vv	83	83	83	83	83	83	83	83	83	83	83	82	83	82	83	82	82	82	83	83	83	83	83	83	83	83	83	83	83	83	83				
Light	5.4	5.4	5.6	5.4	5.2	5.2	5.5	5.6	5.7	5.6	5.4	4.9	4.3	4.7	5.2	5.9	4.9	5.1	4.5	4.6	4.6	4.5	4.8	4.6	5.4	5.3	5.3	5.3	5.2	5.3					
Temperature	3.6	3.8	4.1	4.3	4.3	4.5	3.9	4.2	4.2	4	4	4.5	4.3	4.4	4.6	4.3	5.4	4.4	4.5	4.6	4.6	4.5	4.4	4.6	3.8	3.9	3.8	3.9	4	4					
Continentality	3.8	4.2	4.1	4	4	3.9	3.9	4.4	4.2	3.9	4.1	3.4	3.9	3.6	4.1	4.5	3.9	3.7	3.5	3.6	3.6	3.5	3.8	3.5	4.2	3.8	3.8	3.8	3.9	4.5					
<i>Ranunculus carthusianus</i>	1	.	.	r	1	1	+	+	+	+	+	88	.	.	.	.	*	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Rumex alpinus</i>	.	.	+	+	1	1	+	+	1	+	1	88	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Myosotis sylvatica</i>	1	1	+	+	+	+	+	+	+	+	+	86	.	.	.	1	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Juniperus sibirica</i> b	4	4	+	+	+	1	+	+	1	2	1	82	.	.	.	.	--	.	.	3	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Juniperus sibirica</i> c	1	2	+	+	+	+	+	+	+	+	+	82	.	.	.	.	--	2	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Ribes alpinum</i> b	.	.	+	+	1	.	+	1	1	+	+	82	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Vaccinium vitis-idaea</i>	1	.	2	1	2	1	1	2	2	1	1	75	.	.	.	.	--	2	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Fagus sylvatica</i> c	1	1	+	+	+	+	.	+	+	+	1	75	.	.	.	.	--	.	.	1	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Potentilla haynaldiana</i>	.	.	.	r	r	1	.	1	1	.	r	69	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Scilla bifolia</i>	.	1	2	+	+	+	+	2	+	1	.	68	.	.	.	.	--	.	2	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Veronica officinalis</i>	.	.	1	+	+	+	+	.	1	+	+	65	.	.	.	.	--	.	.	.	--	1	.	.	+	.	+	.	.	.	.	--			
<i>Ajuga reptans</i>	1	.	+	+	+	1	+	.	2	+	1	48	1	.	1	.	--	.	1	.	--	.	.	.	.	.	.	.	.	.	.	--			
<i>Picea abies</i> b	.	1	1	1	1	1	2	2	2	2	1	48	.	.	2	2	--	.	.	.	--	.	.	.	+	.	+	1	1	1	+	+	--		

continued

NIKOLOV I., DIMITROV M.  
BOREO-MONTANE FOREST PHYTOCOENOSES IN CENTRAL STARA PLANINA MTS.

continuation of Annex I

<i>Acer pseudoplatanus</i> b	.	.	.	.	+	1	+	.	.	.	.	47	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Veronica alpina</i>	1	.	.	.	.	.	+	.	.	.	+	47	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Veronica beccabunga</i>	.	.	.	.	.	.	2	.	.	.	2	38	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Juncus effusus</i>	.	.	.	.	.	.	.	.	.	.	1	26	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Geum coccineum</i>	.	1	.	.	.	.	.	.	.	.	.	26	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Dryopteris expansa</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	1	1	66	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Laserpitium krapfii</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	1	.	45	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Galium rotundifolium</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	1	.	45	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Cardamine pectinata</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	45	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Fragaria vesca</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	45	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Geranium pyrenaicum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	45	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Galium odoratum</i>	.	1	.	.	.	.	.	.	.	.	.	--	.	2	.	1	41	.	.	.	--	.	+	.	1	+	.	.	.	.	.	--		
<i>Abies albac</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	1	1	.	23	.	1	.	--	1	+	+	+	.	+	.	.	.	.	--		
<i>Betula pendula</i> a	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	3	1	2	78	.	.	.	.	.	.	.	.	.	.	--		
<i>Betula pendula</i> b	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	1	.	1	78	.	.	.	.	.	.	.	.	.	.	--		
<i>Chamaecytisus austriacus</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	1	1	.	78	.	.	.	.	.	.	.	.	.	.	--		
<i>Potentilla rupestris</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	2	.	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Poa nemoralis</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	52	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Crocus veluchensis</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	1	.	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Corylus avellana</i> c	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	52	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Asarum europaeum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	1	.	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Acer pseudoplatanus</i> a	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	+	52	.	.	.	--	.	.	.	.	.	.	.	.	.	.	--		
<i>Acer pseudoplatanus</i> c	1	.	.	+	+	+	.	.	.	+	.	--	.	.	.	.	--	.	1	1	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Acer heldreichii</i> c	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	1	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Epilobium montanum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	1	.	52	.	.	.	.	.	.	.	.	.	.	--		
<i>Geranium macrorrhizum</i>	.	.	.	.	.	.	.	.	.	.	.	--	2	.	.	.	--	.	1	.	31	.	.	.	.	.	.	.	.	.	.	--		
<i>Abies alba</i> a	.	.	.	.	.	.	.	.	.	.	.	--	2	.	.	.	--	.	.	2	21	.	.	2	.	.	3	.	.	.	.	--		
<i>Luzula luzuloides</i>	1	1	1	+	2	.	+	.	.	+	.	--	1	1	.	1	1	1	2	2	52	2	+	1	1	1	1	1	+	+	+	1	23	
<i>Calamagrostis arundinacea</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	1	--	.	.	.	--	1	+	r	+	+	+	+	+	r	+	1	86	
<i>Hieracium murorum</i> gr.	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	1	--	+	+	1	+	+	+	+	+	+	1	82		
<i>Dryopteris dilatata</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	2	1	2	2	.	.	1	2	1	1	78	
<i>Gentiana asclepiadea</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	+	+	+	+	+	.	1	+	+	.	78	
<i>Adenostyles alliariae</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	2	.	2	+	.	1	1	2	2	2	78	
<i>Stellaria nemorum</i>	.	.	.	2	+	2	+	.	.	+	.	--	.	.	.	.	--	.	.	.	--	1	+	+	+	+	+	1	+	1	+	2	1	76
<i>Lamium garganicum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	1	+	+	+	r	+	.	.	.	.	r	72	
<i>Homogyne alpina</i>	1	.	2	2	.	.	+	.	.	r	.	--	.	.	.	.	--	.	.	.	--	1	+	+	+	+	+	+	+	1	1	+	70	
<i>Pinus peuce</i> a	.	.	.	.	.	.	.	.	.	.	.	--	.	5	.	3	--	.	.	.	--	2	3	2	2	4	2	2	2	3	3	3	2	68

continued

<i>Festuca drymeja</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	1	+	1	+	+	3	.	.	.	.	.	.	66
<i>Abies albab</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	2	+	+	+	+	2	.	.	.	.	.	.	66
<i>Pinus peuce b</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	+	.	.	1	1	1	+	.	59
<i>Pinus peucec</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	+	+	+	+	+	.	59	
<i>Gymnocarpium dryopteris</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	+	.	1	+	.	.	.	+	.	r	.	59
<i>Mycelis muralis</i>	.	.	+	+	+	1	+	.	.	+	.	--	.	.	1	1	--	.	.	.	--	1	+	+	+	+	+	+	+	+	+	+	1	56
<i>Prenanthes purpurea</i>	1	1	.	.	.	.	.	.	.	.	.	--	1	.	1	.	--	.	.	.	--	.	1	1	1	1	.	+	1	+	+	+	1	54
<i>Lerchenfeldia flexuosa</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	r	+	r	.	1	.	.	.	.	.	.	52
<i>Diphyscium foliosum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	+	.	.	+	.	.	.	.	.	+	.	45
<i>Cirsium appendiculatum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	+	1	1	.	.	45
<i>Geranium sylvaticum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	+	+	.	.	36	
<i>Dryopteris filix-mas</i>	1	.	r	+	+	+	+	+	1	.	1	--	.	1	1	1	--	.	.	1	--	1	1	+	+	1	1	+	1	1	1	+	1	36
<i>Pinus sylvestris a</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	3	.	.	.	.	.	3	36
<i>Pinus sylvestris b</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	+	.	.	.	.	+	36	
<i>Pinus sylvestris c</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	+	.	.	.	.	1	36	
<i>Lonicera xylosteumb</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	+	.	.	.	.	.	.	25	
<i>Heracleum verticillatum</i>	.	.	.	.	.	.	.	.	.	.	.	--	.	.	.	.	--	.	.	.	--	.	.	.	.	.	.	.	+	.	.	.	25	
<i>Luzula sylvatica</i>	1	1	.	+	.	.	+	.	.	+	.	--	2	2	.	1	23	.	.	.	--	1	3	2	2	3	1	2	2	2	2	2	2	52
<i>Senecio nemorensis</i>	1	1	+	+	1	1	+	+	2	+	1	35	2	1	1	2	35	.	.	.	--	1	1	+	1	+	.	2	2	2	2	+	1	24
<i>Picea abies a</i>	3	3	3	4	4	4	5	5	4	4	4	26	3	2	5	3	26	.	4	+	--	4	2	3	3	2	+	2	2	2	2	2	2	26
<i>Picea abies c</i>	1	1	+	+	+	.	+	1	1	+	+	50	.	.	.	.	--	.	.	.	--	1	+	+	+	+	+	+	+	1	+	+	1	60
<i>Rubus idaeus</i>	1	1	1	1	1	1	+	+	+	r	1	60	.	.	.	.	--	.	.	1	--	.	+	+	+	1	+	+	+	+	+	+	1	41
<i>Athyrium filix-femina</i>	1	1	+	.	2	+	+	.	.	+	+	40	.	.	.	.	--	.	.	.	--	.	1	+	1	1	.	+	+	2	1	+	1	52
<i>Sorbus aucupariac</i>	1	1																																

**Location:** 1- 42° 42' 45.8964", 24°43' 48.6156"; 2 - 42°42' 58.0248", 24°42' 52.1100"; 3 - 42°42' 28.2960", 24°44' 22.6392"; 4 - 42°42' 47.3868", 24°43' 55.9704"; 5 - 42°42' 28.2960", 24°44' 22.6392"; 6 - 42°42' 20.2932", 24°44' 7.7856"; 7 - 42°42' 46.3320", 24°43' 53.2344"; 8 - 42°42' 46.1880", 24°43' 53.8608"; 9 - 42°42' 43.0308", 24°44' 0.4056"; 10 - 42°42' 46.5300", 24°43' 53.1876"; 11 - 42°42' 38.7972", 24°44' 4.0740"; 12 - 42°44' 3.7019", 24°39' 36.1892; 13 - 42°46' 29.4528", 24°23' 27.6432"; 14 - 42°46' 34.5072", 24°23' 25.0368"; 15 - 42°46' 34.5072", 24°23' 25.0368"; 16 - 42°43' 34.5108", 24°40' 28.0020"; 17 - 42°43' 56.2584", 24°39' 25.9992"; 18 - 42°43' 55.2828", 24°39' 42.2100"; 19 - 42°46' 15.3408", 24°23' 23.1036"; 20 - 42°45' 52.2036", 24°23' 17.0556"; 21 - 42°45' 59.0580", 24°23' 23.5176"; 22 - 42°45' 54.4716", 24°23' 19.5108"; 23 - 42°45' 52.4340", 42°45' 52.4340"; 24 - 42°46' 8.4432", 24°23' 25.4940"; 25 - 42°46' 12.4824", 24°23' 36.7368"; 26 - 42°45' 53.3196", 24°23' 37.2192"; 27 - 42°45' 46.4796", 24°23' 24.3276"; 28 - 42°45' 49.8672", 42°45' 49.8672"; 29 - 42°45' 46.0080", 24°23' 15.8496"; 30 - 42°46' 13.3536", 24°23' 37.8708.

**Date:** 25.06.2013 -1-11; 12.06.2011 -12; 26.06.2012 - 17; 09.07.2011 -13-15; 18.06.2013 - 16; 07.06.2013 - 18; 11.07.2013 -19-30.

## ANALYSIS OF FLORISTIC COMPOSITION OF MOUNTAIN BEECH FOREST ON LIMESTONE AND SERPENTINE IN SERBIA

NOVAKOVIĆ-VUKOVIĆ M., MILOŠEVIĆ R.

University of Belgrade, Faculty of Forestry, Belgrade, Serbia

Corresponding author e-mail address: marijana.novakovic@sfb.bg.ac.rs

**ABSTRACT:** The paper is floristic composition comparative research of mountain beech forests on serpentine in southwestern Serbia, and beech forests on limestone in eastern Serbia. The analysis showed that there are significant differences in the floristic composition. On the serpentine bedrock a low count of plant species was recorded, Jacquard's similarity index of studied stands had a very low value. Within phytocoenological relevés made on limestone bedrock, the most typical species of beech forests appear as isolated, while in the beech forests on serpentine, xerophyl species characteristic to the order *Erico* - *Pinetalia* and *Quercetalia pubescentis* were observed, in addition to "fagetal" species. In the spectrum of life forms on both the geological substrate hemicryptophytes dominated, and phanerophytes and nanophanerophytes of serpentinite had the significant presence, as a result of warmer climate in southwestern Serbia. In geofloristic spectrum, a group of submediterranean floral elements, which are connected with Balkan floral elements, is as twice as numerous in the mountain beech forest on serpentine than on limestone, which is another indicator of southwestern Serbia's exposure to submediterranean influence.

**Keywords:** beech, serpentine, limestone, southwest Serbia, eastern Serbia.

### 1 INTRODUCTION

Beech is the most ubiquitous tree species in Serbia with the widest altitudinal distribution, arising from a broad ecological amplitude in regard to climatic factors (light and temperature), and edaphic factors (geological substrate and soil types), and slightly lower amplitude compared to the humidity of habitat [8]. It occurs in most parts of Serbia, except in the Pannonian Plain, with the exception of Fruška Gora and Vršac Mountains [2]. On the territory of Serbia beech forests cover 660,400 ha (29.3% of the total covered area), out of which is 64.7% is in government property [1]. Considering that the beech communities are widespread in almost all mountain massifs in Serbia, their soils were formed on different rocks which vary in textural characteristics, chemical and mineral composition, as well as in resistance and products created in their degradation process [5]. Within beech forests, several basic types of substrate can be defined, considering the importance of the parent substrate for the genesis and soil properties [6]: carbonate rocks, acidic and neutral silicate rocks and mafic silicate rocks. Beech forests have very diverse botanical composition, considering their prevalence and effect of orographic-edaphic factors, so they include a large number of phytocoenoses. One of them is a mountain beech forest (*Asperulo odoratae-Fagetum moesiacaе* B. Jovanović in 1973., Syn. *Fagetum moesiacaе montanum* B. Jovanović 1953.; *Fagetum montanum asperuletosum* B. Jovanović 1973), which represents a powerful climate-regional belt of vegetation on the territory of Serbia, as well as the most abundant and economically important type of beech forest and Serbian woodland in general [11]. Aim of the paper was to determine the extent of the differences in the floristic composition between mountainous beech forests on serpentine and limestones, which would contribute to a better knowledge of beech forests on different geological substrates, since these forests are of great importance for Serbia.

### 2 MATERIAL AND METHOD

The study of mountain beech forests floristic composition (*Asperulo odoratae-Fagetum moesiacaе* B. Jovanović 1973) was based on 24 phytocoenological relevés that were obtained from literature. 14 phytocoenological relevés were sampled in the mountain beech forest on serpentine bedrock on Crni vrh near Priboj [12], another 10 were sampled in the mountain beech forest on Mount Ozren - Sokobanja, on a limestone substrate [3]. Syntaxonomic names follow [16]. CA vegetation data analysis was performed using the statistical software CANOCO 4.5 [9]. Indexes of diversity and equivalence (Shannon -Wiener diversity index and Evenness) were calculated in the program JUICE 7.0 [15]. Jaccard's similarity index of investigated stands was also calculated [10]. The spectrum of floral elements was made according to the systematization of phytogeographical elements [4], and spectrum of life forms [7].

### 3 RESULTS AND DISCUSSION

64 species were recorded in mountain beech forest on serpentine and 77 species in beech forest on limestone. The conclusion is that the beech forest on limestone is floristically richer than on the serpentine, although the analysis included a smaller number of phytocoenological relevés on limestone than in the serpentine [14] compared the vegetation of limestone and serpentine in Bosnia and came to the conclusion that the serpentine, unlike limestone, sees great uniformity in terms of growing vegetation and poorness in the number of species. It should be noted that the serpentine bedrock is very unfavorable for the development of the plant world, on which only plants that have managed to adapt to life on this inhospitable bedrock can survive. The same conclusion is reached by analyzing the floristic composition of beech forests on serpentine and limestones in Serbia. Floristic composition analysis of mountain beech forests on Pešter, on limestone and silicate bedrock, also showed greater floristic richness of this forest on limestone, but this phenomenon is explained by higher degradation and more open set of studied stands, which enables settlement to a large number of species [13].


Mountain beech forests on both serpentinite and limestone within studied stands occupy mainly same expositions - northern and northwestern. They are also located on similar inclinations - inclinations on serpentinite are in 5-40° range, while inclinations on limestone are in 0-30° range. Mountain beech forests on serpentinite are registered in 560-800 m elevation range, while stands on limestone are registered on elevations between 810-870 m. Based on all of the above-mentioned, it can be concluded that geological substrate is the primary factor that affects floristic composition in the studied stands of mountain beech forest on serpentinite and limestone. Jaccard's index of similarity of studied stands [10] is:

$$J = \frac{UV}{V + U - UV} = 0.24, \text{ where is}$$

UV- the number of common species,  
U, V- the number of species in mountain beech forest on serpentinite and limestone, respectively.

Jaccard's similarity index shows that although this is the same community, floristic similarity among the studied stands is low.

Ordinate system (Fig. 1) shows that, although it is one plant community, phytocoenological relevés diverge depending of the geological basis. Within phytocoenological relevés made on limestone bedrock, in the negative part of the coordinate system, "fagetal" species appear as isolated: *Asperula odorata*, *Isopyrum thalictroides*, *Cardamine bulbifera*, *Anemone ranunculoides*, *Salvia glutinosa* and others. On the other hand, within the beech forests on serpentinite, in addition to "fagetal" species, xerophyl species characteristic for *EricoPinetalia* and *Quercetalia pubescentis* orders: *Fraxinus ornus*, *Campanula patula*, *Polygonatum odoratum*, appear on the graph.



**Figure 1:** CA ordination biplot, fit range for the species 30-100%, 42 species (○-relevé representation, △-species representation; (1-14)- relevés on serpentinite; (15-24)- relevés on limestone

Species abbreviations: *Ace cam*-*Acer campestre*; *Mel uni*-*Melica uniflora*; *Gal lut*-*Galeobdolon luteum*; *Iso tha*-*Isopyrum thalictroides*; *Geu urb*-*Geum urbanum*; *Cor ave*-*Corylus avellana*; *Gal vern*-*Galium vernum*; *Ane ran*-*Anemone ranunculoides*; *Asp odo*-*Asperula odorata*; *Car bul*-*Cardamine bulbifera*; *Hel odo*-*Helleborus odoratus*; *Gal apa*-*Galium aparine*; *Cra mon*-*Crataegus monogyna*; *Fag moe*-*Fagus moesiaca*; *Epi alp*-*Epimedium alpinum*; *Gen asc*-*Gentiana asclepiadea*; *Vib lan*-*Viburnum lantana*; *Cam pat*-*Campanula patula*; *Gle hir*-*Glechoma hirsuta*; *Hep nob*-*Hepatica nobilis*; *Sal glu*-*Salvia glutinosa*; *Pul off*-*Pulmonaria officinalis*; *Her sph*-*Heracleum sphondylium*; *Cor mas*-*Cornus mas*; *Fra orn*-*Fraxinus ornus*; *Vac myr*-*Vaccinium myrtillus*; *Mer per*-*Mercurialis perennis*; *Pol odo*-*Polygonatum odoratum*; *Sax rot*-*Saxifraga rotundifolia*; *Sed tel*-*Sedum telephium*; *Pte aqu*-*Pteridium aquilinum*; *Evo eur*-*Evonymus europaeus*; *Cet off*-*Ceterach officinarum*; *Cor col*-*Corylus colurna*; *Evo ver*-*Evonymus verrucosa*; *Ger mac*-*Geranium machrorrhizum*; *Phy sco*-*Phyllitis scolopendrium* (the abbreviations following the species denotes A-tree layer, B –shrub layer, C-ground flora layer)

Shannon Wiener index (Table I) shows significant differences between the studied stands. This index has a much higher value in the mountain beech forest on limestone than on serpentinite, which is logical if we consider the number of recorded species, which is larger in the stands on limestone. On the other hand, Evenness index shows approximately equal value (Table I), provided that it's slightly higher in the mountain beech forest on limestone. Based on Evenness index it can be concluded that plant species do not have regular spatial distribution. As species that reduce the value of this index, we can mention the ones that are characterized by great number and coverage in a certain number of relevés: *Asperula odorata*, *Allium ursinum*, *Geranium machrorrhizum*, *Cardamine bulbifera* etc.


**Table I:** Indices of the diversity and evenness of mountain beech forest at the study sites

	Average Shannon-Wiener index	Average Evenness index
serpentinite	1.37	0.55
limestone	2.24	0.60

The life form spectrum (Fig. 2) reveals significant differences. Hemicryptophytes are dominant in mountain beech forest on limestone and serpentinite, which is a common occurrence in temperate zone climatic conditions, with the fact that they are about 10% more numerous on limestone than on serpentinite. On the other hand, phanerophytes have significant presence on serpentinite, a lot larger than on limestone (37.5%: 27.3%). Considering that the analysis of the spectrum of life forms shows plant species relation to the climate of a region, we can conclude that this phenomenon is due to a warmer climate in southwestern Serbia. Geophytes also have strong presence both in the beech on the serpentinite and on limestone (20.41%: 19.47%), which was expected, considering very mesophilic nature of beech




forests. Other life forms have approximately equal presence on both the geological substrata.



**Figure 2:** The spectrum of life forms for the community of mountain beech on limestone and serpentinite

Analysis of geofloristic spectrum (Fig. 3) also shows significant differences between the studied stands. In both cases, the most common as collective is Central European floral element. A group of species with wide ecological amplitude (Eurasian areal type) is the second most common on limestone. The biggest differences are reflected in the presence of a collective group of sub-Mediterranean floral elements, which are connected with Balkan floral elements. This group is as twice as numerous in the mountain forest of beech on serpentinite than on limestone, which is another proof that this community is more xerophil on serpentinite than on limestone. Larger presence of group of subatlantic floral elements on limestone than on the serpentinite is indicator of more mesophilic conditions on limestone.



**Figure 3:** Spectrum of floral elements for the community of mountain beech on limestone and serpentinite

#### 4 CONCLUSIONS

The paper is based on a comparison of the floristic composition of mountain beech forests (*Asperula odoratae-Fagetum moesiaca* B. Jovanović 1973) on limestone and serpentinite, based on 14 phytosociological relevés made in the mountain beech forest on serpentinite on Crni Vrh, close to Priboj, and 10 relevés made on limestone in Ozren-Sokobanja. The analysis showed significant differences. Jaccard's index of similarity on studied stands is 0.24, indicating a very low floristic similarity in studied stands. CA floristic data analysis showed relevés grouping into two groups, depending on the geological substrate. Within phytocoenological relevés made on limestone bedrock,

typical species of beech forests appear as isolated: *Asperula odorata*, *Isopyrum th alictrioides*, *Cardamine bulbifera*, *Anemone ranunculoides*, *Salvia glutinosa* and others. On the other hand, within beech forests on serpentinite, in addition to "fagetal" type, xerophil species characteristic to the order *Erico - Pinetalia* and *Quercetalia pubescentis*: *Fraxinus ornus*, *Campanula patula*, *Polygonatum odoratum* are also visualized on the graphics.

#### 5 ACKNOWLEDGMENT

This work was carried out within the project "Sustainable management of the total resources of forests in the Republic of Serbia" -EVBR 37008, the Ministry of Education and Science of the Republic of Serbia.

#### 6 REFERENCES

- [1] S. Banković, Medarević, M., Pantić, D., N. Petrović, Nacionalna inventura šuma Republike Srbije-Šumski fond Republike Srbije, Ministarstvo poljoprivrede, šumarstva i vodoprivrede Republike Srbije-Uprava za šume, Beograd, (2009), pag. 244.
- [2] R. Cvjetičanin, Fitocenoze bukve u Srbiji, Šumarstvo 1-2. UŠITS. Beograd, (2003) pag. 107-112.
- [3] R. Cvjetičanin, M. Novaković, Fitocenoza pripadnost bukovih šuma u istraživanim sastojinama na Ozrenu-Sokobanja, Šumarstvo 3. UŠITS. Beograd, (2004) pag. 97-104.
- [4] M. Gajić, Pregled vrsta flore SR Srbije sa biljnogeografskim oznakama, Glasnik Šumarskog fakulteta, Serija A „Šumarstvo“, br. 54, Beograd, (1980).
- [5] V. Joksimović, M. Zdravković, Uticaj geološke podloge na bukove šume u Srbiji, u: Monografija Bukva u Srbiji, Udruženje šumarskih inženjera i tehničara Srbije, Beograd, (2005) pag. 83-93.
- [6] M. Knežević, O. Košanin. Zemljišta u bukovim šumama Srbije, u: Monografija Bukva u Srbiji, Udruženje šumarskih inženjera i tehničara Srbije, Beograd, (2005) pag. 94-107.
- [7] M. Kojić, R. Popović, B. Karadžić, Vaskularne biljke Srbije, Institut za istraživanja u poljoprivredi „Srbija“, Beograd, (1997).
- [8] M. Krstić, R. Cvjetičanin, Bioekološke karakteristike mezijske bukve (*Fagus moesiaca* /Domin, Maly/Czeczott.), u: Monografija Bukva u Srbiji, Udruženje šumarskih inženjera i tehničara Srbije, Beograd, (2005) pag. 118-124.
- [9] J. Lepš, P. Šmilauer, Multivariate analysis of ecological data, Faculty of biological sciences, University of south Bohemia, České Budějovice, (2002).
- [10] A. E. Magurran, Measuring biological diversity. Oxford: Blackwell Publishing, (2004)
- [11] V. Mišić, Red šuma bukve, Vegetacija Srbije II<sub>1</sub>, Srpska akademija nauka i umetnosti, Beograd, (1997) pag. 159-270.
- [12] M. Novaković, Šumska vegetacija Crnog vrha kod Priboja, magistarski rad-rukopis. Univerzitet u Beogradu- Šumarski fakultet, Beograd, (2008).
- [13] L.J. Rakonjac, M. Ratknić, M. Matović, V. Lavadinović, Fitocenoze karakteristike planinske šume bukve na Pešterskoj visoravni Ass. *Fagetum*

- moesiaca montanum* B.Jov.1953, Šumarstvo 4. UŠITS. Beograd, (2005) pag. 93-110.
- [14] H. Riter-Studnička, Biljni pokrov na serpentinama u Bosni, Godišnjak biološkog instituta u Sarajevu, (1963).
- [15] L. Tichý, JUICE, software for vegetation classification, J. Veg. Sci., Uppsala 13, (2002) pag. 451-453.
- [16] Z. Tomić, LJ. Rakonjac, Šumske fitocenoze Srbije, Institut za šumarstvo, Univerzitet Singidunum-Fakultet za primenjenu Ekologiju Futura, Beograd, (2013).

## FLORISTIC AND VEGETATION DIVERSITY IN KONGURA RESERVE (SOUTH-WEST BULGARIA)

PEDASHENKO H., VASSILEV K., BANCHEVA S., DELCHEVA M., VLADIMIROV V.

*Institute of Biodiversity and Ecosystem Research – Bulgarian Academy of Sciences, Sofia, Bulgaria*

*Corresponding author e-mail address: hristo\_pedashenko@yahoo.com*

**ABSTRACT:** Kongura reserve is situated in the Bulgarian part of Belasitsa mountain. It occupies an area of 1310.80 ha on slopes with northern, north-eastern and north-western exposition from 430 to 1740 m alt. The reserve was established in 1988 mainly to protect the well preserved *Castanea sativa* and *Fagus sylvatica* forests. A total of 300 vascular plant species were identified, more than a quarter of which belong to the families *Asteraceae*, *Rosaceae* and *Fabaceae*. Among the richest genera are *Galium*, *Campanula*, *Acer*, *Hieracium*, *Lathyrus*, *Trifolium* and *Viola*, which comprise more than 12% of the total flora of the study area. Five species are of conservation concern and are included in the *Red Data Book of the Republic of Bulgaria*. Three taxa of these are legally protected by the national Biodiversity Act. Endemism is relatively low (3.3%) since only 10 Balkan endemics have been recorded. Almost the entire territory of the reserve is covered by various forest types. The most widespread vegetation type belongs to *Quercus-Fagetum* class, which is represented by alliance *Fagion sylvaticae*. The rest of the forest vegetation was classified to alliances *Quercion petraeo-cerridis* and *Carpinion orientalis* of *Quercetum pubescentis* class. Wet woodlands of *Salici purpureae-Populetea nigrae* class (alliance *Platanion orientalis*) have only limited distribution in the area. Areas covered by shrubs were classified to *Loiseleurio-Vaccinietea* and *Rhamno-Prunetea* classes. Although areas dominated by herbs or grasses cover only less than two percent of the reserve's territory their diversity was classified into four classes: *Festuco-Brometea*, *Juncetea trifidi*, *Mulgedio-Aconitetea* and *Galio-Urticetea*.

**Keywords:** Belasitsa Mt., endemics, flora, habitats, conservation.

## 1 INTRODUCTION

Kongura reserve was declared as a protected area in 1988, covering the area of the former "Skoshnik" reserve, established in 1954 to preserve the most characteristic for Belasitsa mountain primary sweet chestnut forests. Apart from them slopes of the mountain are covered by variety of mesic and xeric woodlands, creating a rich diversity of habitats and niches. The top of the main ridge of Belasitsa mountain consists of vast mountain pastures.

Despite the high number of researches from Belasitsa Mt. there was no complete study of the reserve's flora and vegetation until now. Studies on the vegetation and habitat diversity of Belasitsa Mt. are clearly unequal. On one hand, there are a considerable number of scientific publications that deal with syntaxonomy [73, 53, 58], ecology [79, 49, 33, 70, 55, 60, 52], resources [81, 72, 71, 47, 45, 46] and ecosystem health of sweet chestnut forests [48, 64, 59], whereas on the other hand, shrub and grassland vegetation and resources were highly neglected [54, 27].

Floristic data were provided by [54, 61, 5, 41], as well as in the above mentioned vegetation studies. However, these publications do not deal with the area of Kongura reserve.

The aim of this study is to reveal the floristic and syntaxonomic diversity of Kongura reserve and to assess their current condition.

## 2 MATERIAL AND METHODS


### 2.1 Study area

The study was conducted in Kongura reserve, situated in the northeast part of Belasitsa Mountain. It covers 1310.8 ha in the valley of Petrichka river at altitude from 430 m to 1740 m a.s.l. (Fig. 1). The average annual air temperature is 9-11°C with average gradient of decline 0.26°C/100m. The highest average monthly temperature is in July (18-23°C) and the lowest in January (0-2°C) [6]. The monthly maximum precipitation rates are in November – December, and the minimum in July – August [21]. Annual precipitation rates are

between 600 and 1100 mm, depending on the altitude. The snow cover is shallow and short lasting (about 20–80 days/year) between November and April. The bedrock type is silicate [30]. Soils are Litosols up to 1000 m a.s.l. and Regosols above [67].

The potential natural vegetation is mostly comprised of south and east Balkan, as well as Crimean-west Caucasian colline Oriental hornbeam-downy oak forests (Mapping unit G57) and Species-rich, eutrophic and eumesotrophic beech and mixed beech forests with two subtypes 1. Colline-submontane, partly with hornbeam, sessile oak (Mapping unit F124) and 2. Montane-altimontane mostly with fir, partly with spruce and other mixed tree species (Mapping unit F154) [69].

The territory of the reserve is part of the Belasitsa Natural Park as well as Site of Community Importance (NATURA 2000 site BG0000167 Belasitsa). Recently, Belasitsa Mt. was declared as Important Plant Area [26].



**Figure 1:** Map of the study area

### 2.2 Sampling of the flora

Field studies were conducted in summer and autumn of 2014 using transect sampling method. For compilation of the list of vascular plant species in the reserve phytosociological relevés from the syntaxonomic studies

as well as some critical editions on the Bulgarian flora were used [18, 76, 77, 62, 19, 78]. The taxonomic scheme adopted follows mostly [13]. Biological types and life-forms are based on field observations and checked with relevant literature sources, e.g. [18, 77, 76, 62, 19, 50, 22]. Conservation significance of the species was determined according to [10] and [2]. Floristic elements follow [7].

### 2.3 Vegetation sampling

The field sampling was carried out during the period July – August 2014 following the Braun-Blanquet approach [35, 83]. We covered the whole range of different vegetation types as well as all grassland fragments on the territory of the reserve. We used 16 m<sup>2</sup>, 64 m<sup>2</sup> and 100 m<sup>2</sup> plots (n = 28) square-shaped and placed in stands that were visually homogeneous in terms of topography, vegetation structure and floristic composition. All relevés were stored in TURBOVEG database [63] and in the Balkan Vegetation Database [38].

### 2.4 Data analysis of relevés

Numerical classification was performed by PC-ORD [9] incorporated in JUICE 7.0 software package [40]. Relative Sørensen was used as distance measure and species similarity was calculated using flexible beta (-0.25).

### 2.5 Habitat classification and mapping

Habitat types were determined according to the Interpretation Manual of European Union Habitats [34, 74]. Assignment of each vegetation unit at association level to a certain Habitat Directive Code (HDC) was given in the text. Mapping was done using ArcGIS 10.0 software [23]. Spatial data was collected in the field using GPS devise Juno BS by Trimble and was later overlaid over the most recent orthophoto images available. Outlining the polygons was done manually by using features collected in the field as well as the orthophoto images. Mapping was done in scale 1:5000.

## 3 RESULTS

### 3.1 Flora

The flora of Kongura reserve comprises 64 families, 184 genera and 300 species. The taxonomic structure of the flora is presented in table I. *Polypodiophyta* comprises 7.81% of the families, 3.80% of the genera and 2.67% of the species from the total flora, *Pinophyta* – 3.13% of families, 1.09% of genera, 0.67 of species, and *Magnoliophyta* – 89.06%, 95.11% and 96.67% respectively. The list of taxa is provided in Annex I.

**Table I:** Taxonomic structure of the flora

Taxon	no. of species	no. of genera	no. of families
<i>Polypodiophyta</i>	8	7	5
<i>Pinophyta</i>	2	2	2
<i>Magnoliophyta</i>	290	175	57
<i>Magnoliopsida</i>	255	151	50
<i>Liliopsida</i>	35	24	7
<b>Total numbers</b>	<b>300</b>	<b>184</b>	<b>64</b>

The richest in species 12 families are: *Asteraceae* – 29 species (9.67% of the total flora), *Rosaceae* – 25

(8.33%), *Fabaceae* – 25 (8.33%), *Lamiaceae* – 23 (7.67%), *Poaceae* – 18 (6.00%), *Caryophyllaceae* – 15 (5.00%), *Scrophulariaceae* – 13 (4.33%), *Rubiaceae* – 9 (3.00%), *Ranunculaceae* – 8 (2.72%), and *Apiaceae*, *Boraginaceae*, *Liliaceae* – each with 6 (2.33%). Altogether these families comprise 57.33% of the species and 52.72% of the genera in the total flora. The largest genera are *Galium* – 7 species, *Campanula* – 6, *Acer*, *Hieracium*, *Lathyrus*, *Trifolium*, *Viola* – each with 5 species, *Genista*, *Ranunculus*, *Rosa*, *Rubus*, *Veronica* – each with 4 species.

The biological spectrum of the flora of Kongura reserve is presented in table II.

**Table II:** Biological spectrum of the flora

Biological type	no. of species	Share (%)
Perennials	198	66.00
Shrubs, incl. semishrubs and lianes	31	10.33
Trees	23	7.67
Biennials and Annuals to Biennials	19	6.33
Annuals	12	4.00
Annuals to perennials and biennials to perennial	11	3.67
Shrub/Tree	6	2.00
<b>total</b>	<b>300</b>	<b>100</b>

The life-form spectrum of the flora of Kongura reserve is presented in table III.

**Table III:** Life-form spectrum of the flora

Life form	no. of species	Share (%)
Hemicryptophytes	172	57.33
Phanerophytes	43	14.33
Geophytes	33	11.00
Chamaephytes	19	6.33
Therophytes	13	4.33
Therophytes/ Hemicryptophytes	9	3.00
Chamaephytes/ Phanerophytes	7	2.33
Geophytes/ Hemicryptophytes	3	1.00
Chamaephytes/ Hemicryptophytes	1	0.33
<b>total</b>	<b>300</b>	<b>100</b>

The phytogeographic composition of the flora of Kongura reserve is presented in Table IV.

**Table IV:** Floristic elements in the flora of Kongura reserve

Floristic element	no. of species	Share (%)
subMed	38	12.67
Eur-As	37	12.33
Eur-Sib	34	11.33
Eur-Med	31	10.33
Eur	30	10.00
Boreal	27	9.00
subBoreal	22	7.33
Bal	10	3.33
Kos	9	3.00
Med	9	3.00

Eur-subMed	8	2.67
Bal-Anat	5	1.67
Carp-Bal	5	1.67
Eur-OT	4	1.33
Pont-Med	4	1.33
Alp-Carp	3	1.00
Spont	3	1.00
Ap-Bal	2	0.67
Other – each with 1 species (Adv (SAm), Alp-Carp-Bal, Arct-Alp, Bal-Carp, Bal-Dac, Bul-Dac, Eur-Med-CAs, Eur- Pont, Eur-WAs, Med-CAs, Med-OT, Med-subMed, Pann- Bal, Pont-OT, Pont-Sib, Pont- subMed, SSib, subMed-As, subMed-CAs)	19	6.34
<b>total</b>	<b>300</b>	<b>100</b>

The endemic taxa in the flora are 10 – *Acer heldreichii*, *Armeria rumelica*, *Chamaecytisus absinthioides*, *Digitalis viridiflora*, *Festuca hirtovaginata*, *F. valida*, *Peucedanum olygophyllum*, *Scabiosa triniifolia*, *Scrophularia aestivalis* and *Viola speciosa*, all Balkan endemics [3].

The species of conservation concern are five. With a national IUCN category are *Viola speciosa* (CR), *Castanea sativa* (EN), *Ilex aquifolium* (EN), *Medicago carstiensis* (EN), *Acer heldreichii* (VU), all included in the *Red Data Book of the Republic of Bulgaria* [15, 14, 12, 11, 17, 16, 64, 65]. Legally protected by the national Biodiversity Act are three species – *Acer heldreichii*, *Ilex aquifolium* and *Viola speciosa*.

### 3.2 Vegetation

Classification of relevés showed a great diversity of communities on the territory of Kongura reserve. The list of established syntaxa consists of 10 classes, 10 orders, 12 alliances, 8 associations and 5 communities at association level. Beech forests are the most widespread vegetation type and cover more than 85% of the reserve's territory. Despite their prevalence they are relatively uniform and were classified into two associations of *Quercus-Fagetum* class (clusters 1-12 Fig. 2).

#### 3.2.1 Ass. *Asperulo odoratae-Fagetum sylvaticae* (HDC: 9130; 1101.0 ha)

This is the most widespread association on the territory of the reserve. Its communities occur at 700-1600 m a.s.l. on moderately steep slopes (20°-30°) with varying aspects. Soils are shallow to moderately deep regosols. The dominant species is *Fagus sylvatica* but at lower altitude other species like *Castanea sativa*, *Quercus dalechampii* and *Carpinus betulus* are found as subdominant with covers between 10% and 30%. The shrub layer is usually missing but undergrowth has cover between 0% and 40%, mainly by *Calamagrostis arundinacea*, *Luzula sylvatica*, *Melica uniflora*, *Euphorbia amygdaloides*, *Festuca heterophylla*, *Geranium robertianum*, *Lapsana communis*.

#### 3.2.2 Ass. *Aremonio agrimonoidis-Fagetum sylvaticae* (HDC: 91W0; 28.4 ha)

This vegetation type was found at 900-1000 m a.s.l. on south facing slopes with inclination of 30°-35°. *Fagus sylvatica* subsp. *moesiaca* is dominant species and

subdominants are *Castanea sativa* and *Quercus dalechampii*. Shrub layer is presented by young trees of above mentioned species as well as *Carpinus betulus*, *Tilia tomentosa*, *Fraxinus ornus*, *Acer platanoides* and *Crataegus monogyna*. Undergrowth is well developed mainly by *Dactylis glomerata*, *Melica uniflora*, *Poa nemoralis*, and *Festuca heterophylla*.

*Quercetea pubescentis* class covers only 11% of the study area but its communities were classified into three different alliances (clusters 13-17 and 24 Fig. 2).

#### 3.2.3 Ass. *Tilio tomentosae-Castanetum sativae* (HDC: 9260; 113.7 ha)

Sweet chestnut forests are distributed from 550-800 m a.s.l. on north facing slopes with inclination of 25°-35°. Soils are moderately deep litosols. *Castanea sativa* is dominating in the tree layer (50%-70%) together with *Fagus sylvatica* subsp. *moesiaca* (20%-40%). Participation of *Quercus dalechampii* and *Carpinus orientalis* is low (up to 10%). Accept for the species from the tree layer the shrub layer is also formed by *Chamaecytisus austriacus*, *Corylus avellana*, *Rubus caesius*, *Rosa* sp. *Brachypodium sylvaticum*, *Melica uniflora*, *Galium pseudoaristatum* and *Festuca heterophylla* are forming the undergrowth.

#### 3.2.4 *Quercus dalechampii* communities (HDC: 9170; 20.5 ha)

Communities of *Quercus dalechampii* have a limited distribution on the territory of the reserve. They cover areas between 750 and 800 m a.s.l. on north facing slopes with inclination between 25°-35°. Soils are moderately deep litosols. Balkan durmast oak is covering between 50%-60% whereas its subdominants (*Castanea sativa* and *Fagus sylvatica*) between 10%-20%. Single trees of *Carpinus orientalis* and *Ostrya carpinifolia* are also typical for the communities. The shrub layer is formed by *Corylus avellana*, *Chamaecytisus austriacus* and *Crataegus monogyna*. Undergrowth is formed mainly by *Poa nemoralis*, *Dactylis glomerata*, *Brachypodium sylvaticum*, *Galium pseudoaristatum*.

#### 3.2.5 *Ostrya carpinifolia* community (HDC: – ; 4.8 ha)

Hop hornbeam communities are locally distributed in the area at 500 m a.s.l. on northern slopes with prevailing inclination of 10°-35°. Soils are shallow, moderately dry litosols. The dominant species in the tree layer is *Ostrya carpinifolia*, but *Quercus dalechampii* is also forming substantial cover. Single trees of *Fagus sylvatica*, *Castanea sativa* and *Carpinus orientalis* are also common. Shrub layer is mainly formed by the same species like in the tree layer together with *Chamaecytisus austriacus*, *Corylus avellana*, *Crataegus monogyna* and *Rosa* sp. The composition of the undergrowth is uniform, with higher abundance of *Dactylis glomerata*, *Galium pseudoaristatum*, *Festuca heterophylla* and *Poa nemoralis*.

*Platanus orientalis* woods (cluster 25 Fig. 2) are only distributed along the Petrichka River at the lower part of the territory and were classified to *Salici-Populetea* class.

#### 3.2.6 Ass. *Petasito hybridi-Platanetum orientalis* (HDC: 92C0; 3.4 ha)

The association is locally distributed at 600-650 m a.s.l. on slightly inclined north facing slopes. Soils are rich, moderately deep, alluvial. *Platanus orientalis* is forming mono-dominant communities with low cover



(less than 10%) of other tree species like *Fagus sylvatica* and *Ostrya carpinifolia*. Shrub layer is formed by young individuals of the above mentioned trees as well as *Rubus caesius*, *Acer platanoides* and *Sambucus nigra*. Species with high cover in the undergrowth are *Dryopteris* spp., *Aegopodium podagraria*, *Geranium robertianum*, *Physospermum cornubiense* and *Lunaria rediviva*.

Shrublands and grasslands are covering very small areas (2% and 0.2% of the total area respectively) due to the intentional selection of the boundaries of the reserve to protect the well preserved *Castanea sativa* and *Fagus sylvatica* forests. Shrubby vegetation (clusters 18-20 Fig. 2) is classified to *Loiseleurio-Vaccinietea* and *Rhamno-Prunetea* classes.

### 3.2.7 Ass. *Festuco-Juniperetum sibiricae* (HDC: 4060; 24.4 ha)

The association is present at the beech forests belt, just above the timberline at 950-1150 m a.s.l. Soils are shallow to moderately deep, regosols. *Juniperus sibirica* is forming the highest cover but species like *Chamaecytisus absinthioides*, *Vaccinium myrtillus*, and *Rubus caesius* are also well presented (10%-35%). The herb layer is dominated by *Calamagrostis arundinacea*, *Festuca valida*, *Thymus jankae* and *Luzula luzuloides*.

### 3.2.8 *Rosa*, *Crataegus*, *Prunus* community (HDC: – ; 1.0 ha)

This community is very locally distributed at the beech forests belt in forest openings at the place of abandoned secondary grasslands at 1300-1700 m a.s.l. Soils are shallow to moderately deep, regosols. Dominant species are *Rosa* sp., *Crataegus monogyna* and *Prunus spinosa*. Other species with higher values are *Chamaecytisus absinthioides*, *Rubus caesius*, *Festuca hirtovaginata*, *Thymus jankae*, *Brachypodium pinnatum*, *B. sylvaticum*, *Dactylis glomerata*. At certain places stands are showing a transition to ass. *Festuco-Juniperetum sibiricae*.

Herb communities (clusters 21-23 and 26-27 Fig. 2) showed greater diversity than other vegetation types and were assigned to five phytosociological classes: *Trifolio-Geranietaea*, *Festuco-Brometea*, *Juncetea trifidi*, *Mulgedio-Aconitetea* and *Galio-Urticetea*.

### 3.2.9 Ass. *Festucetum validae* (HDC: 62D0; 0.9 ha)

Communities of this association are locally distributed in the sub-alpine and beech forests zones on western slopes with inclination 5°-15°. Soils are shallow to moderately deep, regosols. Species with high cover values are: *Festuca valida*, *Thymus jankae*, *Calamagrostis arundinacea*, *Juniperus sibirica*, *Vaccinium myrtillus*, *Luzula luzuloides*.

### 3.2.10 *Calamagrostis arundinacea* community (HDC: – ; 0.4 ha)

These communities are locally distributed in the sub-alpine zone of the reserve above the timber line on north and northwestern slopes with inclination of 10°-25°. *Calamagrostis arundinacea* is a dominant species and subdominants are *Rubus caesius*, *Epilobium angustifolium* and *Festuca valida*. Individuals of shrubs like *Fagus sylvatica*, *Juniperus sibirica* and *Vaccinium myrtillus* are also present in the community.

### 3.2.11 *Festuca hirtovaginata* community (HDC: 6210; 0.06 ha)

These communities are locally distributed in the beech forests zone on southern slopes with inclination 3°-8°. Soils are shallow, dry litosols. Communities have semi-open horizontal structure with no single species dominating. The group of species with high cover values contain *Festuca hirtovaginata*, *Carex caryophyllea*, *Phleum phleoides*, *Agrostis castellana*, *Thymus jankae*.

### 3.2.12 Ass. *Pteridietum aquiline* (HDC: – ; 0.8 ha)

The association is present at the beech forests belt on flat or slightly inclined terrains. Soils are regosols. Horizontal structure is closed with high cover of *Pteridium aquilinum* (90%-100%).

### 3.2.13 Ass. *Urtico-Sambucetum ebuli* (HDC: – ; 0.3 ha)

The association was found in a single locality in the beech zone at 1300 m a.s.l. on flat or slightly inclined terrains. Soils are moderately deep regosols. Communities have a specific species composition, rich in ruderal species. *Sambucus ebulus* is dominating with cover of 80%-90%.

Proposed syntaxonomical scheme:

**Cl.** *Salici purpureae-Populetea nigrae* Rivas-Martinez et Cantó ex Rivas-Martinez et al. 2001

**Ord.** *Populetalia albae* Br.-Bl. ex Tchou 1948

**All.** *Platanion orientalis* Kárpáti et Kárpáti 1961

**Ass.** *Petasito hybridi-Platanetum orientalis* Kárpáti et Kárpáti 1961

**Cl.** *Quercu-Fagetea* Braun-Blanq. et Vlieger in Vlieger 1937

**Ord.** *Fagetalia sylvaticae* Pawł. et al. 1928

**All.** *Fagion sylvaticae* Luquet 1926

**Ass.** *Asperulo odoratae-Fagetum sylvaticae* Sougnez et Thill 1959

**Ass.** *Aremonio agrimonoidis-Fagetum sylvaticae* Boşcaiu in Resmeriță 1972

**Cl.** *Quercetea pubescentis* (Oberd. 1948) Doing Kraft 1955

**Ord.** *Quercetalia pubescenti-petreae* Klika 1933

**All.** *Carpinion orientalis* Horvat 1958

*Ostrya carpinifolia* community

**All.** *Quercion confertae* Horvat 1954

**Ass.** *Tilio tomentosae-Castanetum sativae* Dafis 1973

**All.** *Quercion petraeo-cerridis* (Lakušić et Jovanović 1980) Čarni et al. 2009

*Quercus dalechampii* community

**Cl.** *Loiseleurio-Vaccinietea* Eggli 1952 ex Schub. 1960

**Ord.** *Rhododendro-Vaccinietalia* Braun-Blanq. in Braun-Blanq. et Jenny 1926

**All.** *Juniperion nanae* Br.-Bl. et al. 1939

**Ass.** *Festuco-Juniperetum sibiricae* Roussakova in Tzonev et al. 2009

**facies** of *Chamaecytisus absinthioides*

**Cl.** *Rhamno-Prunetea* Rivas Goday et Borja Carbonell, 1961

**Ord.** *Prunetalia spinosae* Tüxen 1952

**All.** *Berberidion vulgaris* Br.-Bl. 1950

*Rosa*, *Crataegus*, *Prunus* community

**Cl.** *Trifolio-Geranietaea sanguinei* T. Müller 1962

**Ord.** *Melampyro pratensis-Holcetalia mollis* Passarge 1979

**All.** *Holco mollis-Pteridion aquilini* Passarge (1994) 2002

**Ass.** *Pteridietum aquilini* Jouanne & Chouard 1929

**Cl.** *Festuco-Brometea* Braun-Blanq. et Tüxen ex Soó 1947

**Ord.** *Astragalo-Potentilletalia* Micevski 1971

**All.** *Armerio-Potentillion* Mic. 1978

*Festuca hirtovaginata* community

**Cl.** *Juncetea trifidi* Hadač in Klika et Hadač 1944

**Ord.** *Seslerietalia comosae* Simon 1958

**All.** *Poion violaceae* Horvat 1937

**Ass.** *Festucetum validae* Horvat et al. 1937

**Cl.** *Mulgedio-Aconitetea* Hadač et Klika in Klika et Hadač 1944

**Ord.** *Calamagrostietalia vilosae* Pawlowski et al. 1928

**All.** *Calamagrostion arundinaceae* Oberd. 1950


*Calamagrostis arundinacea* community

**Cl.** *Galio-Urticetea* Passarge ex Kopecký 1969

**Ord.** *Lamio albi-Chenopodietalia boni-henrici* Kopecký 1969

**All.** *Geo-Alliarion* Lohm et Oberd. in Görs et Müller 1969

**Ass.** *Urtico-Sambucetum ebuli* Braun-Blanq. et al. 1952



**Figure 2:** Dendrogram of phytosociological relevés (Relative Sorensen clustering method; flexible beta (-0.25) used for similarity). Legend: *Asperulo odoratae-Fagetum sylvaticae* (clusters 1-10), *Aremonio agrimonoidis-Fagetum sylvaticae* (clusters 11 and 12), *Tilio tomentosae-Castanetum sativae* (clusters 13-15), *Quercus dalechampii* community (clusters 16 and 17), *Festuco-Juniperetum sibiricae* (clusters 18-20), *Festucetum validae* (cluster 21), *Calamagrostis arundinacea* community (cluster 22), *Festuca hirtovaginata* community (cluster 23), *Ostrya carpinifolia* community (cluster 24), *Petasito hybridi-Platanetum orientalis* (cluster 25), *Urtico-Sambucetum ebuli* (cluster 26), *Pteridietum aquilini* (cluster 27).

## 4 DISCUSSION


### 4.1 Flora

Kongura reserve comprises 7.3% of the species in the Bulgarian flora. Vascular plant diversity is relatively low compared to some other areas with similar size, e.g. Vitanovo Reserve in Strandzha Mts [36] has smaller size but the number of vascular plant species is higher by ca. 54%. This can be explained by the fact that almost the entire territory of Kongura reserve is covered by beech and sweet chestnut forests, by the siliceous bedrock and by the lack of diversity of substrate and expositions. The share of *Polypodiophyta* in the flora of Kongura reserve – 2.67% of all species – is slightly higher than in some other studied sites, e.g. Vrashka Chuka protected site (0.8%, [82]), Mt Golo Bardo (0.6%, [51]) as well as for the whole country (1.1%, cf. [4]). This fact was noted first by [54] and can be explained with the presence of suitable habitats and environmental conditions, e.g. prevalence of forested habitats, northern expositions, relatively high soil and air humidity.

Regarding the presentation of the different families, the relatively high share of *Rosaceae* and relatively low share of *Asteraceae* and *Poaceae* in the flora of Kongura

reserve are notable if compared with the data for the total Bulgarian flora ([4]). Again, this can be explained with the availability of suitable habitats for *Rosaceae* (especially for shrubs and trees), and lack of such for *Asteraceae* and *Poaceae* (e.g. large open areas). For similar reasons the higher representation of the genera *Campanula*, *Galium* and *Viola* is obvious if compared with data for the whole country [4].

The analysis of the biological types and life forms in the flora of Kongura reserve shows an overall pattern characteristic for the temperate mountainous floras in the Balkans – strong prevalence of herbaceous perennial species, respectively hemicryptophytes [20]. However, it is noteworthy the relatively high representation of shrubs and trees, respectively phanerophytes, and the low presentation of annuals and therophytes which relates to the habitat availability. For example, the phanerophytes in the flora of Mt Golo Bardo are 10% (ca. two times lower than in Kongura) and therophytes are 18% (four times higher than in Kongura) [51].



**Figure 3:** Map of vegetation units on Kongura reserve. Codes in brackets represent the codes of habitats in the Habitat directive: 6210 – Semi-natural dry grasslands and scrubland facies on calcareous substrates (*Festuco-Brometalia*), 62D0 – Oro-Moesian acidophilous grasslands, 9170 – *Galio-Carpinetum* oak-hornbeam forests, 9260 – *Castanea sativa* woods, 9130 – *Asperulo-Fagetum* beech forests, 91W0 – Moesian beech forests, 92C0 – *Platanus orientalis* and *Liquidambar orientalis* woods (*Platanion orientalis*), 4060 – Alpine and Boreal heaths.

The phytogeographical analysis shows that in the flora of Kongura reserve prevail the following floristic elements – subMediterranean, Euro-Asiatic, Euro-Siberian, Euro-Mediterranean, European, Boreal and subBoreal, which altogether comprise nearly 73% of the total flora. This pattern is similar to the pattern in other studied areas, e.g. Mt Golo Bardo [51]. However, the representation of the Mediterranean and subMediterranean chorotypes is obviously lower (21.4% in Golo Bardo) and corresponds better to the share in the mountain regions – 11.9% [20]. This intermediate position of Kongura reserve can be explained with the higher elevation and position further to the south.

The endemism in the flora of Kongura reserve is low – only about 3.33% if compared with the data for the whole country – ca. 11% of the total Bulgarian flora [3]. This corresponds well with the low number of endemic species for the whole mountain and floristic region of Belasitsa as well as with the ecological structure of endemism in Bulgaria. As [80] state, with a few exceptions the Bulgarian endemics grow in open places, and are heliophytes and xerophytes, but this is true also for the Balkan endemics.

Although the number of the species of conservation concern is rather low, the reserve is of crucial importance for the preservation of these species. It harbours some of the best subpopulations at national level, e.g. of *Castanea sativa*, *Medicago carstiensis*, and *Viola speciosa*.

## 4.2 Vegetation

### 4.2.1 Syntaxonomy

This investigation is one of the few comprehensive studies about vegetation diversity in SW part of Bulgaria [25, 68]. In addition this is one of the few studies about fringe [29, 75, 42, 37], ruderal [31, 32, 39, 56, 42, 43, 37, 24] and shrubland vegetation belonging to *Rhamno-Prunetea* class [41] in the country. Due to the small size of the study area as well as the limited distribution of some vegetation types we were unable to classify them to already described associations, so they were classified as communities. Beech and sweet chestnut forests are among the best studied communities of all vegetation types present on the mountain [57, 58]. Further studies should be conducted for analyzing communities of *Festuca hirtovaginata*, *Calamagrostis arundinaceae* and shrublands dominated by *Rosa* sp., *Crataegus monogyna* and *Prunus spinosa*.

Despite the wide distribution of *Quercus dalechampii* forests in Bulgaria their syntaxonomical position according to Braun-Blanquet approach is still uncertain. According to the variety of described formations and associations following the dominant approach, its communities will be probably classified to more than one class. Problem for classification of Balkan durmast oak forests is also the fact that *Quercus dalechampii* is often recorded as *Quercus petraea* agg., including *Quercus petraeae*, *Q. dalechampii* and *Q. polycarpa*, [8] or even misidentified as any of the species of the group. Mesophytic woodlands of *Quercus dalechampii* with *Fagus sylvatica*, *Carpinus betulus*, *Poa nemoralis* and *Luzula luzuloides* might be separated as “mesophytic type” and should be classified to *Querco-Fagetea* class. This vegetation type is known from other parts of the country such as Balkan Range (unpublished data), Rila Mt, Pirin Mt, Sredna Gora Mts (unpublished data) and Western Frontier Mts [25]. On the other hand xerophytic communities of *Quercus dalechampii* are found together

with *Pinus nigra*, *Fraxinus ornus*, *Carpinus orientalis*, *Brachypodium pinnatum* and *Dactylis glomerata*. They should be assigned to *Carpinion orientalis* alliance of *Quercietea pubescenti* class. This “xerophytic type” is known up to now from Balkan Range (Chamdzha reserve, Vassilev & Gavrilova in preparation), Western Frontier Mts (Gabra reserve, [28]) *Quercus dalechampii* communities from Kongura reserve show an intermediate position determined by their mountainous distribution on one hand and the Mediterranean influence on the other. These ecological characteristics as well as the species composition convinced us to consider *Quercion petraeo-cerridis* alliance (*Quercietea pubescenti* class) as suitable for our communities. [1] report communities of the same alliance only 50 km west from the Kongura reserve, on the territory of Macedonia. Future analysis with supranational dataset from the areal of *Quercus dalechampii* in S and SE Europe will give the real syntaxonomical position of its communities.

### 4.2.2 Ecology

Agglomerative classification of the data resulted in sorting of relevés along the gradient of soil trophic status (Fig. 2), which appeared to be the most important environmental factor responsible for their separation. The oligotrophic beech forests of *Asperulo-Fagetum* were positioned at the left edge of the dendrogram whereas eutrophic ruderal communities of *Urtico-Sambucetum ebuli* appeared on the right-most part.

Vegetation belts in Kongura reserve as well as on the whole Belasitsa Mt. are well preserved like in most high Bulgarian mountains (Stara Planina, Rila, Pirin, the Rhodopes). Xero-mesophytic forests of *Quercetea pubescenti* class (*Castanea sativa*, *Quercus dalechampii* and *Ostrya carpinifolia* stands) are distributed at the bottom of the altitudinal gradient up to 800 m a.s.l. Higher parts of the slopes are covered by mesophytic beech forests belonging to *Querco-Fagetea* class, which form the timberline in the reserve at about 1500-1700 m a.s.l. Grasslands of *Juncetea trifidi* class are distributed above the beech woodlands up to the top of the mountain. They are very weakly presented in the reserve because beech and sweet chestnut forests were the target communities subject to protection. They usually grow in a complex with shrublands of *Loiseleurio-Vaccinietea* class outside the reserve's territory. The other herb communities do not form a separate belt but grow on forest openings in the forest zone. *Platanus orientalis* communities are also zonally distributed, intersecting the beech and sweet chestnut belts along the small streams.

Within the reserve's territory we encountered the tendency of shrub encroachment in grasslands. At higher altitude *Festuca valida* grasslands were overgrown by *Juniperus sibirica* whereas at lower places *Festuca hirtovaginata* communities were almost completely replaced by *Rosa* sp., *Crataegus monogyna* and *Prunus spinosa*.

### 4.2.3 Importance for conservation

Looking from conservational point of view, among all 13 communities at association level, there are 8 habitat types (Fig. 3) protected by Appendix 1 of the Bulgarian Biological Diversity Act (2007) and Directive 92/43/EEC. Most habitat types (6210, 62D0, 9170, 9130, 91W0 and 4060) are widespread on the territory of the country, whereas 92C0 *Platanus orientalis* and *Liquidambar orientalis* woods (*Platanion orientalis*) is

distributed only in Macedono-Rhodope mountains and neighboring lowlands but its largest areas are on the territory of Belasitsa Mt. Similar example is habitat type 9260 – *Castanea sativa* woods, 80% of the national cover of which is also on northern slopes of Belasitsa Mt.

## 5 ACKNOWLEDGEMENTS

Authors are grateful to the project “Activities on sustainable management of reserves Sokolata, Kongura, Ali Botush, Orelyak and maintained reserve the Dark forest”, financed under Priority Axis 3 “Preservation and restoration of biodiversity”, Operational Programme Environment, 2007 – 2013.

## 6 REFERENCES

- [1] A. Čarni, P. Košir, B. Karadžić, V. Matevski, S. Redžić, Ž. Škvorc, Thermophilous deciduous forests in Southeastern Europe. *Plant Biosystems* 143(1), (2009), pag. 1–13.
- [2] A. Petrova, V. Vladimirov, (eds), Red List of Bulgarian vascular plants. *Phytol. Balcan.*, 15(1), (2009), pag. 63-94.
- [3] A. Petrova, V. Vladimirov, Balkan endemics in the Bulgarian flora. *Phytol. Balcan.*, 16(2), (2010), pag. 293-311.
- [4] A. Petrova, V. Vladimirov, D. Dimitrova, D. Ivanova, Current state of the Bulgarian fern and seed plant biodiversity. – In: Petrova, A. (ed.), Current state of Bulgarian biodiversity – problems and perspectives, Bulgarian Bioplatform, (2005), Sofia, pag. 75-104.
- [5] A.I. Asenov, D.K. Pavlova, The high-altitude serpentine flora of Mt Belasitsa (Bulgaria). – *Phytol. Balcan.*, 15(2), (2009), pag. 191-198.
- [6] Atlas People's Republic of Bulgaria, General Department of Geodesy and Cartography, (1973), Sofia, pag. 168. [in Bulgarian].
- [7] B. Assyov, A. Petrova, (eds). Conspectus of the Bulgarian Vascular Flora. Distribution Maps and Floristic Elements. 4th revised and enlarged ed. Bulgarian Biodiversity Foundation, (2012) Sofia.
- [8] B. Jovanović, Dendrologija, Naučna knjiga, Beograd, (1991), pag. 231-233.
- [9] B. McCune, M. J. Mefford, PC-ORD. Multivariate analysis of ecological data. Version 4. MjM Software design, Gleneden Beach, (1999).
- [10] Bulgarian Biodiversity Act (Act on Amending and Supplementing), Decree no. 354 accepted by the 40th National Assembly on 01 November 2007, Durzhaven Vestnik, no. 94/16.11.2007,(2007), pag. 2-44 [in Bulgarian].
- [11] Ch. Gushev, *Ilex aquifolium*. – In: D. Peev, et al. (eds), Red Data Book of the Republic of Bulgaria. I Vol. Plants and fungi. BAS and MoEW, (2015) Sofia, pag. 527.
- [12] Ch. Gushev, *Ilex aquifolium*. In: A. Petrova, V. Vladimirov, (eds), Red List of Bulgarian vascular plants. *Phytol. Balcan.*, 15(1), (2009), pag. 77.
- [13] D. Delipavlov, I. Cheshmedzhiev, (eds), Key to the Plants of Bulgaria. Agrarian Univ. Acad. Press, (2011), Plovdiv [in Bulgarian].
- [14] D. Dimitrova, *Acer heldreichii* (pag. 657), *Castanea sativa* (pag. 434). – In: D. Peev, et al. (eds), Red Data Book of the Republic of Bulgaria. I Vol. Plants and fungi. BAS and MoEW, (2015), Sofia.
- [15] D. Dimitrova, *Acer heldreichii* (pag. 81), *Castanea sativa* (pag. 74). – In: A. Petrova, V. Vladimirov, (eds), Red List of Bulgarian vascular plants. *Phytol. Balcan.*, 15(1), (2009).
- [16] D. Ivanova, *Viola speciosa*. – In: D. Peev, et al. (eds), Red Data Book of the Republic of Bulgaria. I Vol. Plants and fungi. BAS and MoEW, (2015), Sofia, pag. 363.
- [17] D. Ivanova, *Viola speciosa*. – In: A. Petrova, V. Vladimirov, (eds), Red List of Bulgarian vascular plants, *Phytol. Balcan.*, 15(1), (2009), pag. 72.
- [18] D. Jordanov, (ed.), Flora Reipublicae Popularis Bulgaricae. I-VII Vols., Akad. Izd. Bulg. Acad. Sci., (1963–1979), Sofia [in Bulgarian].
- [19] D. Peev, (ed.), Flora Reipublicae Bulgaricae. XI Vol. Akad. Izd. “Prof. Marin Drinov”, (2012), Sofia [in Bulgarian].
- [20] D. Uzunov, Ch. Gushev, High mountain flora of Bulgaria – statistics, ecological characteristics and phytogeography. *Bocconeia*, 16(2), (2003), pag. 763-770.
- [21] E. Koleva, R. Peneva, Precipitations. – In: S. Velez, Climate of Bulgaria, Narodna prosveta, (1991), Sofia, pag. 179. [in Bulgarian].
- [22] E.J. Jäger, K. Werner, Exkursionsflora von Deutschland. Band 4, Gefäßpflanzen: Kritischer Band. 10 Aufl. Spektrum, Elsevier GmbH, (2005), München.
- [23] ESRI, ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute, (2011).
- [24] G. Georgiev, Syntaxonomical diversity of grassland vegetation of Haskovo municipality. Master Thesis, Plovdiv University "Paisii Hilendarski", Biological Faculty, Department of Ecology and Conservation, (2015).
- [25] G. Gogushev, Floristic classification of oak woodlands in the Western Frontier Mountains, Forestry ideas, 37, (2009), pag. 54-77.
- [26] G. Spiridonov, Ch. Gushev, B. Assyov, M. Gyosheva, Belasitsa. – In: D. Peev, A. Petrova, I. Apostolova, B. Assyov, (eds), Important Plant Areas in Bulgaria, Pensoft Publishers, (2012), Sofia, pag. 67-70.
- [27] I. Bondev, Vegetation of Bulgaria. Map in scale 1:600000 with explanatory text. St. Kliment Ohridski University Publishing house, (1991), Sofia, [in Bulgarian].
- [28] I. Bondev, I. Lazarov, Vegetation of Gabra reserve. – In: G. Tsankov, et al. (eds), Proceeding of a jubileum symposium (2-3 Jun 1994) 100 years from birthday of the Acad. Boris Stefanov (1894-1979), II Vol. (1995), Sofia, pag. 14-19.
- [29] I. Horvat, B. Pawłowski, J. Walas, Phytosociologische Studien über die Hochgebirgsvegetation der Rila Planina in Bulgarien, Bull. Int. Acad. Polon. Cl. Sci. Math. Sér. B 1, Bot., (1937), pag. 159-189.
- [30] I. Klimov, A. Marinova, R. Marinova, I. Petrov, P. Milovanov, E. Ilieva, Geological map of the Republic of Bulgaria, scale 1:50 000, Map sheet K-34-107-A (Belasitsa). Geocomplex Consortium Ltd., (2010). [in Bulgarian].
- [31] I. Kolev, Phytosociological features of the synanthropic plants in Bulgaria. Ruderals. – Nauchni Trudove Agron. Fak. Vissh Selskosp. Inst. “Georgi Dimitrov”, Ser. Rasteniev., 14, (1965), pag. 77-90. [in Bulgarian].



- [32] I. Kolev, Phytosociological Features of the Synanthropic Plants in Bulgaria. Weeds. Bulg. Acad. Sci. Press, (1976), Sofia. [in Bulgarian].
- [33] I. Velichkov, G. Hinkov, T. Zlatanov, M. Zlatanova, H. Hristova, Area Distribution Dynamics of *Castanea sativa* on the Northern Slopes of Bulgarian part of Belasitsa Mountain. *Silva Balcanica*, 11(1), (2010), pag. 21-26.
- [34] Interpretation Manual of European Union habitats, EUR27. Europea Commission, DG Environment, (2007), Brussels.
- [35] J. Braun-Blanquet, Plant Sociology: The study of Plant Communities, (1965), Hafner, London.
- [36] J. Koeva, (ed.), Floristic analysis of the Vitanovo reserve. Pontica Print, (1997), Burgas, [in Bulgarian].
- [37] K. Vassilev, H. Pedashenko, N. Vele, I. Apostolova, Grassland Vegetation of Ponor Special Protection Area (Natura 2000), Western Bulgaria. *Acta Zoologica Bulgarica*, suppl. 5, (2014), pag. 61-73.
- [38] K. Vassilev, H. Pedashenko, A. Alexandrova, A. Tashev, A. Ganeva, A. Gavrilova, A. Gradevska, A. Assenov, A. Vitkova, B. Grigorov, C. Gushev, E. Filipova, I. Aneva, I. Knollová, I. Nikolov, G. Georgiev, G. Gogushev, G. Tinchev, K. Pachedjieva, K. Koev, M. Lyubenova, M. Dimitrov, N. Apostolova-Stoyanova, N. Vele, P. Zhelev, P. Glogov, R. Natcheva, R. Tzonev, S. Boch, S. Hennekens, S. Georgiev, S. Stoyanov, T. Karakiev, V. Kalniková, V. Shivarov, V. Russakova, V. Vulchev, Balkan Vegetation Database: historical background, current status and future perspectives, *Phytocoenologia*, 46(1), (2016), pag. 89-95.
- [39] L. Mucina, J. Kolbek, Some anthropogenous vegetation types of South Bulgaria. – *Acta Bot. Croat.*, 48, (1989), pag. 83-102
- [40] L. Tichý, JUICE, software for vegetation classification. – *Journal of Vegetation Science*, 13, (2002), pag. 451-453.
- [41] L. Topalova-Zhezhiha, G. Gogushev, R. Ivanova, S. Kostadinova-Ilkova, The interesting plants of Belasitsa Mt. Bulgarian Biodiversity Foundation, (2010), Sofia, [in Bulgarian].
- [42] M. Dimitrov, D. Pavlov, P. Glogov, D. Yordanova, Study on changes of vegetation in territories with continued antropogenic influence in the Vitosha Natural Park. – *Nauka za Gorata*, 3, (2004a), pag. 57-75 [in Bulgarian].
- [43] M. Dimitrov, Floristic and phytosociological investigation of secondary herbal vegetation in coniferous forest ecosystems. – *Lesovudska Missul*, 1, (2004b), pag. 32-42 [in Bulgarian].
- [44] M. Dimitrov, Floristic classification of the vegetation in the Yundola Forestry Experimental Station. PhD. Thesis, Univ. of Forestry Press, (2004a), Sofia [in Bulgarian].
- [45] M. Ljubenova, M. Sokolovska, S. Bratanova-Doncheva, M. Radonova, Content of macro- and micro elements in sweet chestnut phyto mass in Belasitza mountains, Bulgaria. *Silva Balkanica*, 9(1), (2008), pag. 41-58.
- [46] M. Ljubenova, S. Bratanova-Doncheva, V. Dimitrova, M. Grozeva, Main characteristics of biological turn-over in *Castanes sativa* Mill. communities from Belasitza mountain, Bulgaria. – In: G. Bounous, (ed.), Proceedings of the first EU congress on Chestnut – *Castanea* 2009, *Acta Horticulture* No 866, (2010), pag. 275-285.
- [47] M. Lyubenova, N. Chavdarova, V. Dimitrova, S. Bratanova-Doncheva, Overground biomass production and dynamics of productivity of the tree layer in chestnut ecosystem (*Castanes sativa* Mill.) from Belasitza mountain, – In: Annual of University of Shoumen “Episkop K. Preslavski”, v. XIV B4 Nature science, University publisher, (2006), pag. 103-122.
- [48] M. Lyubenova, S. Bratanova-Doncheva, S. Mirchev, Structural-functional approach to risk assessment in the communities of sweet chestnut in Belasitza - applied methodological complex and initial results. – In: Compendium Chestnut forests - combining ecological functions and their use Regional meeting on the management of chestnut forests, Belasitza, (2002), pag. 27-37, [in Bulgarian].
- [49] M. Lyubenova, V. Dimitrova, S. Bratanova-Doncheva, Ecosystem assessment of sweet chestnut forests in Belasitsa. An-Di, (2001), Sofia, [in Bulgarian].
- [50] M.A. Fischer, K. Oswald, W. Adler, Exkursionsflora für Österreich, Liechtenstein und Südtirol. 3 Auflage, Biologiezentrum der Oberösterreichischen Landesmuseen, (2008), Linz.
- [51] N. Apostolova-Stoyanova, N. Stoyanov, Systematical and phytogeographical analysis of the flora of Mt Golo Bardo. *Phytologia Balcanica*, 15(3), (2009), pag. 401-430.
- [52] N. Chihev, V. Dimitrova, S. Bratanova-Doncheva, M. Ljubenova, The ecosystem approach to ecosystem management with an example from sweet chestnut (*Castanea sativa* Mill.) forest in Belasitsa Mountain, Bulgaria. – In: Proceeding 3th Congress of ecologists of the Republic of Macedonia with international participation, October 06-09.2007, Struga, Macedonia, (2008), pag. 69-74.
- [53] N. Ljubenova, V. Dimitrova, V. Vele, S. Bratanova-Doncheva, Phytocoenological investigation of chestnut (*Castanea sativa* Mill.) communities in the Belasitza region. – In Proceeding 2th Congress of ecologists of the Republic of Macedonia with international participation, October 25-29.2003, Ohrid, Macedonia, (2004), pag. 29-42
- [54] N. Stojanoff, Floristische Materialien von dem Belassiza-Gebirge. *God. Sofiisk. Univ. Fiz.-Mat. Fak.*, (1921), Sofia, pag. 15-16, [in Bulgarian].
- [55] P. Zhelyazkov, S. Timushiea, M. Teoharov, Ecological conditions for sweet chestnut in Bulgaria. *Gorsko stopanstvo* 7, (1982), pag. 26-30, [in Bulgarian].
- [56] R. Tzonev, Flora and vegetation of the Middle Danubian Plain between the valleys of Vit and Studena Rivers. PhD. Thesis. Biol. Fak., Sofia Univ. St. Kliment Ohridski, (2002), Sofia, [in Bulgarian].
- [57] R. Tzonev, M. Dimitrov, M. Chytrý, V. Roussakova, D. Dimova, Ch. Gushev, D. Pavlov, V. Vulchev, A. Vitkova, G. Gogushev, I. Nikolov, D. Borisova, A. Ganeva, Beech forest communities in Bulgaria. – *Phytocoenologia*, 36, (2006), pag. 247-279.
- [58] R. Tzonev, M. Lyubenova, G. Hinkov, T. Karakiev, Syntaxonomic investigation of Sweet chestnut

- (*Castanea sativa* Mill.) forests in Belasitsa mountain, Journal of Balkan Ecology 2, (2011), pag. 171-186.
- [59] S. Bratanova-Doncheva, M. Ljubenova, N. Ignatova, Is the chestnut (*Castanea sativa* Mill.) forest in Belasitsa healthy? – In: Proceeding “First national scientific conference of ecology – biodiversity, ecosystems, global change”. Sofia, (2005a), pag. 111-120.
- [60] S. Bratanova-Doncheva, M. Lyubenova, V. Dimitrova, S. Mihaylov, Ecological characteristics, Distribution and Management of *Castanea sativa* Mill. ecosystems in Bulgaria. III International Chestnut congress, Chaves, Portugal. Acta Horticulturae, special issue, (2005b), pag. 355-367.
- [61] S. Kostadinova, D. Dimitrov, New data on the vascular flora of Mt Belasitsa. – Phytol. Balcan., 8(3), (2002), pag. 293-306.
- [62] S. Kožuharov, (ed.), Flora Reipublicae Bulgaricae. X Vol. Akad. Izd. Bulg. Acad. Sci., (1995), Sofia [in Bulgarian].
- [63] S. M. Hennekens, J. H. J. Schaminée, TURBOVEG, a comprehensive data base management system for vegetation data. Journal of Vegetation Science, 12, (2001), pag. 589-591.
- [64] S. Mirchev, M. Lyubenova, S. Bratanova-Doncheva, Dendrochronological investigation on the impact of *Loranthus europeus* L. on growth and condition of the sweet chestnut in Belasitsa. – In: Workshop “Sustainable management of sweet chestnut forests”, 5-6.11.2003, Berkovitsa, (2003.). [in Bulgarian].
- [65] S. Stoyanov, *Medicago carstiensis*. – In: A. Petrova, V. Vladimirov, (ed.), Red List of Bulgarian vascular plants. Phytol. Balcan., 15(1), (2009), pag. 78.
- [66] S. Stoyanov, *Medicago carstiensis*. – In: D. Peev, et al. (eds.), Red Data Book of the Republic of Bulgaria. I Vol., Plants and fungi. BAS and MoEW, (2015), Sofia, pag. 546.
- [67] Soil archive of ISSAPP – N. Pushkarov, Soil survey in Stroumyani Municipality, (1996), [in Bulgarian].
- [68] T. Karakiev, R. Tzonev, *Saxifraga ferdinandicoburgi-Seslerietum actarovii* – a new association from the subalpine belt of the Slavianka (Orvilos) Mt. (Bulgaria). – Hacquetia, 10(1), (2011), pag. 49-71.
- [69] U. Bohn, G. Gollub, C. Hettwer, Y. Neuhäuslova, T. Raus, H. Schlüter, H. Weber, S. Hennekens, (eds.), Map of the natural vegetation of Europe. Scale 1:2 500 000. Interactive CD-ROM: explanatory text, legend, maps. – Bundesamt für Naturschutz, (2004), Bonn.
- [70] V. Dimitrova, Assessment of structure, functional status and degradation processes in ecosystems of sweet chestnut (*Castanea sativa*) in Belasitsa with emphasis of their sustainable use. Dissertation, (2008), Sofia, [in Bulgarian].
- [71] V. Dimitrova, M. Ljubenova, S. Bratanova-Doncheva, Aboveground phytomass and production in the shrub layer of the chestnut (*Castanea sativa* Mill.) communities in Belasitsa mountain. – In: B. Gruev, M. Nikolova, A. Donev, (eds.). Proceeding “Balkan scientific conference of biology”, May 19-20, 2005, Plovdiv, Bulgaria. University of Plovdiv “P. Hilendarski”, F. of biology, Union of scientists of Bulgaria, (2006), pag. 443-451.
- [72] V. Dimitrova, M. Ljubenova, S. Bratanova-Doncheva, Investigation of herb layer phytoproduction in chestnut (*Castanea sativa* Mill.) communities in the region of Belasitsa mountain. – In: Proceeding “First national scientific conference of ecology – biodiversity, ecosystems, global changes”, (2005), Sofia, pag. 121-131.
- [73] V. Dimitrova, N. Apostolova-Stoyanova, M. Ljubenova, N. Chipev, Syntaxonomic analysis of sweet chestnut forests growing on the north slopes of Belasitsa mountain. Comptes rendus de l’Academie Bulgare des Sciences 60 (5), (2007).
- [74] V. Kavrakova, D. Dimova, M. Dimitrov, R. Tzonev, T. Belev, K. Rakovska, (eds.), Manual for Determination of Habitats with European Importance in Bulgaria. Second edition. Sofia, (WWF–Danube Carpathian Programme & Green Balkan Federation), (2009), pag. 131. [in Bulgarian].
- [75] V. Roussakova, Végétation alpine et sous alpine supérieure de la montagne de Rila (Bulgarie). – Braun-Blanquetia, 25, (2000), pag. 3-132.
- [76] V. Velčev, (ed.), Flora Reipublicae Popularis Bulgaricae. IX Vol., Akad. Izd. Bulg. Acad. Sci., (1989), Sofia [in Bulgarian].
- [77] V. Velčev, (ed.), Flora Reipublicae Popularis Bulgaricae. VIII Vol., Akad. Izd. Bulg. Acad. Sci., (1982), Sofia [in Bulgarian].
- [78] V. Velchev, (ed.), Red Data Book of the People’s Republic of Bulgaria. I Vol. Plants. Akad. Izd. Bulg. Acad. Sci., (1984), Sofia [in Bulgarian].
- [79] V. Velchev, I. Bondev, Endangered and rare plant communities in Bulgaria, VIII-IX Vol., (1984), Sofia, [in Bulgarian].
- [80] V. Velchev, S. Kozuharov, Bulgarian endemic higher plants – gene pool, origin and distribution. – In: Petrova, A. (ed.), Atlas of Bulgarian endemic plants, Gea-Libris, (2006), Sofia, pag. 14-33.
- [81] V. Velev, S. Bratanova-Doncheva, R. Vatsava, Distribution of sweet chestnut (*Castanea sativa*) in Bulgaria and its conservation. Jubilee Conference 75 years of higher forestry education, (2000), pag. 85-90, [in Bulgarian].
- [82] V. Vladimirov, Contribution to the study of the vascular flora of Mt Vrashka Chuka. – In: Petrova, A. (ed.), Proc. VII Natl. Conf. Bot., 29-30.09.2011, Sofia. Bulg. Bot. Soc., (2012), Sofia, pag. 191-206.
- [83] V. Westhoff, E. van der Maarel, The Braun-Blanquet approach. – In: R. H. Whittaker, (ed.), Ordination and classification of plant communities, (1973), The Hague, pag. 617-737.

**Annex I:** List of taxa

**POLYPODIOPHYTA**

**Aspidaceae:** *Dryopteris filix-mas* (L.) Schott, *Gymnocarpium dryopteris* (L.) Newman; **Aspleniaceae:** *Asplenium adiantum-nigrum* L., *A. trichomanes* L.; **Athyriaceae:** *Athyrium filix-femina* (L.) Roth, *Cystopteris fragilis* (L.) Bernh.; **Hypolepidaceae:** *Pteridium aquilinum* (L.) Kuhn.; **Polypodiaceae:** *Polypodium vulgare* L.

**PINOPHYTA**

**Cupressaceae:** *Juniperus sibirica* Burgsd.; **Pinaceae:** *Pinus sylvestris* L.

**MAGNOLIOPHYTA**

**Magnoliopsida**

**Aceraceae:** *Acer campestre* L., *A. heldreichii* Orph.; *A. hyrcanum* Fisch. & C.A. Mey; *A. platanoides* L.; *A. pseudoplatanus* L.; **Apiaceae:** *Aegopodium podagraria* L., *Ferulago sylvatica* (Besser) Rchb., *Huetia cynapioides* (Guss.) P. W. Ball, *Peucedanum oligophyllum* (Griseb.) Vandas, *Physospermum cornubiense* (L.) DC., *Sanicula europaea* L., *Seseli libanotis* (L.) Koch; **Aquifoliaceae:** *Ilex aquifolium* L.; **Araliaceae:** *Hedera helix* L.; **Aristolochiaceae:** *Asarum europaeum* L.; **Asteraceae:** *Achillea grandifolia* Friv., *A. millefolium* L., *Anthemis tinctoria* L., *Arctium lappa* L., *Artemisia vulgaris* L., *Centaurea biebersteinii* DC., *C. stenolepis* A. Kern., *C. stoebe* L., *Cichorium intybus* L., *Cirsium candelabrum* Griseb., *C. ligulare* Boiss., *C. vulgare* (Savi) Ten., *Doronicum columnae* Ten., *Erigeron acris* L., *Galinsoga parviflora* Cav., *Hieracium hoppeanum* Schult., *H. murrorum* L., *H. pilosella* L., *H. racemosum* Waldst. & Kit., *H. sparsum* Friv., *Inula conyzia* DC., *Lapsana communis* L., *Leucanthemum vulgare* Lam., *Mycelis muralis* (L.) Dum., *Omalotheca sylvatica* (L.) Sch. Bip. & F.W. Schultz, *Senecio papposus* (Reichenb.) Less., *Tanacetum corymbosum* (L.) Schultz-Bip., *T. macrophyllum* (Waldst. & Kit.) Sch. Bip., *T. vulgare* L.; **Balsaminaceae:** *Impatiens noli-tangere* L.; **Betulaceae:** *Betula pendula* Roth, *Carpinus betulus* L., *C. orientalis* Mill., *Corylus avellana* L., *Ostrya carpinifolia* Scop.; **Boraginaceae:** *Buglossoides purpureo-caerulea* (L.) I.M. Johnst., *Echium vulgare* L., *Myosotis nemorosa* Besser, *Pulmonaria mollis* Hornem., *P. officinalis* L., *P. rubra* Schott, *Symphytum ottomanum* L.; **Brassicaceae:** *Alliaria petiolata* (M.Bieb.) Cavara & Grande, *Arabis glabra* (L.) Bernh., *A. sagittata* (Bertol.) DC., *A. turrita* L., *Cardamine bulbifera* (L.) Crantz, *Lunaria rediviva* L.; **Campanulaceae:** *Campanula glomerata* L., *C. patula* L., *C. persicifolia* L., *C. rapunculoides* L., *C. sparsa* Friv., *C. trachelium* L.; **Caprifoliaceae:** *Sambucus ebulus* L., *S. nigra* L.; **Caryophyllaceae:** *Cerastium alpinum* L., *C. dubium* (Bastard) Guépín, *Dianthus armeria* L., *D. petraeus* Waldst. & Kit., *Lychnis coronaria* (L.) Desr., *L. flos-cuculi* L., *Moehringia trinervia* (L.) Clairv., *Saponaria officinalis* L., *Silene italica* (L.) Pers., *S. latifolia* Poir., *S. vulgaris* (Moench) Garcke, *Stellaria graminea* L., *S. holostea* L., *S. media* (L.) Vill., *Viscaria vulgaris* Röhl.; **Celastraceae:** *Euonymus europaeus* L., *E. latifolius* (L.) Mill.; **Cornaceae:** *Cornus mas* L., *C. sanguinea* L.; **Crassulaceae:** *Sedum cepaea* L., *Umbilicus erectus* DC.; **Cuscutaceae:** *Cuscuta epithymum* (L.) L.; **Dioscoreaceae:** *Tamus communis* L.; **Dipsacaceae:** *Knautia drymeia* Heuff., *Scabiosa columbaria* L., *S. ochroleuca* L., *S. triniifolia* Friv.; **Ericaceae:** *Bruckenthalia spiculifolia* (Salisb.) Rchb., *Vaccinium myrtillus* L., *V. uliginosum* L., *V. vitis-idaea* L.; **Euphorbiaceae:** *Euphorbia amygdaloides* L., *E. polychroma* A. Kern., *Mercurialis perennis* L.; **Fabaceae:** *Astragalus glycyphyllos* L., *Chamaecytisus absinthioides* (Janka) Kuzm., *Ch. austriacus* (L.) Link., *Ch. hirsutus* (L.) Link., *Ch. supinus* (L.) Link., *Coronilla emerus* L. subsp. *emeroides* (Boiss. & Spruner) Holmboe, *C. varia* L., *Genista carinalis* Griseb., *G. depressa* M. Bieb., *G. lydia* Boiss., *G. ovata* Waldst. et Kit., *Lathyrus laxiflorus* (Desf.) Kuntze, *L. niger* (L.) Bernh., *L. pratensis* L., *L. venetus* (Mill.) Wohlf., *L. vernus* (L.) Bernh., *Lotus aegaeus* (Griseb.) Boiss., *L. corniculatus* L., *Medicago carstiensis* Wulf., *Trifolium alpestre* L., *T. arvense* L., *T. aureum* Poll., *T. pratense* L., *T. repens* L., *Vicia cracca* L.; **Fagaceae:** *Castanea sativa* Mill., *Fagus sylvatica* L., *Quercus dalechampii* Ten.; **Geraniaceae:** *Geranium robertianum* L., *G. sylvaticum* L.; **Hypericaceae:** *Hypericum maculatum* Crantz, *H. olympicum* L., *H. perforatum* L.; **Lamiaceae:** *Acinos alpinus* (L.) Moench, *A. arvensis* (Lam.) Dandy, *Ajuga genevensis* L., *Calamintha grandiflora* (L.) Moench, *Clinopodium vulgare* L., *Galeopsis tetrahit* L., *Glehoma hederacea* L., *Lamium galeobdolon* (L.) Crantz, *L. garganicum* L., *L. maculatum* L., *L. purpureum* L., *Melittis melissophyllum* L., *Mentha longifolia* (L.) Huds., *M. spicata* L., *Nepeta nuda* L., *Origanum vulgare* L., *Prunella vulgaris* L., *Salvia glutinosa* L., *Stachys alpina* L., *S. germanica* L., *S. sylvatica* L., *Teucrium chamaedrys* L., *Thymus jankae* Čelak.; **Malvaceae:** *Althaea officinalis* L., *Lavatera thuringiaca* L.; **Oleaceae:** *Fraxinus ornus* L.; **Onagraceae:** *Circaea luteciana* L., *Epilobium angustifolium* L., *E. montanum* L.; **Orobanchaceae:** *Orobanche gracilis* Sm., *O. reticulata* Wallr.; **Oxalidaceae:** *Oxalis acetosella* L.; **Papaveraceae:** *Chelidonium majus* L., *Corydalis solida* (L.) Clairville; **Plantaginaceae:** *Plantago major* L., *P. subulata* Roth; **Platanaceae:** *Platanus orientalis* L.; **Plumbaginaceae:** *Armeria rumelica* Boiss.; **Polygonaceae:** *Bilderdykia convolvulus* (L.) Dumort., *Rumex acetosa* L., *R. acetosella* L., *R. pulcher* L.; **Primulaceae:** *Cyclamen hederifolium* Aiton, *Lysimachia punctata* L., *Primula acaulis* (L.) L., *P. veris* L.; **Pyrolaceae:** *Orthilia secunda* (L.) House; **Ranunculaceae:** *Anemone ranunculoides* L., *Clematis vitalba* L., *Helleborus odoratus* Waldst. & Kit., *Isopyrum thalictroides* L., *Ranunculus acris* L., *R. ficaria* L., *R. nemorosus* DC., *R. repens* L.; **Rosaceae:** *Agrimonia eupatoria* L., *Aremonia agrimonoides* (L.) DC., *Cotoneaster nebrodensis* (Guss.) K. Koch, *Crataegus monogyna* Jacq., *Fragaria moschata* Weston, *F. vesca* L., *F. viridis* Duchesne, *Geum urbanum* L., *Malus sylvestris* Mill., *Potentilla inclinata* Vill., *P. micrantha* DC., *Prunus avium* (L.) L., *P. spinosa* L., *Rosa arvensis* Huds., *R. canina* L., *R. dumalis* Bechst., *R. pendulina* L., *Rubus caesius* L., *R. canescens* DC., *R. idaeus* L., *R. sanguineus* Friv.,

*Sanguisorba minor* Scop., *Sorbus aria* (L.) Crantz, *S. borbasii* Jav., *S. torminalis* (L.) Crantz; **Rubiaceae**: *Asperula aristata* L., *Cruciata glabra* (L.) Opiz, *Galium album* Mill., *G. anisophyllum* Vill., *G. aparine* L., *G. odoratum* (L.) Scop., *G. pseudaristatum* Schur, *G. schultesii* Vest, *G. verum* L.; **Salicaceae**: *Populus tremula* L., *Salix caprea* L.; **Santalaceae**: *Thesium linophyllum* L.; **Saxifragaceae**: *Saxifraga rotundifolia* L.; **Scrophulariaceae**: *Digitalis grandiflora* Mill., *D. viridiflora* Lindl., *Euphrasia liburnica* Wettst., *Lathraea squamaria* L., *Odontites serotina* (Lam.) Dum., *Scrophularia aestivalis* Griseb., *S. scopolii* Hoppe ex Pers., *Verbascum abietinum* Borbas, *V. longifolium* Ten. subsp. *pannosum* (Vis. & Pančić) Murb., *Veronica chamaedrys* L., *V. officinalis* L., *V. verna* L., *V. vindobonensis* (M. Fisch.) M. Fisch.; **Solanaceae**: *Atropabella-donna* L., *Solanum dulcamara* L.; **Tiliaceae**: *Tilia cordata* Mill., *T. tomentosa* Moench; **Urticaceae**: *Urtica dioica* L.; **Valerianaceae**: *Valeriana officinalis* L.; **Violaceae**: *Viola canina* L., *V. odorata* L., *V. riviniana* Rchb., *V. speciosa* Pant., *V. tricolor* L.

#### **Liliopsida**

**Araceae**: *Arum maculatum* L.; **Cyperaceae**: *Carex caryophyllea* Latour., *Carex hirta* L.; **Iridaceae**: *Crocus chrysanthus* (Herb.) Herb.; **Juncaceae**: *Luzula campestris* (L.) DC., *L. luzuloides* (Lam.) Dandy, *L. sylvatica* (Huds.) Gaudin; **Liliaceae**: *Lilium martagon* L., *Ornithogalum nutans* L., *Paris quadrifolia* L., *Polygonatum latifolium* (Jacq.) Desf., *P. odoratum* (Mill.) Druce, *Scilla bifolia* L., *Veratrum lobelianum* Bernh.; **Orchidaceae**: *Cephalanthera longifolia* (L.) Fritsch, *Epipactis helleborine* (L.) Crantz, *Neottia nidus-avis* (L.) Rich.; **Poaceae**: *Agrostis capillaris* L., *A. castellana* Boiss. & Reut., *Alopecurus pratensis* L., *Brachypodium sylvaticum* (Huds.) Beauv., *B. pinnatum* (L.) P. Beauv., *Calamagrostis arundinacea* (L.) Roth, *Dactylis glomerata* L., *Deschampsia flexuosa* (L.) Trin., *Festuca heterophylla* Lam., *F. hirtovaginata* (Acht.) Markgr.-Dann., *F. rubra* L., *F. (Uechtr.) Péntzes*, *Melica uniflora* Retz., *Milium effusum* L., *Phleum phleoides* (L.) H. Karst., *Ph. pratense* L., *Poa annua* L., *P. nemoralis* L.

**JUNCUS FILIFORMIS L. (JUNCACEAE), A NEW SPECIES IN THE FLORA OF MACEDONIA**<sup>1</sup>TEOFILOVSKI A., <sup>2</sup>NIKOLOV Z., <sup>1</sup>MANDŽUKOVSKI D.<sup>1</sup>Public enterprise Makedonski sumi, Skopje, Macedonia<sup>2</sup>Macedonian museum of natural history, Skopje, Macedonia

Corresponding author e-mail address: acoteofilovski@hotmail.com

**ABSTRACT:** In this work *Juncus filiformis* is reported for the first time in the flora of Macedonia. The species was discovered in the subalpine belt of Mt Šar Planina - on Bogovinje Lake coast. Chorological and ecological data regarding the discovered population with additions of a photograph of live specimens, a photograph of fruiting inflorescence of herbarium specimens, and a distribution map are presented. A short discussion of taxonomical position of the species and short description are also given.

**Keywords:** *Juncus filiformis*, new, species, flora, Macedonia.

**1 INTRODUCTION**

The genus *Juncus* L. comprises c. 300 species which are widely distributed in both hemispheres, but most abundantly in the temperate and cold regions. A monographic study of this genus in the flora of Macedonia so far has been not carried out but according to data incorporated in the check lists of some particular regions and in the other floristical works the number of species could be estimated to c. 20 native and one naturalized species. They inhabit mainly moist and wet habitats in mountain areas. Recently, new chorological data regarding several species of this genus in Macedonia have been disclosed (Teofilovski 2011), while *J. minutulus* (Albert & Jahand.) Prain has been reported for the first time in the flora of Macedonia (Matevski & Teofilovski 2011).

During the recent floristic researches on Mt Šar Planina (NW Macedonia), at the Bogovinje Lake coast we have discovered a small population of one representative of the genus *Juncus*, which has been identified as *J. filiformis* L., so far not known from the territory of the Republic of Macedonia.

**2 MATERIALS AND METHODS**

During the field work herbarium material were collected and deposited in the private herbarium of the first author. The determination was performed according to the treatments of this genus in the several standard floras (Snogerup, 1980; Kirschner et al., 2002; Romero Zacro, 2010). The relevant floristic works were checked for possible presence of chorological data regarding the territory of Republic of Macedonia.

**3 RESULTS AND DISCUSSION****3.1 Distribution in Macedonia**

*Juncus filiformis* L. - Mt Šar Planina: on the Bogovinje Lake coast, 1950 m, 09.08.2015, coll. A. Teofilovski, Z. Nikolov & D. Mandžukovski (herb. Teofilovski; herb. Mac. Mus. Nat. Hist., n. 12572). (Fig. 1, 2, 3).

This is the first record of this species for the flora of Macedonia. Only a few patches of individuals was observed in southeast part of the Bogovinje Lake coast, growing in a shallow water and on wet places of the coastline, on siliceous geological substrate. The species is possibly more widespread around this lake, but unfortunately only southeast part of its coast was checked and thus no additional field data were provided.


**3.2 General and regional distribution**

The general range of this boreo-montane species includes the temperate areas of the north hemisphere. In the north parts of the range it occurs from near sea level to low alpine zone, while in the south mostly in high mountains up to 2200 m or more, growing on various at least temporarily wet habitats, usually on acid soils (Kirschner et al., 2002).

In the Balkan Peninsula, *J. filiformis* is a rare species, which is so far known from several localities in Croatia, Bosnia and Hercegovina, Montenegro, Serbia, Bulgaria, and Greece, but it is not yet reported from Kosovo, Albania, and European part of Turkey.

The closest finding sites are located c. 150 km northeast (Vlasina, SE Serbia) (Nikolić, 1976), c. 150 km southeast (NC Greece) (Dimopoulos & al., 2013), c. 180 km northwest (Nikšić - Budoške Bare, W Monte Negro) (Babunja, 2013), and c. 210 km east (Mt Rila and Mt Pirin, W Bulgaria) (Georgiev et Kožuharov, 1964).

Nikolić (1976) reported *J. filiformis* also for Pčinja valley - Rt (S Serbia), apparently by mistake, based on herbarium specimens labeled with "Macedonia, Pčinja - Rt" (coll. P. Černjavski, 26.5.1926, n. 28645). According to the investigation of Niketić (2011) "Rt" is actually misplaced name used by P. Černjavski for locality Banjski Rid, situated above Katlanovska Banja in the Macedonian part of Pčinja valley. We checked a scan of the cited herbarium specimens but they seem undeterminable since their inflorescences are almost fully undeveloped.



**Figure 1:** Distribution of *Juncus filiformis* in Macedonia





**Figure 2:** *Juncus filiformis* (Photo A. Teofilovski)

### 3.3 Description and taxonomy

*Juncus filiformis* L. Sp. Pl.: 326. 1753

Rhizome with short internodes, horizontal. Stems 10-60 cm, the uppermost basal sheath with a short lamina. Non-flowering shoots usually few or absent. Lowest bract ( $\frac{1}{2}$ )1-1 $\frac{1}{2}$  times as long as stem; second bract sometimes leaf-like, though short. Inflorescence with 4-10 flowers, dense. Perianth-segments 2.5-3.5 mm, ovate to narrowly ovate or the inner oblong; outer aristate; inner obtuse. Anthers c. 0.5 mm,  $\frac{1}{3}$ - $\frac{1}{2}$  as long as filaments. Capsule (2.5-)3-3.5(-4.5) mm, trigonous ovoid, usually shortly mucronate, light brown, about equaling perianth. Seeds 0.5 mm, obliquely ovoid, faintly reticulate, with one inconspicuous appendage (Snogerup, 1980).

*J. filiformis* is taxonomically well defined species belonging to the *J.* subgen. *Agathryon* sect. *Juncotypus* Dumort. This section has a world-wide distribution comprising 67 perennial species which are characterised with pseudo lateral inflorescence, lower inflorescence bract terete, erect, seeming to be a continuation of stem, basal leaves bladeless, cauline leaves  $\pm$  absent, sterile shoots terete, stem-like (Kirschner et al., 2002). Another three species of this section also occur in Macedonia (*J. inflexus* L., *J. effusus* L., and *J. conglomeratus* L.).

### 4 CONCLUSIONS

*J. filiformis* is recorded for the first time from the territory of the Republic of Macedonia. A small population of this species has been discovered in the subalpine belt of Mt Šar Planina – on the Bogovinje Lake

coast, on 1950 m a.s.l. It grows in shallow water and on wet places of the lake coastline, on siliceous geological substrate. The nearest known finding sites are located c. 150 km northeast (Vlasina, SE Serbia) and c. 150 km southeast (NC Greece).

### 5 ACKNOWLEDGEMENTS

The authors are grateful to the reviewers M. Kostadinovski (Skopje) and M. Niketić (Beograd) for useful suggestions which improved the manuscript, and to M. Niketić also for sending us a scan of the cited herbarium specimens of *Juncus* sp.



**Figure 3:** *Juncus filiformis* - inflorescences (herbarium specimens)

## 6 REFERENCES

- [1] Babunja, N., The flora of wetlands and aquatic habitats in the vicinity of Nikšić, *Natura Montenegrina, Nat. Hist. Mus. Monten., Podgorica*, (2013), 12(1): 13-41.
- [2] Dimopoulos, P., Raus, Th., Bergmeier, E., Constantinidis, Th., Iatrou, G., Kokkini, S., Strid, A., Tzanodacis, D. Vascular flora of Greece: an annotated checklist, *Englera*, (2013), 31: 1- 372.
- [3] Georgiev, T., Kožuharov, St., *Juncus* L. In: Jordanov, D. (ed.): *Flora Republicae popularis Bulgaricae*, (1964), 2: 149-167. (In Bulgarian)
- [4] Kirschner, J. et al., *Juncaceae* 3: *Juncus* subg. *Agathryon*, *Species Plantarum: Flora of the World*, (2002), 8: 1-192.
- [5] Matevski, V., Teofilovski, A., New species in the Flora of the Republic of Macedonia, *Biol. Macedonica*, (2011), 62: 49-54.
- [6] Niketić, M., Problematika učitavanja, transfera i standardizacije podataka sa herbarskih etiketa na primeru tribusa *Sileneae* (*Caryophyllaceae*) u zbirci Prirodnjačkog muzeja u Beogradu. – Prirodnjački muzej Beograd, *Habilitacioni rad.*, (2011), p. 81.
- [7] Nikolić, V., *Juncaceae* A. L. Juss. In: Josifović, M. (ed.): *Flora de la Republique socialiste de Serbie*. ASSA, Beograd, (1976), 8: 116-151. (In Serbian)
- [8] Romero Zacro, C., *Juncus* L. In: Talavera et al. (eds.): *Flora Iberica*, Real Jardín Botánico, Madrid, (2010), 17: 123-187.
- [9] Snogerup, S., *Juncus* L. In Tutuin et al. (eds): *Flora Europaea*, Cambridge, (1980), 5: 102-111.
- [10] Teofilovski, A., Contributions to the flora of the Republic of Macedonia, Skopje, (2011), p. 142. (In Macedonian, with English summary).

## FLORA, HABITATS AND VEGETATION OF CHAMDZHA MANAGED RESERVE, CENTRAL BALKAN RANGE

<sup>1</sup>VASSILEV K., <sup>2</sup>GAVRILOVA A.

<sup>1</sup>*Institute of Biodiversity and Ecosystem Research – Bulgarian Academy of Sciences, Sofia, Bulgaria*

<sup>2</sup>*Department of Dendrology, Faculty of Forestry, University of Forestry – Sofia., Bulgaria*

Corresponding author e-mail address: [kiril5914@abv.bg](mailto:kiril5914@abv.bg)

**ABSTRACT:** Chamdzha reserve is a managed reserve, situated on the southern slopes of Central Balkan Range, which occupies an area of 66.4 ha. The aim of this study was to explore floristic, habitat and vegetation diversity of the reserve. Totally the flora is represented by 51 families, 184 genera and 299 vascular plants (without mosses). The richest families are *Asteraceae*, *Poaceae*, *Fabaceae*, *Rosaceae*, *Caryophyllaceae*, *Brassicaceae*, *Boraginaceae* and *Apiaceae*. One hundred and eleven medical plants and 7 species of conservation concern were found (5 Balkan endemics, 3 included in the Red List of Bulgarian vascular plants and in the National Biological Biodiversity Act). Fourteen relevés were collected during 2014 following to the Braun-Blanquet approach. The vegetation diversity is represented by 5 classes (*Erico-Pinetea*, *Quercus-Fagetum*, *Quercetum pubescentis*, *Koelerio-Corynephorum* and *Festuco-Brometum*), 4 alliances (*Erico-Fraxinion orni*, *Carpinion orientalis*, *Carpinion betuli*, and *Festucion valesiacae*), 1 association and 5 communities. Woodland vegetation covers 61.2 ha whereas grassland vegetation includes only 0.15 ha. Natural communities represent 5 habitats protected by Directive 92/43/EEC and the Bulgarian Biodiversity Act.

**Keywords:** managed reserve, Bulgaria, Stara planina, *Erico-Pinetea*, *Koelerio-Corynephorum*

### 1 INTRODUCTION

Chamdzhia managed reserve has a territory of 66.4 ha. It was established to protect a rare natural formation of black pine (*Pinus nigra* Arnold) in the Balkan Range (Stara planina), situated on its southern slopes. The category “managed reserve” according to the Bulgarian legislation is closest to the definition of the IUCN category IV - Habitat/Species Management Area. Various natural as well as semi-natural and anthropogenic phytocenoses surround the reserve.

The floristic complex of Chamdzha reserve was poorly studied in the past. Some old data about the diversity of vascular plants in the region including the territory of the managed reserve are represented in the studies of [24] and [35]. In addition, some general information about the floristic composition of the natural black pine forests can be found in [24] and [48]. Vegetation diversity in the reserve has not been studied before.

The aim of this study is to reveal the floristic and syntaxonomic diversity of Chamdzha managed reserve and to assess their current condition.

### 2 MATERIAL AND METHODS


#### 2.1 Study area

The study was conducted in Chamdzha managed reserve, situated in the southern slope of central part of the Balkan Range (Stara planina), near Hristo Danovo village. It covers 66.4 ha at altitude from 550 to 775 m a.s.l. (Fig. 1). Prevailing slope incline is between 15 and 35°. The climate is temperate to continental, characterized by warm summer and cold winter [40]. The precipitation maximum is in May-June and minimum in January and February. The bedrock type is granite and soils are Luvisols and Vertisols [11].

The potential natural vegetation is mostly comprised of south and east Balkan, as well as Crimean-west Caucasian colline Oriental hornbeam-downy oak forests (Mapping unit G34) [43].

The territory of the reserve is also part of NATURA 2000 site BG000494 Centralen Balkan – buffer.

(<http://natura2000.moew.government.bg/Home/ProtectedSite?code=BG0001493&siteType=HabitatDirective>).



**Figure 1:** Map of the study area

#### 2.2 Sampling of the flora

The field studies of the flora were conducted in June-August 2014. The transect sampling method was applied. The taxonomic scheme adopted generally follows [9]. Biological types and life-forms are based on field observations and checked with relevant literature sources, e.g. [12, 13, 14, 38, 48, 51]. Conservation status of the species was determined according to [1, 2, 3, 7, 32]. Floristic elements follow [4]. The group of medicinal plants was determined according [8] and Regulation №RD-83/3.02.2014 of the MOEW (Ministry of Environment and Water) regarding the special regimes of protection and management of medicinal plants in 2014 on the territory of the country.

#### 2.3 Vegetation sampling

During the vegetation season 2014 a total of 14 relevés were collected following the Braun-Blanquet approach [26, 54]. The sample plots were set in the most homogenous part of communities in a manner to cover all vegetation types in the study area. We used square-shaped form of sample plots with plot size of 16 m<sup>2</sup> for



grassy and 100 m<sup>2</sup> for woody vegetation. All relevés were stored in TURBOVEG database [39] and in the Balkan Vegetation Database (GIVD ID: EU-00-019) [29] and the Balkan Dry Grassland Database (GIVD ID: EU-00-013) [27]

Altitude and location were measured by Garmin eTrex Vista (GPS). The exposition was determined by compass. Slope was assessed visually in categories (1) < 5 °, (2) between 5-10 ° and (3) > 10 °. Soil are classified as shallow (<10 cm depth), (2) moderately deep (10-20 cm) or deep (> 20 cm).

#### 2.4 Data analysis of relevés

The classification was made using the program PC-ORD [6] incorporated in JUICE 7.0 software package [30]. Relative Sorensen was used for calculating of species similarity and flexible beta (-0.25) as distance measure.

#### 2.5 Habitat classification and mapping

Habitat types were determined according to Habitat Directive [25, 44]. Assignment of each vegetation unit to a certain Habitat Directive Code (HDC) was given in the text. Mapping was done using ArcGIS 10.0 software [16]. Spatial data was collected in the field using Garmin eTrex Vista (GPS) and was later overlaid over the most recent orthophoto images available. Outlining the polygons was done manually by using data collected in the field as well as the orthophoto images. Mapping was done in scale 1:5000.

### 3 RESULTS

#### 3.1 Flora

The flora of Chamdzha managed reserve comprises of 51 families, 187 genera and 299 species, which sums up 32.1% of the family diversity in the country, 20.3 % of the genera and 7.6 % of the species. Most of the inventoried species are spermatophytes – 46 families (90.2%), 179 genera (95.7 %) and 291 species (97.3%). On the territory of Chamdzha managed reserve no representatives of *Lycopodiophyta* or *Equisetophyta* were found. The taxonomic structure of the flora is presented in Tabl.1. *Polypodiophyta* comprises 5.9% of the total number of families, 2.7% of the genera and 2.3% of the species in the reserve. On the other hand *Pinophyta* comprises of 3.9% of all families, 1.6% of the genera and 1.3% of the species and *Magnoliophyta* presents 90.2%, 95.7% and 96.3% families, genera and species respectively. The list of taxa is provided in Annex I.


**Table I:** Taxonomic structure of the flora

Taxon	no. of species	no. of genera	no. of families
<i>Polypodiophyta</i>	7	5	3
<i>Pinophyta</i>	4	3	2
<i>Magnoliophyta</i>	288	179	46
<i>Magnoliopsida</i>	239	152	42
<i>Liliopsida</i>	49	27	4
<b>Total number</b>	<b>299</b>	<b>187</b>	<b>51</b>

The richest in species families are *Asteraceae* – 42 species (14%), *Poaceae* – 38 species (12.7%), *Fabaceae* – 35 species (11.7%), *Lamiaceae* and *Rosaceae* – 19 species each (6.4%), *Caryophyllaceae* and *Brassicaceae*

– 10 species each (3.3%), *Boraginaceae* and *Apiaceae* – 8 species each (2.7%).

The life forms of the flora of Chamdzha managed reserve is presented in Fig. 2.



**Figure 2:** Life forms spectrum of the flora of the Chamdzha managed reserve

The phytogeographic composition of the flora of Chamdzha managed reserve is presented in Table 3.

**Table III:** Floristic elements in the flora of Chamdzha managed reserve

Floristic element	no. of species	Share (%)
subMed	45	15.1
Eur-Med	41	13.7
Eur-As	37	12.4
Eur	21	7
Boreal	19	6.4
Eur-Sib	18	6
SBoreal	17	5.7
Kos	16	5.4
Pont-Med	10	3.3
subBoreal	7	2.3
Bal	4	1.3
Med	5	1.7
SPont	5	1.7
Others – Eur-OT, Med-As, Ap-Bal, Pan-Bal, Pont-OT, Med-CAs, subBal, Bal-Anat, Eur-Med-CAs, CSEur, Bal-Dac, Carp-Bal, Pont-subMed, Alp-Med, Eur-CAs, Alp-Carp, Pont, Med-OT, SSib, subMed-CAs, subMed-As, Pont-CAs	54	18
<b>Total number</b>	<b>299</b>	<b>100</b>

There are five Balkan endemics [3] – *Anthemis macedonica*, *Scabiosa triniifolia*, *Chamaecytisus jankae*, *Hypericum umbellatum*, *Campanula lanata* found in the reserve. Three species are included in the Bulgarian Red List of vascular plants [2] – *Campanula lanata* (VU), *Jovibarba heuffelii* (NT) and *Minuartia saxifraga* (LC). One species (*Campanula lanata*) is enlisted in App. 3 of [7] and [32] assigned as “endangered”.

#### 3.2 Vegetation

Classification of relevés did not show a great diversity of communities on the territory of Chamdzha managed reserve. The list of established syntaxa consists of 5 classes, 5 orders, 4 alliances, 1 association and 5

communities (clusters 1-12 Fig. 3, Annex II). *Pinus nigra* forests are the most widespread vegetation type and cover more than 63% of the reserve's territory. They are following by *Quercus dalechampii* woodlands covering 31%.

### 3.2.1 Ass. *Cetrario aculeatae-Plantagininetum subulatae* (HDC: 8230; 0.09 ha)

This association is locally distributed on the territory of the reserve. Its communities occur on eroded, slightly to moderately inclined slopes (10°-30°) with varying expositions. Soils are shallow and rocky. Stands have open horizontal structure and low amount of accumulated litter. Bedrock type is granite. The most frequent species are *Scleranthus perennis*, *Sedum hispanicum*, *Agrostis capillaris*, *Hieracium pilosella*, *Cladonia foliacea*, *Cetraria aculeata*, *Polytrichum piliferum*, *Racomitrium canescens* agg. This association is found at a small scale within stands of woodland vegetation types. The total cover of vegetation is predominantly between 65-80% as cover of herb and cryptogam layers are almost equal (between 30 and 50%).

### 3.2.2 *Pinus nigra* community (HDC: 9530; 41.78 ha)

This vegetation type has widest distribution in the reserve and was found on slopes with varying exposition and inclination between 5 and 40°. Soils are shallow to moderately deep. *Pinus nigra* (cover 50-100%) is the dominant species in the tree layer and subdominants are *Quercus dalechampii*, *Carpinus orientalis* and *Quercus cerris* with cover between 10-30%. Shrub layer has cover between 10-25% and is formed by shrubs of the same species as well as *Coryllus avellana*, *Chamaecytisus jankae*, *C. supinus*, *Crataegus monogyna*, *Prunus spinosa* are found. Herb layer has low cover (10-30%) and is formed mainly by *Poa nemoralis*, *Dactylis glomerata*, *Brachypodium sylvaticum*, *Moehringia pendula*, *Lerchenfeldia flexuosa* and *Galium flavescens*.

### 3.2.3 *Quercus dalechampii* community (HDC: 9170; 7.67 ha)

Communities of *Quercus dalechampii* are found in northern part of the reserve on slopes with north and north-east exposition. Soils are moderately deep. Balkan durmast oak is the dominant species in the tree layer with cover between 60 and 80%, whereas in shrub layer dominant species is *Carpinus betulus* with cover between 20-50%. Single shrubs of *Pinus nigra*, *Quercus cerris*, *Carpinus orientalis*, *Crataegus monogyna*, *Prunus spinosa*, *Chamaecytisus jankae* are forming the shrub layer. Herb layer is formed mainly by *Poa nemoralis*, *Lerchenfeldia flexuosa*, *Festuca heterophylla*, *Galium flavescens*, *Moehringia pendula*.

### 3.2.4 *Carpinus orientalis-Pinus nigra* community (HDC: 9530)

Communities of *Carpinus orientalis-Pinus nigra* (cluster 10, Fig. 2) have limited distribution on the territory of the reserve. They form transitional vegetation types with *Pinus nigra* community. *Carpinus orientalis-Pinus nigra* communities are found on slightly to moderately steep slopes (10-25°) with southern exposition. Soils are shallow to moderately deep. There is not clear dominant species, but subdominants are *Carpinus orientalis* (30-60%) and *Pinus nigra* (20-60%).

Shrub layer has low cover (10-15%) and is formed by the same species as well as some single shrubs (*Fraxinus ornus*, *Chamaecytisus calcareous*, *Crataegus monogyna*). Herb layer has low cover, which varies between 5-15%. Undergrowth is formed mainly by *Melica uniflora*, *Dactylis glomerata*, *Brachypodium sylvaticum*.

### 3.2.5 *Carpinus orientalis - Quercus dalechampii* community (HDC: 91M0; 12.68 ha)

This vegetation type is locally found in the reserve on eastern and southern slopes with prevailing inclination 10-15°. The dominant species are *Quercus dalechampii* and *Carpinus orientalis*. Shrub layer is formed by young individuals of above mentioned trees as well as *Crataegus monogyna*, *Prunus spinosa*, *Fraxinus ornus*, *Acer campestre* and *Quercus cerris*. Species with higher cover (5-10%) in herb layer are *Poa nemoralis*, *Lerchenfeldia flexuosa*, *Festuca heterophylla*, *Arabis procurrens*, *A. sagittata*, *Galium flavescens* and *Moehringia pendula*.

### 3.2.6 *Festuca valesiaca* community (HDC: 6210; 0.06 ha)

This community is locally distributed on slightly inclined, south facing slopes, close to abandoned agricultural areas. Soils are rich, moderately deep. This community has close horizontal structure with total cover of vegetation 90-95%. Dominant species are *Festuca valesiaca*, *Dichanthium ischaemum* and *Koeleria nitidula*. As a result of abandonment of pastures during last 20 years cover of shrubs (*Prunus spinosa*, *Crataegus monogyna*) was increasing and reached 20-25%.

### Proposed syntaxonomical scheme:

**Cl. *Erico-Pinetea*** Horvat 1959

**Ord. *Erico-Pinetalia*** Horvat 1959

**All. *Erico-Fraxionion orni*** Horvat 1959  
*Pinus nigra* community

**Cl. *Quercetea pubescentis*** (Oberd. 1948) Doing Kraft 1955

**Ord. *Quercetalia pubescenti-petreae*** Klika 1933

**All. *Carpinion orientalis*** Horvat 1958

*Carpinus orientalis-Pinus nigra* community  
*Carpinus orientalis-Quercus dalechampii* community

**Cl. *Querco-Fagetea*** Braun-Blanquet et Vlieger in Vlieger 1937

**Ord. *Fagetalia sylvaticae*** Pawł. et al. 1928

**All. *Carpinion betuli*** Issler 1931

*Quercus dalechampii* community

**Cl. *Koelerio-Corynephoretea*** Klika in Klika et Novák 1941

**Ord. *Sedo-Scleranthetalia*** Br.-Bl. 1955

**Ass. *Cetrario aculeatae-Plantagininetum subulatae***  
Pedashenko et al. 2013


**Cl. *Festuco-Brometea*** Br.-Bl. & Tüxen 1943 ex Soó 1947

**Ord. *Festucetalia valesiaca*** Br.-Bl. & Tüxen 1943


**All. *Festucion valesiaca*** Klika 1931

*Festuca valesiaca* community 1952





**Figure 3:** Dendrogram of phytosociological relevés (Relative Sorensen used for similarity and flexible beta (-0.25) as clustering method). Legend: *Cetrario aculeatae-Plantaginietum subulatae* (clusters 1-2), *Pinus nigra* community (clusters 3-7), *Quercus dalechampii* community (clusters 8-9), *Carpinus orientalis-Pinus nigra* community (cluster 10), *Carpinus orientalis- Quercus dalechampii* community (clusters 11-12).



**Figure 4:** Map of vegetation units of Chamdzha managed reserve. Codes in brackets represent the codes of habitats in the Habitat directive: 8230 – Siliceous rock with pioneer vegetation of the *Sedo-Scleranthion* or of the *Sedo albi-Veronicion dillenii*, 6210 – Semi-natural dry grasslands and scrubland facies on calcareous substrates (*Festuco-Brometalia*), 9170 – *Galio-Carpinetum* oak-hornbeam forests, 91M0 – Pannonian-Balkan turkey oak-sessile oak forests, 9530 – \*(Sub-) Mediterranean pine forests with endemic black pines

## 4 DISCUSSION

### 4.1 Flora

Chamdzha managed reserve comprises 7.6% of the species in the Bulgarian flora. Vascular plant diversity is relatively low compared to some other areas characterized with similar vegetation diversity, e.g. Chervenata Stena Reserve (631 species) in Rhodope Mts [34] and Ali Botush Reserve in Mt. Slavyanka (more than 600 species, unpublished data). This can be explained by the fact that Chamdzha reserve has smaller size and bedrock type is granite but not limestone like in other two reserves. It is well known that calcareous substrates are characterized by rich flora and a lot of endemic and rare plants [15,49]. On the other hand silicate substrates are characterized by poorer flora and less number of endemic and rare plants [15]. According to results about taxonomic structure of flora of the reserve the presence of *Pteridiophyta* is 2.3 % of all species, which is higher from the average for the country [4]. This can be explained by the prevalence of forest and rocky habitats, where environmental conditions are suitable for distribution of such species.

Regarding the presentation of the different families, share of *Rosaceae* and *Caryophyllaceae* is relatively higher compare with the data for the total Bulgarian flora [4]. This can be explained with presence of many woody species of *Rosaceae* and availability of suitable habitats for many species of *Caryophyllaceae*.

The analysis of the biological types in the flora of Chamdzha managed reserve shows the prevalence of hemicriptophytes, which is typical for regions with continental climate in Europe and the Balkans. Also the presence of shrubs and trees, respectively phanerophytes is high because the domination of woody habitats. Therophytes are represented by 18.4%, which is high taking into account the size of the territory covered by grassland vegetation.

The phytogeographical analysis shows that in the flora of Chamdzha managed reserve prevail the following floristic elements – subMediterranean, Euro-Mediterranean, Euro-Asiatic, European, Boreal, subBoreal and Euro-Siberian, which comprise nearly 66.4% of the total flora. This pattern is similar to the pattern in other area in the country [18, 20, 28, 34]. The high number of the subMediterranean, Euro-Mediterranean and Euro-Asiatic groups (41.2 %) is a result of wide distribution of xerothermic vegetation types in the reserve.

The presence of endemic species is low- only 1.7%, comparing with data for the whole country – ca. 11% of the total Bulgarian flora [4]. Endemic species and species of conservation concern represent 2.4% of all species. Some species (*Campanula lanata*, *Jovibarba heuffelii* and *Minuartia saxifraga*) are restricted to rocky cliffs locally found in the reserve. On the other hand *Chamaecytisus jankae* is widespread in shrub layer of woody vegetation types.

### 4.2 Vegetation

#### 4.2.1 Syntaxonomy

This investigation is the first study about vegetation diversity in Chamdzha managed reserve. Due to the small size of the study area as well as the limited distribution of some vegetation types we were unable to classify them to already described associations, so majority of them were

classified as communities. Black pine and Balkan durmast oak forests are the most widely distributed vegetation types.

*Pinus nigra* woodlands are distributed and studied mainly in southern part of the country - Rhodope mts [10, 21, 34, 42], West Frontier Mts [20], Pirin mt [45], Mt Slavyanka (unpublished data) and more limited in northern Bulgaria - Vrachanska mt [47]. Similar results were found about the national distribution of habitat 9530 \*(Sub-) Mediterranean pine forests with endemic black pines in NATURA 2000 network in Bulgaria [[http://natura2000.moew.government.bg/Home/Protected Site?](http://natura2000.moew.government.bg/Home/ProtectedSite?)]. It is located in 24 NATURA 2000 sites with national coverage of 25045 ha.

From syntaxonomical point of view black pine forest are still poorly studied in the country. All studies have been done following the Dominance approach and provide only descriptive information about communities [10, 20, 21, 22, 34, 41, 42, 45]. This is the first research of syntaxonomy of *Pinus nigra* communities in the country following the Braun-Blanquet approach. Future analysis with other unclassified relevés from other parts of the country will reveal the existing diversity of this vegetation in Bulgaria. However, based on data collected up to now, we found that there are 2 types of *Pinus nigra* communities in Bulgaria based on different bedrock type – “species rich communities” found on calcareous substrates and “species poor communities” distributed on silicate substrates.

*Quercus dalechampii* forests on the territory of the reserve are belonging to xero-mesophytic type analyzed by [19] from Kongura reserve.

On the other hand there are some transitional vegetation types between *Pinus nigra*, *Quercus dalechampii* and *Carpinus orientalis* phytocoenosis in the reserve. This is also found in West Frontier Mts [17] and Rhodope Mts [22]. *Pinus nigra* forms communities also with *Fagus sylvatica* [10, 41], *Picea abies* [10, 34], *Pinus sylvestris* [34], *P. heldreichii* [45], *Abies alba* [10], *Ostrya carpinifolia* [41], *Juniperus communis* & *Cotinus coggygrya* [21], *Genista carinalis* [22], *Quercus pubescens* [42] and *Festuca balcanica* [47].

Grassland vegetation has limited distribution on the territory of the reserve. *Cetrario aculeatae-Plantaginetum subulatae* association was originally described by [18] from the area of Beklemeto mountain pass, which is only several kilometers away from the managed reserve. It has wider distribution on south slopes of the Balkan Range (Stara planina) on silicate terrains. Future analysis of collected data in Balkan Dry Grassland Database [27] will reveal its actually existing distribution.

#### 4.2.2 Ecology

Agglomerative classification of the data resulted in sorting of relevés along the gradient of soil depth (Fig. 2), which appeared to be the most important environmental factor responsible for their separation. On shallow soils, are found *Cetrario aculeatae-Plantaginetum subulatae* association and *Pinus nigra* communities. On the other vegetation types are found on deeper soils.

Due to the small difference of altitudinal range (225 m.) all vegetation types are found in the belt of Xero-mesophytic oak forests. As a result ecological conditions are similar and there is no much difference in species

composition of shrub and herb layer in different woody vegetation types.

#### 4.2.3 Importance for conservation

Looking from conservational point of view, among all 6 communities at association level, there are 5 habitat types (Fig. 3) protected by Appendix 1 of the Bulgarian Biological Diversity Act (2007) and Directive 92/43/EEC. All habitat types (9530, 9570, 91M0, 8230 and 6210) are widespread on the territory of the country. However, territories covered by habitat 8230 Siliceous rock with pioneer vegetation of the *Sedo-Scleranthion* or of *Sedo albi-Veronicion dillenii* in the country are limited to very small areas. On the territory of the reserve habitat 9530 \*(Sub-) Mediterranean pine forests with endemic black pines represents relict locality of this vegetation. All habitats types are well preserved and subjected on low anthropogenic pressure.

#### 5 ACKNOWLEDGEMENTS

Authors are grateful to the project “DIR – 5113325-12-109 Central Balkan – park for everyone”, financed under Priority Axis 3 “Preservation and restoration of biodiversity”, Operational Programme Environment, 2007 – 2013 and Bulgarian Government.

#### 6 REFERENCES

- [1] A. Petrova, (ed.), Atlas of Bulgarian endemic plants. Gea-Libris Publishing House, (2006), Sofia.
- [2] A. Petrova, V. Vladimirov, (eds.), Red List of Bulgarian vascular plants. Phytol. Balcan., 15(1), (2009), pag. 63-94.
- [3] A. Petrova, V. Vladimirov, Balkan endemics in the Bulgarian flora. Phytol. Balcan., 16(2), (2010), pag. 293-311.
- [4] A. Petrova, V. Vladimirov, D. Dimitrova, D. Ivanova, Current state of the Bulgarian fern and seed plant biodiversity. – In: Petrova, A. (ed.), Current state of Bulgarian biodiversity – problems and perspectives, Bulgarian Bioplatform, (2005), Sofia, pag. 75-104.
- [5] B. Assyov, A. Petrova, (eds.), Conspectus of the Bulgarian Vascular Flora. Distribution Maps and Floristic Elements. 4th revised and enlarged ed. Bulgarian Biodiversity Foundation, (2012), Sofia.
- [6] B. McCune, M. J. Mefford, PC-ORD. Multivariate analysis of ecological data. Version 4. MjM Software design, Gleneden Beach, (1999).
- [7] Bulgarian Biodiversity Act (Act on Amending and Supplementing), Decree no. 354 accepted by the 40th National Assembly on 01 November 2007, Durzhaven Vestnik, no. 94/16.11.2007, (2007), pag. 2-44, [in Bulgarian].
- [8] Bulgarian Medicinal plants Act (Act on Amending and Supplementing), accepted by the 38th National Assembly, Durzhaven Vestnik, 29/07.04.2000, (2000), pag. 9-21, [in Bulgarian].
- [9] D. Delipavlov, I. Cheshmedzhiev, (eds.), Key to the Plants of Bulgaria. Agrarian Univ. Acad. Press, (2011), Plovdiv, [in Bulgarian].
- [10] D. Gorunova, H. Kochev, Vegetation cover of Chepelarska river basin (Western Rhodope) II. Fitologija, 41, (1991), pag.3-27, [in Bulgarian].
- [11] D. Zlatinova, Abiotic characteristics. In: R.Rashid (ed.), Management plan of Chamdzhia managed reserve, (2015), pag. 24-32, [in Bulgarian].
- [12] D. Jordanov, (ed.), Flora Reipublicae Popularis Bulgariae. I-VII Vols., Akad. Izd. Bulg. Acad. Sci., (1963–1979), Sofia, [in Bulgarian].
- [13] D. Mueller-Dombois, H. Ellenberg, Aims and Methods of Vegetation Ecology, (1974), John Wiley and Sons, New York.
- [14] D. Peev, (ed.), Flora Reipublicae Bulgariae. XI Vol. Akad. Izd. “Prof. Marin Drinov”, (2012), Sofia, [in Bulgarian].
- [15] D. Uzunov, Ch. Gushev, High mountain flora of Bulgaria – statistics, ecological characteristics and phytogeography. Boccone, 16(2), (2003), pag. 763-770.
- [16] ESRI, ArcGIS Desktop: Release 10. Redlands, CA: Environmental Systems Research Institute, (2011).
- [17] G. Gogushev, Floristic classification of oak woodlands in the Western Frontier Mountains, Forestry ideas, 37, (2009), pag. 54-77, [in Bulgarian].
- [18] H. Pedashenko, I. Apostolova, S. Boch, A. Ganeva, M. Janišova, D. Sopotlieva, S. Todorova, A. Ūnal, K. Vassilev, N. Velev, J. Dengler, Dry grasslands of NW Bulgarian mountains: first insights into diversity, ecology and syntaxonomy. Tuexenia, 33, (2013), pag. 309-346.
- [19] H. Pedashenko, K. Vassilev, S. Bancheva, M. Delcheva, V. Vladimirov, Floristic and vegetation diversity in Kongura reserve (South-west Bulgaria), Šumarski pregled, (submitted).
- [20] I. Bondev, I. Lazarov, Vegetation of Gabra reserve. In: G. Tsankov, et al. (eds), Proceeding of a jubileum symposium (2-3 Jun 1994) 100 years from birthday of the Acad. Boris Stefanov (1894-1979), II Vol. (1995), Sofia, pag. 14-19, [in Bulgarian].
- [21] I. Bondev, M. Lyubenova, The vegetaion communities in the reserve “Borovo” in east Rhodopes mountains. In: S. Nedyalkov, K. Baeva, B. Nikolov, R. Radev, T. Michev, Y. Danchev, V. Velev (eds), International symposium “Conservation of Natural areas and of the genetic material they contain” – Project 8 on the Programme “Man and the Biosphere” (MAB) of Unesco, 23-28.09.1985, Blagoevgrad, Bulgaria, Collection of reports, vol. 2, (1985a), pag.142-147, [in Bulgarian].
- [22] I. Bondev, M. Lyubenova, The vegetation cover of the reserve “Chamlaka” in the east part of Rodopa mountain. In: S. Nedyalkov, K. Baeva, B. Nikolov, R. Radev, T. Michev, Y. Danchev, V. Velev (eds), International symposium “Conservation of Natural areas and of the genetic material they contain” – Project 8 on the Programme “Man and the Biosphere” (MAB) of Unesco, 23-28.09.1985, Blagoevgrad, Bulgaria, Collection of reports, vol. 3, (1985b), pag.71-79, [in Bulgarian].
- [23] I. Bondev, Vegetation of Bulgaria. Map in scale 1:600000 with explanatory text. St. Kliment Ohridski University Publishing house, (1991), Sofia, [in Bulgarian].
- [24] I. Urumov, Flora of Karlovo municipality. Sbornik BAN, 25, (1929), pag. 6-132.
- [25] Interpretation Manual of European Union habitats, EUR27. Europea Commission, DG Environment, (2007), Brussels.

- [26] J. Braun-Blanquet, Plant Sociology: The study of Plant Communities, (1965), Hafner, London.
- [27] K. Vassilev, Z. Dajič, R. Cušterevska, E. Bergmeier, I. Apostolova, Balkan Dry Grasslands Database. – In: J. Dengler, J. Oldeland, F. Jansen, M. Chytrý, J. Ewald, M. Finckh, F. Glöckler, G. Lopez-Gonzalez, R. K. Peet, J.H.J. Schaminée, (Eds.), Vegetation databases for the 21st century, Biodiversity & Ecology, 4, (2012), pag. 330–330.
- [28] K. Vassilev, H. Pedashenko, N. Veleve, I. Apostolova, Grassland Vegetation of Ponor Special Protection Area (Natura 2000), Western Bulgaria. Acta Zoologica Bulgarica, suppl. 5, (2014), pag. 61–73.
- [29] K. Vassilev, H. Pedashenko, A. Alexandrova, A. Tashev, A. Ganeva, A. Gavrilova, A. Gradevska, A. Assenov, A. Vitkova, B. Grigorov, C. Gushev, E. Filipova, I. Aneva, I. Knollová, I. Nikolov, G. Georgiev, G. Gogushev, G. Tinchev, K. Pachedjieva, K. Koev, M. Lyubanova, M. Dimitrov, N. Apostolova-Stoyanova, N. Veleve, P. Zhelev, P. Glogov, R. Natcheva, R. Tzonev, S. Boch, S. Hennekens, S. Georgiev, S. Stoyanov, T. Karakiev, V. Kalnikova, V. Shivarov, V. Russakova, V. Vulchev, Balkan Vegetation Database: historical background, current status and future perspectives, Phytocoenologia, (2016), [in press].
- [30] L. Tichý, JUICE, software for vegetation classification. Journal of Vegetation Science, 13, (2002), pag. 451–453.
- [31] M. Chytrý, Z. Otýpková, Plot sized used for phytosociological sampling of European vegetation. Journal of Vegetation Science, 14, (2003), pag. 563–570.
- [32] M. Ančev, V. Goranova, Campanula lanata. In: D. Peev, et al. (eds.), Red Data Book of the Republic of Bulgaria. I Vol. Plants and fungi. BAS and MoEW, (2015), Sofia, pag. 665.
- [33] N. Apostolova-Stoyanova, N. Stoyanov, Systematical and phytogeographical analysis of the flora of Mt Golo Bardo. Phytol. Balcan, 15(3), (2009), pag. 401–430.
- [34] N. Andreev, V. Nikolov, Endemic and relict taxa and vegetational communities in the biosphere reservation “Červenata stena”. In: S. Nedyalkov, K. Baeva, B. Nikolov, R. Radev, T. Michev, Y. Danchev, V. Veleve (eds), International symposium “Conservation of Natural areas and of the genetic material they contain” – Project 8 on the Programme “Man and the Biosphere” (MAB) of Unesco, 23–28.09.1985, Blagoevgrad, Bulgaria, Collection of reports, vol. 1, (1985), pag. 88–96, [in Bulgarian].
- [35] S. Baev, Botanical excursions, (1947), BAN Publishing House, Sofia. [in Bulgarian].
- [36] P. Vassilev, Botanical reserve “Ostritsa” in Mt Golo Brdo near Pernik. Regional Symposium under project 8-MAB “Conservation of natural areas and of the genetic material they contain”, 20–24 October 1980, Blagoevgrad, Bulgaria, (1980), pag. 205–218, [in Bulgarian].
- [37] P. Vassilev, N. Andreev, Floristic analysis of vegetation of Ostritsa reserve in Mt Golo Brdo. Phytologia, 9, (1978), pag. 41–50.
- [38] S. Kožuharov, (ed.), Flora Reipublicae Bulgaricae. X Vol. Akad. Izd. Bulg. Acad. Sci., (1995), Sofia, [in Bulgarian].
- [39] S. M. Hennekens, J. H. J. Schaminée, TURBOVEG, a comprehensive data base management system for vegetation data. Journal of Vegetation Science, 12, (2001), pag. 589–591.
- [40] S. Veleve. Climatic zoning. – In: I. Kopralev. (ed.) Geography of Bulgaria. Physical and socio-economic geography, Forkom Publisher, (2002), Sofia, pag. 155–156, [in Bulgarian].
- [41] T. Meshinev, Phytocoenological characterization of the reserve “Kastrakli”. In: S. Nedyalkov, K. Baeva, B. Nikolov, R. Radev, T. Michev, Y. Danchev, V. Veleve (eds), International symposium “Conservation of Natural areas and of the genetic material they contain” – Project 8 on the Programme “Man and the Biosphere” (MAB) of Unesco, 23–28.09.1985, Blagoevgrad, Bulgaria, Collection of reports, vol. 2, (1985), pag. 115–123, [in Bulgarian].
- [42] T. Meshinev, Rare plant communities to the south of the town of Smolyan. In: G. Tsankov, S. Iliev, B. Bojinov, B. Bogdanov, P. Zhelev (eds), Proceeding of a jubileum symposium (2–3 June 1994) 100 years from birthday of the acad. Boris Stephanov (1894–1979), vol. II, (1995), PSSA Publishing House, Sofia, pag. 54–57, [in Bulgarian].
- [43] U. Bohn, G. Gollub, C. Hettwer, Y. Neuhäuslova, T. Raus, H. Schlüter, H. Weber, S. Hennekens, (eds.), Map of the natural vegetation of Europe. Scale 1:2 500 000. Interactive CD-ROM: explanatory text, legend, maps. – Bundesamt für Naturschutz, (2004), Bonn.
- [44] V. Kavrakova, D. Dimova, M. Dimitrov, R. Tzonev, T. Belev, K. Rakovska, (eds.), Manual for Determination of Habitats with European Importance in Bulgaria. Second edition. Sofia, (WWF–Danube Carpathian Programme & Green Balkan Federation), (2009), pag. 131., [in Bulgarian].
- [45] V. Rusakova, E. Genova, Z. Cherneva, Phytocoenological and ecological peculiarities of the bear burry (*Arctostaphylos uva-ursi* L. Spreng) in some regions of Bulgaria, Fitologija, 39, (1991), pag. 33–53, [in Bulgarian].
- [46] V. Rusakova, V. Vulchev. 36G3. *Pinus nigra* woodlands (*Pinus nigra* subsp. *pallasiana*) – In: V. Biserkov, C. Gusev, V. Popov, G. Hibaum, V. Roussakova, I. Pandurski, Y. Uzunov, M. Dimitrov, R. Tzonev, S. Tzoneva (eds). Red Data Book of Bulgaria, vol.3. Natural Habitats. BAS and MoEW, (2015), Sofia, pag. 353–355, [in Bulgarian].
- [47] V. Velčev, The vegetation cover of Vratsa mountain, Publishing House of BAS, (1971), Sofia, [in Bulgarian].
- [48] V. Velčev, (ed.), Flora Reipublicae Popularis Bulgaricae. IX Vol., Akad. Izd. Bulg. Acad. Sci., (1989a), Sofia [in Bulgarian].
- [49] V. Velchev, Floral and plant diversity on calcareous terrains in Bulgaria, Phytol. Balcan., 4(1–2), (1989b), pag. 81–92.
- [50] V. Velčev, (ed.), Flora Reipublicae Popularis Bulgaricae. VIII Vol., Akad. Izd. Bulg. Acad. Sci., (1982), Sofia, [in Bulgarian].
- [51] V. Velchev, (ed.), Red Data Book of the People’s Republic of Bulgaria. I Vol. Plants. Akad. Izd. Bulg. Acad. Sci., (1984), Sofia, [in Bulgarian].

- [52] V. Velchev, I. Bondev, Endangered and rare plant communities in Bulgaria, VIII-IX Vol., (1984), Sofia, [in Bulgarian].
- [53] V. Velchev, S. Kozuharov, Bulgarian endemic higher plants – gene pool, origin and distribution. – In: Petrova, A. (ed.), Atlas of Bulgarian endemic plants, Gea-Libris, (2006), Sofia, pag. 14-33.
- [54] V. Westhoff, E. van der Maarel, The Braun-Blanquet approach. In: R. H. Whittaker, (ed.), Ordination and classification of plant communities, (1973), The Hague, pag. 617-737.



**Annex I:** List of taxa, distributed on the territory of Chamdzha managed reserve

## POLYPODIOPHYTA

**Aspleniaceae:** *Asplenium adianthum-nigrum* L., *A. ruta-muraria* L., *A. trichomanes* L., *Ceterach officinarum* DC, *Phyllitis scolopendrium* (L.) Newman; **Athyriaceae:** *Cystopteris fragilis*(L.) Bernh.; **Polypodiaceae:** *Polypodium vulgare* L.

## PINOPHYTA

**Cupressaceae:** *Juniperus oxycedrus* L.; **Pinaceae:** *Abies alba* Mill., *Pinus nigra* Arnold, *P. sylvestris* L.

## MAGNOLIOPHYTA

### Magnoliopsida

**Aceraceae:** *Acer campestre* L., *A. platanoides* L.; *A. tataricum* L.; **Anacardiaceae:** *Cotinus coggygia* Scop;

**Apiaceae:** *Anthriscus sylvestris* (L.) Hoffm., *Bupleurum rotundifolium* L., *Daucus carota* L., *Eryngium campestre* L., *Myrrhoides nodosa* (L.) Cannon, *Orlaya grandiflora* (L.) Hoffm., *Physospermum cornubiense* (L.) DC., *Seseli rigidum* Waldst. & Kit. **Apocynaceae:** *Vinca herbacea* Walds. & Kit. **Araceae:** *Arum maculatum* L. **Araliaceae:** *Hedera helix* L.; **Asclepiadaceae:** *Vincetoxicum hirundinaria* Medic. **Asteraceae:** *Achillea collina* J. Becker ex Reichenb., *A. crithmifolia* Waldst. et Kit., *A. millefolium* L., *A. setacea* Waldst. & Kit., *Anthemis austriaca* Jacq., *A. macedonica* Boiss., *A. tinctoria* L., *Artemisia vulgaris* L., *Carlina vulgaris* L., *Centaurea cyanus* L., *C. deusta* Ten., *C. diffusa* Lam., *Centaurea stoebe* L. (Syn. *C. rhenana* Boreau), *Chamomilla recutita* (L.) Rauschert, *Chondrilla juncea* L., *Cichorium intybus* L., *Cirsium arvense* (L.) Scop., *Crepis sancta* (L.) Bab., *C. setosa* Haller f., *Filago eriocephala* Guss., *Galinsoga parviflora* Cav., *Hieracium glaucinum* gr., *H. hoppeanum* Schultes, *H. olympicum* Frein., *H. praealtum* Vill. Ex Goch., *H. praealtum* Vill. Ex Goch. ssp. *bauchinii*, *H. pilosella* L., *H. piloseloides* Vill., *H. schmidtii* Tausch, *Lactuca perennis* L., *L. saligna* L., *Lapsana communis* L., *Leontodon crispus* Vill., *Logfia arvensis* (L.) Holub, *Mycelis muralis* (L.) Dumort., *Senecio jacobaea* L., *S. vernalis* Waldst. & Kit., *Sonchus arvensis* L., *Taraxacum officinale* F. H. Wigg., *Xanthium strumarium* L., *Xeranthemum annuum* L.; **Betulaceae:** *Carpinus betulus* L., *C. orientalis* Mill., *Corylus avellana* L., *Ostrya carpinifolia* Scop.; **Boraginaceae:** *Buglossoides arvensis* (L.) I. M. Johnst., *B. purpureoacerulea* (L.) I. M. Johnst., *Cynoglossum officinale* L., *Echium vulgare* L., *Myosotis ramosissima* Rochel, *Nonea atra* Griseb., *Symphytum officinale* L., *S. ottomanum* Friv.

**Brassicaceae:** *Alliaria petiolata* (M.Bieb.) Cavara & Grande, *Alyssum parviflorum* Bieb., *Arabis procurrens* Waldst. & Kit., *A. sagittata* (Bertol) DC., *A. turrita* L., *Berteroa incana* (L.) DC, *Capsella bursa-pastoris* (L.) Medic., *Erophila verna* (L.) Chevall., *Erysimum diffusum* Ehrh., *Thalspi kovatsii* Heuffel; **Campanulaceae:** *Campanula glomerata* L. spp. *hispida* (Witašek) Hayek, *C. persicifolia* L., *C. lanata* Friv., *C. rapunculoides* L., *C. rapunculus* L., *C. trachelium* L., *Jasione heldreichii* Boiss. et Orph.;

**Caprifoliaceae:** *Sambucus ebulus* L., *S. nigra* L.; **Caryophyllaceae:** *Cerastium luridum* Guss., *Dianthus petraeus* Waldst. & Kit., *Minuartia saxifraga* (Friv.) Graebner, *Moehringia pendula* (Waldst. & Kit.) Fenzl, *Petrorhagia prolifera* (L.) P. W. Ball & Heywood, *Scleranthus perennis* L., *Silene armeria* L., *S. bupleuroides* L. Chater et Walters, *S. italica* (L.) Pers., *Viscaria vulgaris* Röhl. ssp. *atropurpurea* (Griseb.) Stoj.; **Celastraceae:** *Euonymus europaeus* L., *E. verrucosus* Scop.; **Cistaceae:** *Helianthemum nummularium* (L.) Mill.; **Convolvulaceae:** *Convolvulus arvensis* L.; **Cornaceae:** *Cornus mas* L.; **Crassulaceae:** *Jovibarba heuffelii* (Schott) A. et D. Löve, *Sedum album* L., *S. hispanicum* L., *S. urvillei* DC; **Dioscoreaceae:** *Tamus communis* L.;

**Dipsacaceae:** *Cephalaria transsylvanica* (L.) Roem. & Schult., *Scabiosa trinitifolia* Friv.; **Euphorbiaceae:** *Euphorbia amygdaloides* L., *E. cyparissias* L., *Mercurialis perennis* L.; **Fabaceae:** *Astragalus onobrychis* L., *Chamaecytisus jankae* (Velen.) Rothm., *C. supinus* (L.) Link, *Chamaespartium sagittale* (L.) Gibbs, *Coronilla varia* L., *Dorycnium herbaceum* Vill., *Genista januensis* Viv., *G. tinctoria* L., *Lathyrus pratensis* L., *L. tuberosus* L., *L. vernus* Bernh., *Lotus corniculatus* L., *Medicago falcata* L., *M. lupulina* L., *M. minima* (L.) Bartal., *Melilotus alba* Medicus, *Robinia pseudoacacia* L., *Trifolium alpestre* L., *T. angustifolium* L., *T. arvense* L., *T. campestre* Schreb, *T. diffusum* Ehrh., *T. dubium* Sibth., *T. medium* L., *T. ochroleucon* Huds., *T. pratense* L., *T. repens* L., *T. setiferum* Boiss, *T. striatum* L., *Vicia cracca* L., *V. grandiflora* Scop., *V. sativa* L., *V. tetrasperma* (L.) Schreb., *V. varia* Host, *V. villosa* Roth; **Fagaceae:** *Fagus sylvatica* L., *Quercus cerris* L., *Q. Dalechampii* Ten., *Q. pubescens* Willd., *Q. rubra* L.; **Geraniaceae:** *Erodium cicutarium* (L.) L'Hér., *Geranium lucidum* L., *G. molle* L., *G. robertianum* L., *G. rotundifolium* L.; **Hypericaceae:** *Hypericum linarioides* Bosse, *H. perforatum* L., *H. umbellatum* A. Kern.; **Lamiaceae:** *Acinos alpinus* (L.) Moench ssp. *hungaricus* (Simonkai) Sojak, *A. rotundifolius* Pers., *Ajuga reptans* L., *Ballota nigra* L., *Calamintha nepeta* (L.) Savi, *Clinopodium vulgare* L., *Galeopsis ladanum* L., *G. tetrachit* L., *Glechoma hederacea* L., *Marrubium peregrinum* L., *Lamium purpureum* L., *Origanum vulgare* L., *Prunella vulgaris* L., *Satureja coerulea* Janka, *Teucrium chamaedrys* L., *Thymus callieri* Borbas ex Velen., *T. pulegioides* L., *T. sibthorpii* Benth., *T. striatus* Vahl; **Malvaceae:** *Malva sylvestris* L.; **Oleaceae:** *Fraxinus excelsior* L., *F. ornus* L., *Ligustrum vulgare* L.; **Papaveraceae:** *Chelidonium majus* L., *Papaver laevigatum* M. Bieb.; **Plantaginaceae:** *Plantago lanceolata* L.; **Polygonaceae:** *Polygonum aviculare* L., *Rumex acetosa* L., *R. acetosella* L., *R. crispus* L.; **Primulaceae:** *Lysimachia punctata* L.; **Pyrolaceae:** *Orthilia secunda* (L.) House; **Ranunculaceae:** *Clematis vitalba* L., *Helleborus odoratus* Waldst. & Kit.; **Rosaceae:** *Agrimonia eupatoria* L., *Aremonia agrimonoides* (L.) DC., *Crataegus monogyna* Jacq., *Filipendula vulgaris* Moench, *Fragaria vesca* L., *F. Viridis* Duchesne, *Geum urbanum* L., *Potentilla argentea* L., *P. laciniosa* Waldst. & Kit. ex Nestl., *P. reptans* L., *Prunus avium* (L.) L., *P. cerasifera* Ehrh., *P. mahaleb* L., *P. spinosa* L., *Pyrus pyraeaster* Burgsd., *Rosa canina* L., *R. micrantha* Borrer ex Sm., *Rubus discolor* Weihe & Nees, *Sanguisorba minor* Scop. **Rubiaceae:** *Cruciatata glabra* (L.) Ehrend., *C. laevipes* Opiz, *Galium aparine* L., *G. flavescens* Borbas, *G. lucidum* All., *G. odoratum* (L.) Scop., *G. pseudoaristatum* Schur, *G. spurium* L., *G. verum* L., *Sherardia arvensis* L.;

**Saxifragaceae:** *Saxifraga rotundifolia* L.; **Scrophulariaceae:** *Digitalis lanata* Ehrh., *Euphrasia pectinata* Ten., *Verbascum densiflorum* Bertol., *V. lychnitis* L., *V. nigrum* L., *Veronica chamaedrys* L., *V. verna* L., *V. vindobonensis* (M. A. Fisch.) M.

A. Fisch.; **Tiliaceae:** *Tilia cordata* Mill.; **Urticaceae:** *Parietaria officinalis* L., *Urtica dioica* L.; **Verbenaceae:** *Verbena officinalis* L.; **Violaceae:** *Viola arvensis* Murr., *V. canina* L., *V. reichenbachiana* Jord. ex Boreau;

**Liliopsida**

**Cyperaceae:** *Carex caryophylla* Latourr., *C. digitata* L., *C. divulsa* Stokes ex With., *C. echinata* Murr., *C. remota* L.; **Juncaceae:** *Luzula campestris* (L.) DC., *L. forsteri* (Sm.) DC., *L. pilosa* (L.) Willd.; **Liliaceae:** *Muscari neglectum* Guss. ex Ten., *M. tenuiflorum* Tausch, *Ruscus aculeatus* L.; **Poaceae:** *Agrostis capillaris* L., *Anthoxanthum odoratum* L., *Apera spica-venti* (L.) P. Beauv., *Arrhenatherum elatius* (L.) P. Beauv. ex J & C. Presl, *Brachypodium pinnatum* (L.) P. Beauv., *B. sylvaticum* (Huds.) P. Beauv., *Bromus commutatus* Schrad., *B. mollis* L., *B. sterilis* L., *Chrysopogon gryllus* (L.) Trin., *Cynodon dactylon* (L.) Pers., *Cynosurus cristatus* L., *C. echinatus* L., *Dactylis glomerata* L., *Dasypyrum villosum* (L.) Cand., *Dichanthium ischaemum* (L.) Roberti, *Elymus repens* (L.) Gould, *Festuca heterophylla* Lam., *F. rubra* L., *F. valesiaca* Schleich. ex Gaudin, *Hordelymus europaeus* (L.) Harz., *Koeleria macrantha* (Ledeb.) Schult., *K. nitidula* Velen., *Lerchenfeldia flexuosa* (L.) Schur, *Lolium perenne* L., *Melica ciliata* L., *M. uniflora* Retz., *Phleum phleoides* (L.) Karsten, *P. pratense* L., *Poa angustifolia* L., *P. annua* L., *P. bulbosa* L., *P. compressa* L., *P. nemoralis* L., *P. sylvicola* Guss., *P. trivialis* L., *Setaria pumila* (Poir.) Schult., *Vulpia myurus* (L.) C. C. Gmel.

**Annex II:** Synoptic table of vegetation syntaxa of Chamdzha managed reserved. About woody species were used following abbreviations: tr - for tree layer, sh - for shrub layer and juv - for juvenile species.

<b>Relevé No</b>	1	2	3	4	5	6	7	8	9	10	11	12	13	14
<b>Altitude [m]</b>	779	880	748	847	611	924	664	923	920	923	850	582	640	660
<b>Exposure</b>	90	135	45	360	90	180	225	180	135	180	180	90	360	45
<b>Inclination [degree]</b>	15	5	30	10	25	15	30	10	10	10	15	20	30	60
<b>Plot size [m<sup>2</sup>]</b>	16	16	100	100	100	100	100	100	100	100	100	100	100	100
<b>Total coverage [%]</b>	70	80	85	80	90	75	85	75	80	85	75	85	85	90
<b>Cover of tree layer [%]</b>	0	0	65	70	80	60	60	50	50	70	70	60	70	60
<b>Cover of shrub layer [%]</b>	0	8	15	10	40	10	50	20	15	5	35	65	60	40
<b>Cover of herb layer [%]</b>	35	80	25	15	30	15	40	35	80	30	5	15	30	20
<b>Cover of mosses [%]</b>	40	20	3	3	2	0	35	1	70	1	0	2	15	70
<b>Cover of lichens [%]</b>	10	0	2	2	0	0	0	0	0	0	0	0	0	0
<b>Latitude</b>	42,73813	42,74239	42,73492	42,73933	42,73296	42,74742	42,73096					42,73137	42,73386	42,73416
<b>Longitude</b>	24,58528	24,58161	24,58913	24,58279	24,59299	24,57252	24,59228					24,59372	24,59178	24,59085
<b>Species number</b>	23	23	16	12	22	13	19	8	10	14	11	24	27	21

**Ch. species of cl. *Koelerio-Corynephoretea***

Veronica verna	+	+	.	.	.	.	.	.	.	.	.	.	.	.
Poa bulbosa	+	1	.	.	.	.	.	.	.	.	.	.	.	.
Trifolium arvense	+	1	.	.	.	.	.	.	.	.	.	.	.	.
Scabiosa triniifolia	+	+	.	.	.	.	.	.	.	.	.	.	.	.
Chamomilla recutita	2	2	.	.	.	.	.	.	.	.	.	.	.	.
Ceratodon purpureus	2	1	.	.	.	.	+	.	.	.	.	.	.	.
Scleranthus perennis	3	2	1	.	.	.	2	.	.	.	.	.	.	.
Polytrichum piliferum	1	2	+	.	.	.	2	.	.	.	.	.	.	.
Hieracium pilosella	.	2	.	.	+	.	.	.	.	.	.	.	.	.
Syntrichia ruralis	2	.	.	.	.	.	.	.	.	.	.	.	.	.

**Ch. species of cl. *Erico-Pinetea* & all. *Erico-Fraxionion orni***

Pinus nigra (tr)	.	.	3	3	4	4	4	4	2	.	2	4	.	.
Pinus nigra (sh)	.	.	2	.	.	.	2	.	.	.	.	.	.	.

**Ch. species of cl. *Quercu-Fagetea* & all. *Carpinion betuli***

Carpinus betulus (sh)	.	.	.	.	.	.	.	.	.	3	2	.	.	.
Quercus dalechampii (tr)	.	.	3	3	2	2	.	.	.	3	4	.	3	4
Quercus dalechampii (sh)	.	.	+	2	2	.	.	2	2	+	2	.	.	.
Melica uniflora	.	.	.	.	.	.	.	.	.	2	2	2	+	.

continuation of Annex II

**Ch. species of cl. *Quercetea pubescentis* & all. *Carpinion orientalis***

Carpinus orientalis (sh)	.	.	1	.	1	.	.	+	2	.	.	4	3	4
Acer campestre (sh)	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Prunus mahaleb (juv)	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Tamus communis	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Acer campestre (tr)	.	.	.	.	.	.	.	.	.	.	.	.	2	.
Poa nemoralis	.	.	.	2	.	.	.	.	.	.	+	.	2	+

**transgressive ch. species of cl. *Erico-Pinetea* & *Quercetea pubescentis***

Fraxinus ornus (sh)	.	.	+	.	2	.	.	.	.	.	.	1	2	2
---------------------	---	---	---	---	---	---	---	---	---	---	---	---	---	---

**Other species**

Fraxinus ornus (juv)	.	.	.	.	.	.	.	.	+	+	.	.	.
Ruscus aculeatus	.	.	.	.	.	.	.	.	3	.	.	.	.
Helleborus odoratus	.	.	.	.	.	.	.	.	+	+	.	.	.
Carpinus orientalis (tr)	.	.	.	.	2	.	.	.	2	.	.	.	.
Abies alba (juv)	.	.	.	.	.	.	.	.	+	.	.	.	.
Pinus nigra (juv)	1	1	.	.	.	.	.	.	.	.	.	.	.
Quercus cerris (tr)	.	.	.	.	.	.	.	.	2	2	.	.	.
Lerchenfeldia flexuosa	.	.	2	+	+	2	+	3	2	.	.	.	2
Geranium lucidum	.	.	.	.	.	.	.	.	+	+	.	.	.
Geranium molle	.	.	.	.	.	.	.	.	.	.	+	+	+
Ceterach officinarum	.	.	.	.	+	.	.	.	.	.	+	+	+
Campanula rapunculoides	.	.	.	.	.	.	.	.	.	.	.	+	+
Porella platyphylla	.	.	.	.	.	.	.	.	.	.	+	.	+
Viola reichenbachiana	.	.	.	.	.	.	.	.	+	.	+	+	.
Arabis procurrens	.	.	.	.	.	.	.	.	.	.	.	+	2
Arabis sagittata	.	.	.	.	.	.	.	.	.	.	.	2	+
Galeopsis tetrachit	.	.	.	.	.	.	.	.	.	.	+	1	.
Asplenium adiantum-nigrum	.	.	.	.	.	.	.	.	.	.	+	.	+
Hypericum linarioides	+	.	.	.	.	.	+	.	.	.	.	.	.
Genista januensis (sh)	.	.	+	+	.	.	+	.	.	.	.	.	.
Quercus dalechampii (juv)	.	1	.	.	.	.	.	.	.	.	.	.	.
Polypodium vulgare	.	.	.	.	.	.	.	+	+	.	.	2	.
Achillea collina	.	.	.	.	.	.	+	.	.	.	.	.	.
Quercus cerris (sh)	.	.	.	.	.	.	2	.	.	.	.	.	.
Campanula persicifolia	.	.	.	.	.	.	.	+	.	.	+	.	.
Hieracium olympicum	.	.	.	.	.	+	.	+	+	.	.	.	.

VASSILEV K., GAVRILOVA A.  
FLORA, HABITATS AND VEGETATION OF CHAMDZHA MANAGED RESERVE, CENTRAL BALKAN RANGE

*continuation of Annex II*

Lamium purpureum	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Sedum hispanicum	+	.	.	.	.	+	1	.	.	.	.	.	1	.
Acer platanoides (sh)	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Silene bupleuroides	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Galium flavescens	.	.	.	.	+	.	2	.	.	.	.	.	+	.
Dianthus petraeus	.	.	1	.	.	.	+	.	.	.	.	.	.	.
Lathyrus vernus	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Bryum argenteum	.	.	.	.	.	.	+	.	.	.	.	.	.	.
Luzula pillosa	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Agrimonia eupatoria	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Lapsana communis	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Hieracium piloselloides	.	.	2	2	+	.	.	.	.	.	.	.	.	+
Grimmia pulvinata	.	.	.	.	.	.	+	.	.	.	.	.	.	.
Chamaecytisus jankae (sh)	+	1	+	1	2	2	+	.	.	.	.	.	.	.
Hedera helix	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Veronica chamaedrys	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Galium aparine	.	.	.	.	+	+	.	.	.	.	.	+	.	+
Phleum phleoides	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Atrichum undulatum	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Festuca heterophylla	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Dactylis glomerata	.	.	.	.	+	.	.	.	.	.	+	.	.	+
Quercus rubra (juv)	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Hypnum cupressiforme	.	.	+	1	+	.	.	.	.	.	.	+	.	2
Saxifraga rotundifolia	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Viscaria vulgaris	.	.	.	+	+	.	.	.	.	.	.	.	.	.
Moehringia pendula	.	.	.	.	2	+	+	.	.	.	.	.	+	2
Agrostis capillaris	.	2	.	.	.	.	1	.	.	.	.	.	.	.
Carex digitata	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Lophocolea minor	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Grimmia anodon	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Jovibarba heuffellii	.	.	1	.	.	.	.	.	.	.	.	.	.	.
Aremonia agrimonoides	.	.	.	+	.	.	.	.	.	.	.	.	.	+
Galium pseudaristatum	.	.	.	+	.	.	.	.	.	.	.	.	.	+
Tamus communis	.	.	.	.	.	.	.	.	.	.	.	.	+	.
Verbascum lychnitis	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Achillea crithmifolia	.	.	.	.	.	+	.	.	.	.	.	.	.	.
Rosa canina (sh)	.	.	.	.	.	.	.	.	.	.	.	.	+	.
Campanula rapunculus	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Melica ciliata	.	.	.	.	.	.	.	.	.	.	.	.	+	.



*continuation of Annex II*

Metzgeria conjugata	.	.	.	.	.	.	.	.	.	.	.	.	.	+
Moehringia trinervia	.	.	+	.	.	.	.	.	.	.	.	.	.	.
Festuca rubra	.	3	.	.	.	.	.	.	.	.	.	.	.	.
Silene armeria	+	.	.	.	.	+	.	.	.	.	.	.	.	.
Galium odoratum	.	.	2	.	.	.	.	.	.	.	.	.	.	.
Myrrhoides nodosa	.	.	.	.	.	.	.	.	.	.	+	.	.	.
Filago eriocephala	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Luzula luzuloides	.	.	.	.	.	.	.	.	+	.	.	.	.	.
Muscari tenuiflorum	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Luzula forsteri	.	.	.	.	.	.	.	.	+	.	.	.	.	.
Rumex acetosella	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Chondrilla juncea	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Jasione heldreichii	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Acinos rotundifolius	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Hieracium hoppeanum	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Dicranum scoparium	.	.	+	.	+	.	2	.	.	.	.	.	.	+
Carex caryophylla	.	1	.	.	.	.	.	.	.	.	.	.	.	.
Euphrasia pectinata	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Bromus sterilis	.	.	.	.	.	+	.	.	.	.	.	.	.	.
Hedwigia ciliata	2	.	.	.	.	.	.	.	.	.	.	.	.	.
Symphytum ottomanum	.	.	.	.	.	.	.	.	.	.	+	.	+	.
Mercurialis perennis	.	.	.	.	.	.	.	.	.	.	.	+	.	.
Hieracium species	.	.	.	.	.	.	+	.	.	.	.	.	.	.
Arabis turrata	.	.	.	.	.	.	.	.	.	+	.	.	.	.
Thlaspi kovatsii	.	.	.	.	+	.	.	.	.	.	.	.	.	.
Thymus sibthorpii	.	2	.	.	.	.	.	.	.	.	.	.	.	.
Trifolium alpestre	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Luzula campestris	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Hypericum umbellatum	.	.	.	.	.	+	.	.	.	.	.	.	.	.
Vulpia myuros	.	1	.	.	.	.	.	.	.	.	.	.	.	.
Muscari neglectum	.	.	.	.	.	.	.	+	.	+	.	.	.	.
Racomitrium canescens	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Mycelis muralis	.	.	.	.	.	.	.	.	.	+	.	.	.	.
Trifolium medium	.	.	.	+	.	.	.	.	.	.	.	.	.	.
Cystopteris fragilis	.	.	.	.	.	.	.	.	.	+	.	.	.	.
Koeleria nitidula	+	.	.	.	.	.	.	.	.	.	.	.	.	.
Chamaespartium sagittale	.	+	.	.	.	.	.	.	.	.	.	.	.	.
Genista tinctoria (sh)	.	.	.	.	.	+	.	.	.	.	.	.	.	.

## FLORISTIC-VEGETATIONAL VARIABILITY OF THE ASSOCIATION *EPIMEDIO-CARPINETUM BETULI* (HORVAT 1938) BORHIDI 1963 IN THE NORTH OF CROATIA

VUKELIĆ J., ŠAPIĆ I., BARIČEVIĆ D.

University of Zagreb, Faculty of Forestry, Zagreb, Croatia

Corresponding author e-mail address: dario.baricevic@zg.htnet.hr

**ABSTRACT:** The paper analyzes 246 phytocoenological relevés of the association *Epimedio-Carpinetum betuli* (I. Horvat 1938) Borhidi 1963 in northern Croatia using the standard principles of the Central European Phytocoenological School [5] and applying recent statistical methods [10, 6, 30]. The association is divided into six subassociations with sociologically closely affiliated differential species. The first part of the research implies formalized classification to analyze and correct the selection of diagnostic and differential species in traditional subassociations formed on the basis of previous studies. The second part uses a statistical method to form three clusters, determine their differential species and compare them with the earlier classification of the association. The results of the analysis have revealed a relatively low floristic variability, whereas structural classification has confirmed a smaller number of the described associations. In addition to providing an objective classification of the community *Epimedio-Carpinetum*, the study will also contribute to future analyses of Illyrian oak-hornbeam forests in their distribution range, as well as to the synthesis of the *Carpinetum* communities of Europe.

**Keywords:** *Epimedio-Carpinetum betuli*, northern Croatia, statistical analysis, differential species.

### 1 INTRODUCTION

The forests of sessile oak and common hornbeam in the north of Croatia belong to the earliest descriptions of forest communities of south-eastern Europe [11]. The most distinctive features of the association *Quercus-Carpinetum croaticum* I. Horvat 1938 are its wealth of flora and diagnostic value of the species of the Illyrian floristic element. On these bases Horvat [12] establishes a separate alliance of oak-hornbeam forests of the Illyrian region (*Carpinion betuli illyrico-podolicum* Horvat 1958), thus delineating them from the Central European forests. His attitude provoked a number of discussions [3, 4, 8, 14, 34], but phytocoenologists of south-eastern Europe retained this alliance in their later works. They pointed out the absence of their characteristic species, but attributed diagnostic importance to the present species from the Illyrian beech forests of the alliance *Fagion illyricum* (= *Aremonio-Fagion* /Horvat 1938/ Borhidi in Török, Podani et Borhidi 1989). Particularly active in the study of the *Carpinion* communities of south-eastern Europe were Slavic phytocoenologists, so Marinček [in 33], in line with the then valid code of phytocoenological nomenclature, named the alliance *Carpinion illyrico-podolicum* as *Erythronio-Carpinion betuli* (Horvat 1938) Marinček 1993 in Wallnöfer et al. 1993; however, this alliance was soon afterwards divided into three sub-alliances [16, 17]. The alliance *Erythronio-Carpinion* is currently recognized in the majority of the surveys of forest vegetation of Europe [25, 1 et al.].

Following research by Horvat, the Illyrian oak-hornbeam forests were nomenclaturally defined by Borhidi [3], who named the association *Epimedio-Carpinetum betuli*. Research into this association has continued in Croatia, resulting in about five hundred relevés to date [23, 24, 29, 15, 27, 21, 31, 2, 18, 28, 26 et al.]. In addition to the association *Epimedio-Carpinetum betuli*, the relevés also describe the association *Festuco drymeiae-Carpinetum betuli*, but it is not analyzed in this paper.


In the cited works, the association *Epimedio-Carpinetum betuli* is divided into six subassociations: *erythronietosum* Horvat 1938, on moderately acidophilic podzolic soils of north-western Croatia, *staphyletosum* Horvat 1938, on neutrophil-alkaline carbonate soils of the

same area, the typical and most common subassociation *caricetosum pilosae* Horvat 1963, mesophilic *asperuletosum* Wraber 1961 and two acidophilic *castanetosum* Wraber 1958 and *luzuletosum albidae* Wraber 1961 with a more pronounced anthropogenic impact. A comparative analysis of the differences and justifiability of the subassociations was not made, while the criteria used to describe them frequently relied on a small number of differential species determined by the first authors of the subassociations I. Horvat and M. Wraber. Since such research activities are supported by new statistical techniques, the goal of our paper was to, with the help of these techniques, analyze the variability and distribution of the association *Epimedio-Carpinetum* in northern Croatia. The results should critically question the justifiability of the traditional classification from previous studies (1), assess the value of diagnostic species (2), possibly define a new statistically justified classification (3), but also answer other important questions concerning the composition and structure of this well known association. A statistical analysis of the overall phytocoenological material of a larger area has an advantage, since it eliminates the narrowly distributed local syntaxa which were developed under the impact of some local factors. On the other hand, it provides a more comprehensive picture of the impact of the macro-climate, other regional factors and naturally, the biogeographic area. Finally, the present European classifications, Natura 2000 in the first place, are based on such regional types.

### 2 RESEARCH AREA

The association *Epimedio-Carpinetum betuli* is a zonal association of the colline belt of a larger part of continental Croatia. The most important areas are located in the mountains between the rivers Sava and Drava in Croatia, then on the northern boundary part of the Dinaric mountain range south of Karlovac, and on the slopes of karst fields in the Dinaric region. It occurs between 150 and 450 m above sea level, most frequently on slope pseudogley, eutric cambisol and luvisol above different parent bedrock. The average annual temperature in the eastern part of the distribution range is 10.5 °C, in the central part it is 10.3 °C, and in the southern part it

amounts to 10.0°C. Precipitation also ranges in the same direction from 700 mm annually, to 1,100 mm (central part), to 1,500 mm at Ogulin. Generally, the climate is moderately warm and humid, while in the eastern part of the distribution range it is semi-humid. Precipitation distribution is favourable (over 50 % in the vegetation period), making the climate conducive to the development of forest vegetation. Apart from climate, there are other ecological factors that are suitable for the life of man, and therefore sessile oak - hornbeam forests were massively cut down very early in history. In northern Croatia (Fig. 1) they inhabit the lower slopes of higher mountains in Pannonia and surround them in a ring-like fashion. In the lower hills (up to 350 m), they occupy comprehensive complexes which are interrupted by colline-submontane beech forests on the northern sides.



**Figure 1:** Map of the research area

### 3 MATERIAL AND METHODS

A phytocoenological analysis of the association *Epimedio-Carpinetum betuli* is based on the principles of the Zurich-Montpellier or the standard Central European School [5]. All the cited research was conducted in accordance with the mentioned School. The most important feature is the unique methodology of field phytocoenological sampling as a prerequisite for comparison and analysis of the results. Two basic analyses were conducted in the project: the first analyzes and evaluates the past (traditional) phytocoenological classification of the association *Epimedio-Carpinetum betuli*, and the second divides the same relevés into clusters formed by means of statistical classification.

A total of 246 phytocoenological relevés were used to analyze previous phytocoenological classification. In all the original papers they were classified into the association *Epimedio-Carpinetum betuli*, or into some of its subassociations:

- *erythronietosum* – 33 relevés, [11, 23, 15, 31]
- *staphyletosum* – 41 relevés, [11, 23, 29, 15, 31]
- *caricetosum pilosae* – 107 relevés, [23, 15, 31, 2, 28]
- *asperuletosum* – 23 relevés, [27, 28]
- *castanetosum sativae* – 25 relevés, [18]
- *luzuletosum albidiae* – 17 relevés, [29, 27, 15]

Vegetation relevés were entered into TURBOVEG database [10]. Cluster analysis was performed in PRIMER 6 software [6], with Euclidian distance as a measure of similarity. Diagnostic species were

determined by means of JUICE 7.0 software package [30] on the basis of the analysis of Fidelity measure. The plant covers that occur in more layers were combined. Each plant species was considered with total cover, regardless of the number of structural layers in which it occurred in a particular relevé. Mosses were not evidenced in the majority of studies and were therefore not taken into consideration.

In the first analysis the relevés were classified into six groups (Table I) according to the original subassociations described by the previous authors of research. Species participating with over 30% and fidelity index above 30 were taken as their differential species.

The second analysis was performed on the basis of statistical classification of the same relevés using cluster analysis, which resulted in their grouping within 3 clusters.

Research results (Tables I and II) show only differential, constant, dominant and Illyrian floral element species. Constant species were defined as those with a frequency  $\geq 50\%$  inside the vegetation unit (cluster), whereas dominant species were those with a cover value  $> 25\%$  in at least 20% of the relevés belonging to a particular vegetation unit. Differential species were listed separately. The analysis of ecological conditions using Ellenberg's ecoindicator coefficients [7] was performed in JUICE 7.0 programme [30]. The plant nomenclature was adjusted to the *Flora Croatica* database [19]. The nomenclature of plant communities was adjusted to the survey of forest vegetation of Croatia [32] and subassociation as determined by Croatian phytosociologies in the cited papers [11, 13, 18, 27, 28].

### 4 RESULTS AND DISCUSSION

A total of 276 species of higher plants were recorded in 246 relevés of the association *Epimedio-Carpinetum betuli*. The most important species of the Illyrian floral element distributed in all the six subassociations include *Lonicera caprifolium*, *Epimedium alpinum*, *Ruscus hypoglossum*, *Cyclamen purpurascens*, *Aposeris foetida*, *Knautia drymeia* ssp. *drymeia* and *Primula vulgaris*. A major part of the subassociations also contain *Lamium orvala*, *Erythronium dens-canis*, *Aremonia agrimonoides* and *Vicia oroboides*. The species *Omphalodes verna*, *Cardamine trifolia*, *Helleborus purpurascens* and *Crocus vernus* ssp. *vernus* occur in only one subassociation. In two subassociations in the eastern part of the study area there is an abundance of the species *Helleborus odoratus*. It is the presence of these species in oak-hornbeam forests of the Illyrian floral province that constitutes the main reason for the establishment of the alliance *Erythronio-Carpinion betuli*.

From a relatively large number of the species, 55 species were observed in all the subassociations. Constant species of the association are *Carpinus betulus*, *Quercus petraea*, *Prunus avium*, *Fagus sylvatica*, *Crataegus monogyna*, *Corylus avellana*, *Ligustrum vulgare*, *Cornus sanguinea*, *Acer campestre*, *Fraxinus ornus*, *Viola reichenbachiana*, *Pulmonaria officinalis*, *Galium odoratum*, *Asarum europaeum*, *Stellaria holostea*, *Hedera helix*, *Carex sylvatica*, *Rubus hirtus*, *Epimedium alpinum* and *Galium sylvaticum*. They are generally not differential for particular subassociations. The dominant species are *Carpinus betulus* and *Quercus petraea*.

VUKELIĆ J., ŠAPIĆ I., BARIČEVIĆ D.  
FLORISTIC-VEGETATIONAL VARIABILITY OF THE ASSOCIATION *EPIMEDIO-CARPINETUM BETULI* (HORVAT 1938)  
BORHIDI 1963 IN THE NORTH OF CROATIA

**Table I:** Schortened synoptic table with percentage frequency and modified fidelity phi coefficient based on the traditional division of associations *Epimedio-Carpinetum betuli*

Group No.	1	2	3	4	5	6
No. of relevés	107	23	33	25	17	41
Subassociations	caric	aspe	eryt	casta	luzu	staph
Differential species						
<i>Tilia tomentosa</i>	28 49.5	. ---	. ---	. ---	. ---	. ---
<i>Melica uniflora</i>	52 45.5	13 ---	6 ---	. ---	12 ---	10 ---
<i>Carex pilosa</i>	60 40.1	13 ---	9 ---	32 10.3	6 ---	15 ---
<i>Viola hirta</i>	41 37.4	26 17.4	. ---	. ---	6 ---	5 ---
<i>Glechoma hirsuta</i>	44 36.6	22 8.7	3 ---	12 ---	6 ---	2 ---
<i>Acer tataricum</i>	45 40.9	30 22.0	6 ---	. ---	. ---	. ---
<i>Helleborus odoros</i>	37 39.0	26 22.6	. ---	. ---	. ---	. ---
<i>Galeobdolon luteum</i>	42 17.3	65 41.1	12 ---	12 ---	18 ---	2 ---
<i>Prunella vulgaris</i>	5 ---	35 35.6	3 ---	. ---	18 10.6	2 ---
<i>Galium odoratum</i>	65 6.9	91 30.3	61 2.5	44 ---	29 ---	56 ---
<i>Asarum europaeum</i>	41 ---	96 30.1	67 3.2	48 ---	47 ---	80 16.1
<i>Sanicula europaea</i>	30 ---	87 29.8	52 ---	60 5.6	35 ---	59 4.3
<i>Erythronium dens-canis</i>	2 ---	. ---	64 60.2	20 5.9	6 ---	. ---
<i>Crocus vernus agg.</i>	3 ---	. ---	58 50.8	16 ---	. ---	20 4.3
<i>Anemone nemorosa</i>	24 ---	. ---	82 41.1	56 17.2	6 ---	56 17.3
<i>Milium effusum</i>	11 ---	35 7.3	58 30.1	. ---	18 ---	44 16.4
<i>Doronicum austriacum</i>	3 ---	. ---	52 41.8	36 23.2	. ---	10 ---
<i>Convallaria majalis</i>	12 ---	. ---	64 37.9	48 22.0	. ---	34 7.9
<i>Luzula pilosa</i>	4 ---	. ---	58 34.7	52 28.9	18 ---	15 ---
<i>Fragaria vesca</i>	20 ---	30 ---	82 33.7	20 ---	41 ---	73 25.9
<i>Frangula alnus</i>	1 ---	. ---	33 29.7	8 ---	12 ---	17 7.2
<i>Cephalanthera longifolia</i>	11 ---	. ---	. ---	60 66.6	. ---	. ---
<i>Castanea sativa</i>	10 ---	9 ---	24 ---	100 55.2	53 12.2	41 1.7
<i>Melampyrum pratense</i>	10 ---	22 ---	33 ---	88 50.9	41 6.7	10 ---
<i>Euonymus latifolius</i>	1 ---	. ---	. ---	32 47.3	. ---	5 ---
<i>Prenanthes purpurea</i>	2 ---	. ---	9 1.4	36 45.2	. ---	2 ---
<i>Potentilla micrantha</i>	47 19.3	26 ---	6 ---	68 40.6	6 ---	12 ---
<i>Lamium orvala</i>	6 ---	. ---	42 12.8	68 37.9	12 ---	49 19.0
<i>Fraxinus ornus</i>	64 19.1	9 ---	21 ---	80 34.0	18 ---	63 19.0
<i>Hedera helix</i>	46 ---	48 ---	52 ---	96 31.4	47 ---	83 19.4
<i>Acer platanoides</i>	7 ---	4 ---	6 ---	36 30.1	6 ---	20 8.3
<i>Gentiana asclepiadea</i>	6 ---	9 ---	61 22.6	76 36.9	35 ---	32 ---
<i>Hieracium murorum</i>	9 ---	. ---	33 22.4	36 25.7	6 ---	7 ---
<i>Luzula luzuloides</i>	11 ---	. ---	48 8.8	64 23.0	100 56.0	10 ---
<i>Hieracium racemosum</i>	19 ---	13 ---	15 ---	40 16.8	47 24.2	10 ---
<i>Staphylea pinnata</i>	3 ---	. ---	. ---	. ---	. ---	66 76.4
<i>Hacquetia epipactis</i>	. ---	. ---	9 ---	28 13.5	. ---	63 55.9
<i>Scilla bifolia agg.</i>	. ---	. ---	6 ---	. ---	. ---	34 49.1
<i>Helleborus atrorubens</i>	4 ---	. ---	9 ---	. ---	6 ---	44 49.0
<i>Melampyrum nemorosum</i>	9 ---	. ---	21 5.8	. ---	12 ---	56 47.9
<i>Campanula trachelium</i>	3 ---	4 ---	24 8.8	8 ---	6 ---	56 46.8
<i>Vicia oroboides</i>	12 4.9	. ---	3 ---	. ---	. ---	39 46.8
<i>Tamus communis</i>	36 10.6	. ---	21 ---	28 2.0	. ---	71 45.5
<i>Solidago virgaurea</i>	6 ---	9 ---	15 ---	. ---	12 ---	49 42.3
<i>Mercurialis perennis</i>	7 ---	. ---	9 1.0	. ---	. ---	34 41.3
<i>Salvia glutinosa</i>	7 ---	13 ---	. ---	24 10.4	. ---	49 41.0
<i>Viburnum lantana</i>	20 5.3	. ---	6 ---	20 5.8	. ---	46 38.5
<i>Acer pseudoplatanus</i>	19 ---	9 ---	52 10.9	56 15.0	29 ---	73 30.7
<i>Senecio nemorensis agg.</i>	2 ---	. ---	9 ---	40 31.3	. ---	39 30.1
Dominant species						
<i>Carpinus betulus</i>	100 6.4	100 6.4	100 6.4	88 ---	100 6.4	100 6.4
<i>Quercus petraea</i>	100 14.6	83 ---	100 14.6	68 ---	94 5.7	98 10.9
Constant species						
<i>Fagus sylvatica</i>	90 5.6	87 2.1	70 ---	92 8.5	100 18.6	73 ---
<i>Corylus avellana</i>	48 ---	83 ---	91 10.1	88 6.7	94 13.9	90 9.3
<i>Carex sylvatica</i>	65 ---	91 18.5	70 ---	52 ---	76 3.5	83 10.0
<i>Prunus avium</i>	79 8.4	65 ---	82 11.6	72 2.1	41 ---	80 10.4

<i>Pulmonaria officinalis</i>	55	---	65	2.7	79	15.2	68	5.3	24	---	83	19.1
<i>Ligustrum vulgare</i>	50	---	48	---	58	4.7	44	---	41	---	73	18.6
<i>Crataegus monogyna</i>	54	---	48	---	61	1.8	64	4.9	47	---	78	17.6
<i>Viola reichenbachiana</i>	58	6.0	43	---	73	19.2	32	---	35	---	66	13.1
<i>Acer campestre</i>	79	16.4	74	12.2	55	---	72	10.5	24	---	61	---
<i>Epimedium alpinum</i>	66	14.1	65	13.0	61	8.9	28	---	47	---	37	---
<i>Cornus sanguinea</i>	62	2.9	57	---	55	---	72	12.2	24	---	83	22.1
<i>Rubus hirtus s.lat.</i>	86	35.4	30	---	24	---	64	15.7	35	---	39	---
<i>Stellaria holostea</i>	60	---	74	13.5	88	26.2	24	---	53	---	56	---
<i>Galium sylvaticum</i>	40	---	13	---	82	26.5	60	7.0	35	---	83	27.5
<b>Illyrian floral element</b>												
<i>Lonicera caprifolium</i>	14	---	65	13.8	61	9.7	52	2.0	53	2.9	54	3.5
<i>Ruscus hypoglossum</i>	14	---	22	4.2	3	---	20	2.2	35	19.9	15	---
<i>Primula vulgaris</i>	22	---	39	---	73	29.5	28	---	24	---	56	14.4
<i>Aposeris foetida</i>	11	---	52	---	91	27.5	76	13.9	47	---	88	24.7
<i>Knautia drymeia</i>	24	---	43	6.5	42	5.5	16	---	29	---	63	25.0
<i>Cyclamen purpurascens</i>	10	---	17	---	33	---	48	14.3	35	2.2	54	19.7

Differential species of the subassociations are decisive for the classification of the association. According to the studies of the cited authors, the differential species of the compared subassociations are:

- *erythronietosum*: *Erythronium dens-canis*, *Lathyrus montanus*,
- *staphyletosum*: *Staphylea pinnata*, *Hacquetia epipactis*, *Craex digitata*, *Salvia glutinosa*, *Vicia oroboides*, *Aconitum vulparia*, *Rhamnus cathartica*;
- *caricetosum pilosae*: *Carex pilosa*, *Potentilla micrantha*, *Hepatica nobilis*;
- *asperuletosum*: *Galium odoratum*, *Asarum europaeum*, *Sanicula europaea*;
- *castanetosum*: *Castanea sativa*, *Melampyrum pratense*, *Gentiana asclepiadea*, *Pteridium aquilinum*, *Luzula luzuloides*, *Serratula tinctoria*;
- *luzuletosum*: *Luzula luzuloides*.

In the first statistical analysis the subassociations were distributed into six groups (subassociations), according to the standpoints of the phytocoenologists in the cited works. The goal of our analysis was to re-examine their standpoints and use a statistical method to determine differential species. Our subjective criterion guiding the selection was the following: participation in more than 30% of the relevés of a particular subassociation and the fidelity index above 30. The results of the analysis are given in Table I. Of 21 differential species determined by the original researchers, our analysis confirmed 14 and grouped as many as 37 new ones. Since the majority of the species had broader distribution, the previous phytocoenologists did not consider them sociologically significant for discriminating the subassociations. These are, for example, *Tilia tomentosa*, *Melica uniflora*, *Viola hirta*, *Hedera helix*, *Prenanthes purpurea*, *Fraxinus ornus*, *Senetio ovatus*, *Convallaria majalis* etc. On the other hand, a differential status was not confirmed for a smaller number of species (e.g. *Lathyrus montanus*, *Carex digitata*, *Hepatica nobilis*, *Serratula tinctoria*). In the primary research, these species had received this status on the basis of their higher participation, but in a small number of relevés.

In the second analysis we wanted to highlight some new possibilities of classifying the association *Epimedio-Carpinetum betuli* on the basis of the same phytocoenological relevés. Cluster analysis was employed to group the relevés into three clusters, which

were then used for further analysis.

Cluster I: comprises 36 relevés formed for the most part of the former groups 3 and 6, i.e. of the subassociations *erythronietosum* (13 relevés) and *staphyletosum* (21 relevés). The differential species of the new cluster include all the differential species of the former subassociation *staphyletosum* and the majority of the differential species of the former subassociations *erythronietosum* and *asperuletosum*. This cluster is the richest in the species of the Illyrian floral geoelement. A total of 11 species occur with over 30% in all the relevés in the cluster.

Cluster II: comprises 99 relevés, of which 90% belonged to the subassociation *caricetosum pilosae* in the previous analysis, and the rest mainly to the subassociation *asperuletosum*. The subassociation *caricetosum pilosae* had 7 differential species and they all regained their differential status. Along with these, the following species were also determined as differential: *Quercus cerris*, *Euphorbia amygdaloides*, *Lathyrus vernus*, *Lamium galeobdolon* and *Potentilla micrantha*. In the previous analysis, the first three did not prove to be differential species, *Potentilla micrantha* was found in the subassociation *castanetosum*, while *Lamium galeobdolon* was found in the subassociation *asperuletosum*.

Cluster III: comprises 111 phytocoenological relevés grouped from all the subassociations. This heterogeneous cluster contains all the relevés of the subassociations *castanetosum* and *luzuletosum*, while of other subassociations, the relevés lacking differential species determined in the two remaining clusters. It was also found that relevés with acidothermophilic species were more fully present, where *Castanea sativa* and *Luzula luzuloides* feature as differential species. Acidophytes *Melampyrum pratense*, *Hieracium racemosum*, *Hieracium murorum*, *Gentiana asclepiadea*, *Aposeris foetida* and other species have distinctly high participation. The backbone of this cluster is formed by the subassociations *castanetosum* and *luzuletosum*, and six species of the Illyrian floral geoelement are present in more than 30% of all the relevés.

According to the results of the analysis, the former subassociations *caricetosum pilosae* in Cluster II and *castanetosum* and *luzuletosum* in Cluster III manifest the highest degree of homogeneity.

The differential species of the subassociation *asperuletosum* are widely distributed and had a lower fi -



VUKELIĆ J., ŠAPIĆ I., BARIČEVIĆ D.  
 FLORISTIC-VEGETATIONAL VARIABILITY OF THE ASSOCIATION *EPIMEDIO-CARPINETUM BETULI* (HORVAT 1938)  
 BORHIDI 1963 IN THE NORTH OF CROATIA

**Table II:** Shortened synoptic table with percentage frequency and modified fidelity phi coefficient identified by cluster analysis

Group No. No. of relevés	1 36	2 99	3 111
Differential species			
<i>Heracleum sphondylium</i>	81 65.1	12 ---	16 ---
<i>Ranunculus lanuginosus</i>	67 64.4	1 ---	12 ---
<i>Euonymus europaeus</i>	83 60.7	27 ---	13 ---
<i>Milium effusum</i>	75 58.5	14 ---	17 ---
<i>Crocus vernus</i> agg.	58 58.4	1 ---	11 ---
<i>Campanula trachelium</i>	61 58.4	2 ---	13 ---
<i>Lamium orvala</i>	72 57.0	1 ---	29 ---
<i>Aegopodium podagraria</i>	61 55.5	3 ---	15 ---
<i>Melampyrum nemorosum</i>	58 52.4	3 ---	16 ---
<i>Staphylea pinnata</i>	50 52.1	2 ---	9 ---
<i>Scilla bifolia</i> agg.	39 52.0	2 ---	. ---
<i>Hacquetia epipactis</i>	53 50.8	. ---	15 ---
<i>Daphne mezereum</i>	64 49.2	1 ---	30 ---
<i>Helleborus atrorubens</i>	44 48.4	8 ---	2 ---
<i>Galanthus nivalis</i>	31 46.3	. ---	1 ---
<i>Ranunculus ficaria</i>	44 44.0	10 ---	5 ---
<i>Frangula alnus</i>	36 41.5	2 ---	7 ---
<i>Arum maculatum</i>	39 41.4	10 ---	2 ---
<i>Convallaria majalis</i>	58 41.1	6 ---	30 ---
<i>Paris quadrifolia</i>	31 39.9	4 ---	2 ---
<i>Glechoma hederacea</i>	36 37.3	8 ---	5 ---
<i>Malus sylvestris</i>	39 37.1	4 ---	13 ---
<i>Carex digitata</i>	36 36.7	5 ---	9 ---
<i>Mercurialis perennis</i>	33 35.8	4 ---	8 ---
<i>Ulmus glabra</i>	39 35.7	3 ---	15 ---
<i>Vicia oroboides</i>	36 35.0	6 ---	10 ---
<i>Prunus spinosa</i>	31 34.5	7 ---	4 ---
<i>Scrophularia nodosa</i>	44 33.5	9 ---	19 ---
<i>Moehringia trinervia</i>	33 33.5	10 ---	5 ---
<i>Geranium phaeum</i>	33 32.9	8 ---	7 ---
<i>Serratula tinctoria</i>	39 30.2	. ---	25 6.6
<i>Helleborus odoratus</i>	. ---	45 58.7	1 ---
<i>Rubus hirtus</i> s.lat.	6 ---	87 55.1	51 4.8
<i>Acer tataricum</i>	6 ---	52 55.1	4 ---
<i>Melica uniflora</i>	. ---	53 52.1	14 ---
<i>Viola hirta</i>	3 ---	46 51.8	5 ---
<i>Glechoma hirsuta</i>	3 ---	46 47.5	10 ---
<i>Carex pilosa</i>	17 ---	56 37.0	22 ---
<i>Quercus cerris</i>	6 ---	38 36.1	12 ---
<i>Tilia tomentosa</i>	3 ---	26 36.0	3 ---
<i>Galeobdolon luteum</i>	11 ---	47 35.6	18 ---
<i>Potentilla micrantha</i>	8 ---	49 33.9	26 ---
<i>Euphorbia amygdaloides</i>	. ---	30 30.6	14 ---
<i>Lathyrus vernus</i>	22 ---	60 30.4	34 ---
<i>Castanea sativa</i>	19 ---	2 ---	57 49.4
<i>Luzula luzuloides</i>	25 ---	6 ---	45 32.0
<i>Sanicula europaea</i>	47 2.2	23 ---	67 29.8
Dominant species			
<i>Carpinus betulus</i>	100 6.7	100 6.7	97 ---
<i>Quercus petraea</i>	100 15.1	96 1.1	91 ---
Constant species			
<i>Fagus sylvatica</i>	58 ---	89 17.0	90 19.1
<i>Galium odoratum</i>	58 ---	69 11.6	55 ---
<i>Carex sylvatica</i>	83 15.7	70 ---	68 ---
<i>Prunus avium</i>	94 26.3	79 ---	65 ---
<i>Hedera helix</i>	78 23.0	41 ---	67 6.9
<i>Asarum europaeum</i>	100 50.3	42 ---	57 ---
<i>Pulmonaria officinalis</i>	100 45.1	57 ---	57 ---
<i>Ligustrum vulgare</i>	92 42.8	57 ---	39 ---

<i>Galium sylvaticum</i>	89	41.9	37	---	53	---
<i>Corylus avellana</i>	100	39.0	43	---	86	16.5
<i>Cornus sanguinea</i>	94	38.3	64	---	50	---
<i>Acer campestre</i>	97	36.9	76	2.1	50	---
<i>Stellaria holostea</i>	89	33.2	64	---	48	---
<i>Crataegus monogyna</i>	86	31.3	57	---	52	---
<i>Viola reichenbachiana</i>	83	31.1	58	---	45	---
<b>Illyrian floral elemente</b>						
<i>Ruscus hypoglossum</i>	3	---	17	9.6	18	11.4
<i>Erythronium dens-canis</i>	14	4.0	6	---	16	9.0
<i>Epimedium alpinum</i>	50	---	71	22.8	43	---
<i>Knautia drymeia</i>	58	27.6	16	---	43	5.8
<i>Lonicera caprifolium</i>	86	54.0	9	---	49	1.0
<i>Cyclamen purpurascens</i>	56	34.6	8	---	34	2.4
<i>Primula vulgaris</i>	81	48.8	18	---	40	---
<i>Aposeris foetida</i>	100	60.4	4	---	69	16.6

delity index than all the differential species of other subassociations. As it turned out, this subassociation does not show individuality in northern Croatia. It was primarily established because the sampled plots lacked the favoured differential Illyrian species from other subassociations (*staphyletosum* and *erythronietosum* in the first place), while species of beech forests listed in Table I dominated. It should be mentioned that the association *Asperulo-Carpinetum* Wraber 1969 was described in Slovenia.

The subassociation *luzuletosum* follows a similar pattern. In the first analysis it had only two differential species (*Luzula luzuloides* and *Hieracium racemosum*) which occur in the new Cluster III with a high percentage of participation. Since chestnut was singled out as the differential species, the subassociation *castanetosum* should be given priority.

In relation to the previous insights, what is most surprising is the absence of the subassociation *erythronietosum*. Its 33 relevés were for the most part grouped into Cluster I and III. Of its differential species, *Erythronium dens-canis* and *Luzula pilosa* do not have diagnostic importance, whereas all the other species have differential importance for Cluster I, i.e. for the subassociation *staphyletosum*. Evidently, the diagnostic species from the previous studies could not accentuate the identity of the subassociation *erythronietosum* clearly enough to form its own cluster. However, it may have local importance, because the difference between the subassociations *staphyletosum* and *erythronietosum* was also confirmed by pedological research [9]. Undoubtedly, in the process of merging the majority of the relevés of these two subassociations into Cluster I, an important role was played by the biogeographic position of the sampled forest stands. Namely, the study area in the north-western Croatia is the richest in the species of the Illyrian floral geoelement, and so these relevés were grouped into Cluster I.

The former subassociation *staphyletosum* is represented by 41 relevés. Of this, 21 relevés was placed in Cluster I, which gathered all the differential species of the subassociation, including *Staphylea pinnata*. The remaining relevés were grouped into Cluster III. These relevés are characterized by a poorer floristic composition, lessened participation of Illyrian species, absence of diagnostic species of the subassociation *staphyletosum* and higher participation of the species *Castanea sativa*, *Solidago virgaurea*, *Sanicula europaea* and others. This can well be illustrated with a concrete

example of the percentage ratio of some particular species of the former subassociation *staphyletosum* after it was grouped into two clusters. The participation ratio of the species *Staphylea pinnata* in Cluster I in relation to Cluster III is 86:45 %, that of the species *Heracleum sphondylium* is 90:25 %, of *Ranunculus lanuginosus* it is 81:20 %, of *Hacquetia epipactis* it is 90:35 %, etc. All the relevés from the Samobor Mountains were grouped into Cluster III [29].

The subassociation *caricetosum pilosae* underwent the least changes in relation to the traditional standpoints. All the differential species remained the same in the new analysis as well, but some new ones were also added. The number of Illyrian species decreases from the west towards east, while some, such as *Lamium orvala* and *Hacquetia epipactis*, are completely absent. However, it is in this subassociation that important Illyrian species *Epimedium alpinum* and *Helleborus odoratus* are the best represented, despite being the only ones that occur in more than 30% of the relevés in Cluster II. Important differential species in the central and eastern distribution range of the community are *Tilia tomentosa*, *Heleborus odoratus*, *Acer tataricum* and *Quercus cerris*. It will be interesting to compare the relationship of these 99 relevés towards the related associations *Festuco drymeiae-Carpinetum* Vukelić ex Marinček 1994, and *Carici pilosae-Carpinetum betuli* Neuhäusl et Neuhäuslova-Novotna 1964. They are characterized by the smaller number and cover of Illyrian species and high participation of the species *Carex pilosa* and *Festuca drymeia*.

The analysis of ecological conditions conducted with Ellenberg's ecoindicator coefficients [7] did not show significant differences between individual clusters. According to these conditions, the association *Epimedio-Carpinetum betuli* occurs in semi-light (4.58 to 4.81), moderately warm (5.54 to 5.73) and fresh (4.81 to 5.04) habitats. They are moderately acidophilic to moderately basophilic (6.41 to 6.80) and moderately rich in nutrients (4.88 to 5.21).

By summarizing the problem of differential species in this manner, it is evident that they were determined primarily by means of numerical methods. In these methods all the species are put on the same level, regardless of their sociological affiliation. In this sense, it is possible to combine research methodologies for the purpose of achieving the best possible results. By analyzing as many relevés or syntaxa as possible, the fidelity value of more widely distributed ("common")

species is lessened, while the sociologically more closely affiliated species gain increasing importance. Moreover, an unbalanced number of relevés of a particular taxon may give a distorted image, which is also the problem of this analysis. This is the reason that we recommend our results primarily as guidelines for future research.

## 5 CONCLUSIONS

Based on the analysis of 246 relevés and participation of 276 species of higher plants of the association *Epimedio-Carpinetum betuli*, we re-examined its classification into traditional subassociations and proposed a possible new classification. The analysis is based first and foremost on differential species, for which of decisive importance was their participation and fidelity index in a particular syntaxon, i.e. cluster. A small number of differential species which were sociologically closely affiliated to a particular subassociation was frequently the reason for describing a larger number of subassociations, but this often did not correspond to the real field condition. In our study, the fund of differential species in traditionally formed subassociations was significantly increased, which facilitates field research and mapping.

In the second statistical analysis all the relevés were grouped into three clusters. In these clusters, the subassociations *caricetosum pilosae* and *staphyletosum* retained their basic identity, regardless of the fact that a part of the relevés was grouped into other clusters. The subassociation *asperuletosum* does not have distinct differential species and does not show an independent character. Its species are relatively abundantly present in other subassociations. The subassociation *luzuletosum* coincides with the subassociation *castanetosum* in terms of differential species and character, but the latter should be given priority. The relevés of the subassociation *erythronietosum* on deeper, more humid and shadier soils can be added to the subassociation *staphyletosum*, while those on steeper, drier and moderately acidophilic to neutrophilic soils with the species *Erythronium dens-canis* require more detailed study and analysis within the stands grouped in Cluster III.

The results of research suggest the need for further studies and a new syntaxonomic classification of the association *Epimedio-Carpinetum betuli* on the northern boundary of the Illyrian floral province. The results will also be useful for future research into the Illyrian oak-hornbeam forests in their overall distribution range, but also the *Carpinetum* communities in a wider European space.

## 6 REFERENCES

- [1] D. Aeschimann, K. Lauber, D. M. Moser, J. P. Theurillat, *Flora Alpina*, Bd 1. 2. 3. (2004), Haupt Verlag, Bern, Stuttgart, Wien.
- [2] D. Baričević, Sinakološko-fitocenološke značajke šumske vegetacije Požeške i Babjegore, Disertacija, Šumarski fakultet Sveučilišta u Zagrebu (2002), pag. 175, Zagreb.
- [3] A. Borhidi, Die Zönologie des Verbandes *Fagion illyricum*. 1. Allg. Teil. *Acta Bot. Acad. Sci. Hung.* 9 (1963), pag. 259–297.
- [4] A. Borhidi, Die geobotanischen Verhältnisse der Eichen- und Hainbuchenwälder Südosteuropas, *Feddes Rep.* 78/1-3 (1968), pag. 109–130.
- [5] J. Braun-Blanquet, *Pflanzensoziologie, Grundzüge der Vegetationskunde* (1964), pag. 865, Springer, Wien – New York.
- [6] K. Clarke, R. N. Gorley, *PRIMER v5. User manual/Tutorial*, Primer-E Ltd, (2001), Plymouth.
- [7] H. Ellenberg, C. Leuschner, *Vegetation Mitteleuropas mit den Alpen - Zeigerwerte der Pflanzen Mitteleuropas*, (2010), pag. 109, Eugen Ulmer, Stuttgart.
- [8] V. Glavač, Über Eichen-Hainbuchenwälder Kroatiens, *Feddes Rep.* 79/1-2 (1968), pag. 115–138.
- [9] M. Gračanin, Tipovi šumskih tala Hrvatske, *Glas. šum. pokuse* 9 (1948), pag. 95–119.
- [10] S. M. Hennekens, J. H. J. Schaminée, TURBOVEG, a comprehensive data base management system for vegetation data. *J. Veg. Sci.*, 12 (2001) pag. 589–591.
- [11] I. Horvat, Biljnoscijološka istraživanja šuma u Hrvatskoj. *Glas. šum. pokuse* 6 (1938), pag. 127–279.
- [12] I. Horvat, Laubwerfende Eichenzonen Südosteuropas in pflanzensoziologischer, klimatischer und bodenkundlicher Betrachtung, *Angew. Pflanzensoziologie* 15 (1958), pag. 50–62.
- [13] I. Horvat, Šumske zajednice Jugoslavije. *Šumarska enciklopedija*, I. izdanje, 2 (1963), pag. 560–590.
- [14] I. Horvat, V. Glavač, H. Ellenberg, *Vegetations Südosteuropas*. G. Fischer Verlag (1974), pag. 768, Stuttgart.
- [15] K. Hruška-Dell'Uomo, Biljni pokrov Moslavačke gore. Disertacija, Prirodoslovno-matematički fakultet Sveučilišta u Zagrebu (1974), pag. 312, Zagreb.
- [16] L. Marinček, Zur Nomenklatur der Hainbuchenwälder des *Erythronio-Carpinion*. Zbornik simpozija „Pevalek“, Flora i vegetacija Hrvatske, Šumarski fakultet Sveučilišta u Zagrebu i JP „Hrvatske šume“ (1994), pag. 57–62, Koprivnica – Zagreb.
- [17] L. Marinček, A. Čarni, Die Unterverbände der Hainbuchenwälder des Verbandes *Erythronio-Carpinion betuli* (Horvat 1938) Marinček in Walnöffler, Mucina et Grass 1993, *Scopolia* 45 (2000), pag. 1–20.
- [18] J. Medak, J. Fitocenološke značajke šuma pitomoga kestena u sjeverozapadnoj Hrvatskoj, Magistarski rad, Šumarski fakultet Sveučilišta u Zagrebu (2004), pag. 126, Zagreb.
- [19] T. Nikolić, (ur.), *Flora Croatica*, baza podataka. On-line (<http://hirc.botanic.hr/fcd>). Botanički zavod, Prirodoslovno-matematički fakultet, Sveučilište u Zagrebu, Accessed december, 2015.
- [20] J. Podani, SYN-TAX 2000. Computer Programs for Data Analysis in Ecology and Systematics, User's Manual (2001), pag. 53, Budapest.
- [21] Đ. Rauš, N. Šegulja, J. Topić, Vegetacija sjeveroistočne Hrvatske, *Glas. šum. pokuse* 23 (1985), pag. 225–355.
- [22] Lj. Regula-Bevilacqua, Biljni pokrov Strahinšćice u Hrvatskom zagorju, Disertacija, Prirodoslovno-matematički Sveučilišta u Zagrebu (1978), pag. 261, Zagreb.
- [23] R. Rizovski, Cenoze hrasta kitnjaka s običnim grabom i hrasta kitnjaka u centralnoj Makedoniji i centralnoj Hrvatskoj, Magistarski rad, Šumarski

- fakultet Sveučilišta u Zagrebu (1969), pag. 102, Zagreb.
- [24] R. Rizovski, Sporeduvanje na gorun-gaberovi šumi od severna Hrvatska i SR Makedonija, Šumarskiot fakultet na Univerzitet vo Skopje, Godišen zbornik na zemjodelsko 23 (1970), pag. 55–63, Skopje.
  - [25] J. S. Rodwell, H. J. Schamine, L. Mucina, S. Pignatti, J. Dring, D. Moss, The Diversity of Europaen Vegetation. An overview of phytosociological alliances and their relationships to EUNIS habitats, National Reference Centre for Agriculture, Nature and Fisheries, Wageningen, (2002), pag. 163, The Netherlands.
  - [26] I. Šapić, I., 2012: Šumska vegetacija Zrinske gore, Disertacija, Šumarski fakultet Sveučilišta u Zagrebu (2012), pag. 214, Zagreb.
  - [27] N. Šegulja, Biljni pokrov Vukomeričkih gorica. Disertacija, Prirodoslovno-matematički fakultet Sveučilišta u Zagrebu (2004), pag. 232, Zagreb.
  - [28] Ž. Škvorc, Florističke i vegetacijske značajke Dilja. Disertacija, Šumarski fakultet Sveučilišta u Zagrebu (2006), pag. 221, Zagreb.
  - [29] I. Šugar, Biljni svijet Samoborskog gorja. Disertacija, Sveučilište u Zagrebu, (2012), pag. 179, Zagreb.
  - [30] L. Tichý, JUICE, software for vegetation classification, Journal of Vegetation Science 13 (2002), pag. 451–453.
  - [31] J. Vukelić, Šumska staništa i zajednice hrasta kitnjaka (*Quercus petraea* Liebl.) u gorju sjeverozapadne Hrvatske, Glas. šum. pokuse 27 (1991), pag. 1–82.
  - [32] J. Vukelić, Šumska vegetacija Hrvatske, Sveučilište u Zagrebu, Šumarski fakultet, Državni zavod za zaštitu prirode (2012), pag. 403, Zagreb.
  - [33] S. Wallnöfer, L. Mucina, G. Grabherr, Die Pflanzengesellschaften Österreichs. Teil III, (1993), Gustav Fischer Verlag, pag. 85-236, Jena.
  - [34] M. Wraber, Über die Verbreitung, Ökologie und Systematische Gliederung der Eichen-Hainbuchenwälder Slowenien, Feddes Rep. 79/6, (1969), pag. 373–389.

## INSTRUCTIONS TO AUTHORS

Author(s)  
University(ies) / Institute(s)  
Address(es)

**ABSTRACT:** These notes provide important information on how to prepare and submit your paper. Read the notes carefully and follow them as precisely as possible. **Any inaccuracy will cause delay at the Technical Editors and in the publication of the Forest Review.** Your paper must be **written in English (UK)** and the layout should be exactly the same as this master document. **In order to prepare your layout, save this document with a new name and use it as a guide. Replace the text of this document with the text of your paper without changing the layout, font type and size, line spacing, page margins and structure of this template** (see section 3). **Do not insert page numbers or page headers/footers.** If you have any question, please do not hesitate to contact us (see section 7.3).

Keywords: select 3 to 6 keywords.

### 1 DEADLINES AND DELIVERY

**Papers will be published in the Forest Review only if correctly submitted.**

The electronic version of your paper must be submitted to the Technical Editors by e-mail according to the dates (see section 5, note 3), by one of the authors, together with **two suggested reviewers** (see section 11), and the **Copyright Transfer Agreement Form** duly filled-in and signed (this will be available and sent to the corresponding author before Review print).

Your original manuscript must be delivered in the following formats:

- **Microsoft Word** (97/2003, .docx), and **Adobe Acrobat PDF** by e-mail.

Please make sure that the paper you submit is the **final version** with all **numberings in the correct order**. **Do not submit the paper more than once.**

### 2 PREPARING THE MANUSCRIPT

#### 2.1 Volume and length of the paper

Please consider that the complete paper in pdf format, including illustrations, may not exceed **10 A4 pages**. This is a very good proven capacity for final papers.

#### 2.2 Organisation of the paper

The **title** of the paper should be informative and concise. It must be followed by the author(s) name(s) – listing the principal author first, organisation, complete address, telephone, fax and e-mail address.

No logos may appear in the title.

The **abstract** preceding the body of your paper should give a brief account of the most relevant aspects of your paper, in 200/250 words. Please avoid using symbols, graphics and text formatting (bold, italic, underline) in this part of the document.

Next, in order of importance, select three to six of the most relevant **keywords** and include them in your paper. The keywords should be separated with commas.

The **body** of the text must be in **two columns**. Number each heading using decimal numbering. Follow the layout specifications in section 3 below.

### 3 LAYOUT SPECIFICATIONS

**The layout of your paper should have exactly the same format as this master document.**

**Before** you start working on your paper, if you use Adobe Acrobat, select the printer option “Acrobat Distiller” (version 5.0) or Adobe pdf (version 6.0), in order to avoid accidental misplacement of layout elements afterwards when converting the Word file into PDF format.

#### 3.1 Font type and size

Font type: **Times New Roman**. Font size: **9pt**. Line spacing: **single**. Text alignment: **justified left and right**. Captions should have the same font and size as the typeface used for the text. Make sure that illustrations are clear and easy to read. Please do not use any other font than Times New Roman.

#### 3.2 Page size

Page size must be **A4** (210 mm x 297 mm). Margins: top: 32 mm; bottom: 19 mm; left and right: 25 mm.

The **body of the text must be in two equal columns** of 73,6 mm each. All written parts and images must fit **inside** these margins (for further details see subsection 4.1 about figures, and subsection 4.2 about tables).

#### 3.3 Typing the text

Begin at the top of the first page with the **title** of the paper in bold capital letters and centered.

Leave one blank line between the title and the name of the author(s).

List the surname preceded by the initial of the first name; when several authors prepare a paper, the name of the main one should appear first. On the following lines, give the name of the company or institute, wherever applicable, with the full address; the name of each organisation should be easy to depict. This paragraph must be centered and without any blank space.

Next, leave two blank lines and then type an **abstract** of no more than 250 words (keep the indent of this block on both sides, as shown on this document). At the end of the abstract give your 3/6 **keywords** on the last line.

Leave two blank lines between the abstract and the body of the text of your paper, which must be in two columns.

#### 3.4 Headings

Leave one blank line before each section and one blank line before the heading of each sub-section. Headings and sub-headings should be numbered (e.g. 3, 3.1, 3.2). Separate the numbers from the text of the

heading with two spaces.

There should be no blank line after the title of the sub-sections but only an indentation to indicate the beginning of a paragraph. Section headings should be in capital letters. Sub-section headings should be in upper and lower case. Headings should be normal text – not underlined or in bold.

## 4 ADDITIONAL COMPONENTS

### 4.1 Illustrations

Illustrations (photographs, drawings, graphs, charts, etc.) should not exceed 50% of the whole paper and should be placed as near as possible their citation. Illustrations must not be taken from previously-printed materials.

Illustrations should have a resolution of **300 dpi** using simple colors (Standard+RGB) and be placed at **100% scale** (i.e. if an illustration covers the full column width, it should be of approx. 860 pixel).


Illustrations should have the layout in line with the text (right click on the object – Format object – Layout – in line with the text, which should be the left-most option in the layout dialogue window).

All illustrations must be numbered progressively in bold decimals (e.g. **Figure 1:**) and have a reference in the text (e.g. Fig. 1). Captions should be as clear as possible, to allow comprehension of the illustration without reference to the text.


Graphs and charts must not be imported from Excel, but should be inserted as a picture (.jpg, .bmp or .gif). Please, use simple contrasting colors and effects instead of fill patterns. See Fig. 1 for good/bad example.

Illustrations must be clear also when printed in black and white.

Good example with contrasting colors:



Bad example with fill patterns:



**Figure 1:** Clear line drawings are essential

### 4.2 Tables

Tabular presentation of data is an easy way to condense many items. Tables must be numbered in bold Roman numerals (e.g. **Table I**), and have a reference in the text. Captions should be as clear as possible, for an easy comprehension of the tables.

**Table I:** Overview of biomass resources available

Biomass Sources	Quantity	Moisture	Residue
Sewage Sludge	1.86	1.73	1.40
Septage	0.32	0.28	0.16
Fruit Pulp	3.78	3.89	4.02

### 4.3 References and notes

References and notes must not appear as footnotes in the pages, but should be listed together at the end of the text, in the dedicated sections.

When referring to them in the text, please type the corresponding reference number in brackets. Use round brackets for the notes (1) and square brackets for the references [1].

To make them easier to find, indent your notes and references from the second line, as in the examples (see sections 5 and 6).

### 4.4 Acknowledgements

Any acknowledgement should be added before the references and/or notes, in a dedicated section, as in the example (see section 9).

## 5 NOTES

- (1) This section should have the progressive number before the title, exactly as for the previous ones.
- (2) Do not add any unnecessary space between the listed numbers of your notes.
- (3) Important dates of the Forest Review are:
  - Abstract submission March 15<sup>th</sup>
  - Abstract acceptance March 30<sup>th</sup>
  - Paper submission May 15<sup>th</sup>
  - Paper acceptance June 15<sup>th</sup>
  - Forest Review Online/Print October 15<sup>th</sup>

## 6 REFERENCES

- [1] This section should have the progressive number before the title, exactly as for the previous ones.
- [2] Do not add any unnecessary space between the listed numbers of your references.
- [3] G. Campolmi, Proceedings of the 3<sup>rd</sup> World Biomass Conference – Biomass for Energy, Industry and Climate Protection, III Vol. (2005), pag. 981.
- [4] D. Reed, Evaluation of Biomass Resources in the southern regions in Nigeria, (2007), pag. 124.
- [5] O. Vecchi, Biofuel Production in central Italy, (2008), pag. 45.

## 7 OTHER POINTS

### 7.1 Permissions

Authors are fully responsible for their manuscripts. They must take the necessary steps to obtain permission for using any material that might be protected by copyright.



## 7.2 Copyright

Please be aware that on delivery of your manuscript, you transfer the **copyright** to **Ss. Cyril and Methodius University in Skopje, Faculty of Forestry in Skopje**, publishers of the Forest Review.

## 7.3 Further information

We are available to assist you. Please, do not hesitate to contact us for any query; please address your e-mails to **sumpregled@sf.ukim.edu.mk**.

## 8 IMPORTANT SPECIFICATIONS FOR THE PDF

### 8.1 Printer

If you are using Adobe Acrobat, please select Acrobat Distiller (version 5.0) or Adobe Pdf (version 6.0) as printer option to convert your Word file into pdf format. Go to Printer Properties and click on “Acrobat pdf settings”; in Conversion Settings select “Press” quality.

If you are using **special fonts** make sure that the checkbox “Do not send fonts to distiller (“adobe pdf” for version 6.0)” is **not** selected.

### 8.2 Password / Security

Please **do not use any Password / Security when making the Pdf file**. As the header and page numbers will be added by us, we need to ensure that your pdf file is not password protected.

## 9 ACKNOWLEDGEMENTS

- We thank you for successfully following these instructions. This will make the Reviewers easier to read and avoid queues at the Technical Editors!
- The Authors are grateful to the students-members of DREN - NGO of students of Forestry at the Faculty of Forestry in Skopje for their helpful cooperation.
- .....

## 10 LOGO SPACE

If you wish, you may add your logos (e.g. Organisations, Project Partners, Supporters, Brands etc) at the end of the paper.

## 11 IMPORTANT NOTICE FOR THE SUBMISSION

To ensure a convenient publication process, the authors are requested to follow these instructions. Submission of a paper implies that it reports unpublished work and that it is not under consideration for publication elsewhere. If previously published tables, illustrations or more than 200 words of text are to be included, then the copyright holder's written permission must be obtained. Copies of any such permission letters should be scanned and attached with the paper.

A **condition of submission** (with the abstract submission) is that the author nominates **two people** who are qualified to act **as reviewers** and who have not previously been involved with the paper in any way. Both of the nominated reviewers must be working in a different country to the first named author.

Before the publication, the paper will be reviewed by the members of the Editorial Board and a notification of

acceptance will be send to the corresponding author. Editors may request that authors submit a revised version of their paper before it can be accepted for publication. The Editor's decision on all submissions is final.

Finally, the authors will receive **Copyright Transfer Agreement Form** which they need to fill-in and sign for the publication to be complete.





## ШУМАРСКИ ПРЕГЛЕД

Меѓународно научно списание  
Год. 46 / Стр. 1-95  
Скопје, 2015

## FOREST REVIEW

International Scientific Journal  
Vol. 46 / Pag. 1-95  
Skopje, 2015

Online ISSN 1857-9507  
УДК 630  
УДК 635.9  
УДК 674

Online ISSN 1857-9507  
UDC 630  
UDC 635.9  
UDC 674

### Издавач

Универзитет „Св. Кирил и Методиј“ во Скопје  
Шумарски факултет во Скопје  
Декан  
Д-р Јане Ацевски

### Publisher

Ss. Cyril and Methodius University in Skopje  
Faculty of Forestry in Skopje  
Dean  
Jane Acevski PhD

### Главен и одговорен уредник

Д-р Бојан Симовски

### Editor-in-chief

Bojan Simovski PhD

### Уредувачки одбор

Д-р Љупчо Несторовски (Скопје, Македонија)  
Д-р Марилена Ицџотиќ (Загреб, Хрватска)  
Д-р Милосав Анђелиќ (Подгорица, Црна Гора)  
Д-р Милорад Даниловиќ (Белград, Србија)  
Д-р Ирена Папазова-Анакиева (Скопје, Македонија)  
Д-р Роберт Брус (Љубљана, Словенија)  
Д-р Чиприан Палагиану (Сучава, Романија)  
Д-р Влатко Андоновски (Скопје, Македонија)  
Д-р Саша Орловиќ (Нови Сад, Србија)  
Д-р Маргарита Георгиева (Софија, Бугарија)  
Д-р Зоран Говедар (Бања Лука, Р. Српска, БИХ)  
Д-р Јасминка Ризовска Атанасовска (Скопје, Македонија)  
Д-р Мариус Димитров (Софија, Бугарија)  
М-р Дејан Манџуковски (Скопје, Македонија)

### Editorial board

Ljupčo Nestorovski PhD (Skopje, Macedonia)  
Marilena Idžojtić PhD (Zagreb, Croatia)  
Milosav Anđelić PhD (Podgorica, Montenegro)  
Milorad Danilović PhD (Belgrade, Serbia)  
Irena Papazova-Anakieva PhD (Skopje, Macedonia)  
Robert Brus PhD (Ljubljana, Slovenia)  
Ciprian Palaghianu PhD (Suceava, Romania)  
Vlatko Andonovski PhD (Skopje, Macedonia)  
Saša Orlović (Novi Sad, Serbia)  
Margarita Georgieva PhD (Sofia, Bulgaria)  
Zoran Govedar PhD (Banja Luka, R. Srpska, BiH)  
Jasminka Rizovska Atanasovska PhD (Skopje, Macedonia)  
Marius Dimitrov PhD (Sofia, Bulgaria)  
Dejan Mandžukovski MSc (Skopje, Macedonia)

### Технички уредник

М-р Дејан Манџуковски

### Technical editor

Dejan Mandžukovski MSc

### Корица и насловна фотографија

Д-р Бојан Симовски, *Ramonda nathaliae*

### Cover page and photography

Bojan Simovski PhD, *Ramonda nathaliae*

Излегува еднаш годишно

Published once a year

### Интернет-страница

[www.sf.ukim.edu.mk/sumarski\\_pregled.htm](http://www.sf.ukim.edu.mk/sumarski_pregled.htm)

### Web page (on-line)

[www.sf.ukim.edu.mk/sumarski\\_pregled.htm](http://www.sf.ukim.edu.mk/sumarski_pregled.htm)

### Адреса на издавачот

УКИМ-Шумарски факултет во Скопје  
Редакција на Шумарски преглед  
Ул. „16 Македонска бригада“ бр. 1  
(П. факс 235)  
1 000 Скопје  
Република Македонија  
Е-пошта: [sumpregled@sf.ukim.edu.mk](mailto:sumpregled@sf.ukim.edu.mk)  
[www.sf.ukim.edu.mk](http://www.sf.ukim.edu.mk)

### Publisher's address

UKIM Faculty of Forestry in Skopje  
Editorial Board of the Forest Review  
Ul. 16 Makedonska brigada br. 1  
(P.O. box 235)  
MK-1000 Skopje  
Republic of Macedonia  
E-mail: [sumpregled@sf.ukim.edu.mk](mailto:sumpregled@sf.ukim.edu.mk)  
[www.sf.ukim.edu.mk](http://www.sf.ukim.edu.mk)

Шум. преглед (Šum. pregled)  
For. review

Год. 46  
Vol. 46

Стр. 1-95  
Pag. 1-95

Скопје, 2015  
Skopje, 2015

